	DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
			ANTIBIOTICS		
PENICILLIN	**Amoxicillin**	Oral	*Neonates:* 10-15mg/kg BD *Paediatrics:* 15-30 mg/kg TDS	1g	Acute OM/severe pneumonia: dose 90mg/kg/**DAY.** May prescribe 20-45mg/kg BD for compliance
	Amoxicillin/ clavulanic acid (Augmentin)	Oral	Suspension: Augmentin 228mg/5mL (amoxicillin **200mg**, clauvunate 28mg per 5mL): 25mg/kg/dose BD-TDS Augmentin ES 642.9mg/5mL (amoxicillin **600mg**, clauvunate 42.9mg per 5mL): 48mg/kg/dose BD	12.5mL 15mL (package insert)	Dose is based on **Amox/Clav component** and it depends on the **ratio formulation of Amoxicillin/Clavulanate.** Max **clavulanic** dose **10mg/kg/day** (500mg/day)
			Tablets (>12yo): Augmentin 625mg tablet (amoxicillin **500mg**, clauvunate 125mg) – 1 tablet BD-TDS Augmentin 1g Tablet (amoxicillin **875mg**, clauvunate 125mg) – 1 tablet BD	625mg 1g	
	Cloxacillin	Oral	6.25 – 12.5mg/kg QDS	500mg	
CEPHALOSPORIN	**Cefaclor**	Oral	10-15mg/kg TDS	500mg	Potential for serum sickness
	Cefuroxime	Oral	Suspension: 10-15mg/kg BD Tablets (>12yo): 250-500mg BD	500mg	Suspension and tablets not interchangeable.
	Cephalexin	Oral	10-15mg/kg QDS Severe: 15-25mg/kg QDS	500mg	Split total daily dose to 3 divided doses for compliance

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

	DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
MACROLIDE	**Azithromycin**	Oral	10mg/kg 24H for 3 days Alternative: 10mg/kg on Day 1, then 5mg/kg on Days 2-5	500mg	For age > 28 days
	Clarithromycin	Oral	7.5mg/kg BD	500mg	
QUINO-LONE	**Ciprofloxacin**	Oral	10-20mg/kg BD	750mg	Not routinely first-line therapy but can be considered a reasonable alternative for some situations after assessment of risks and benefits.
SULPHO-NAMIDE	**Trimethoprim/ sulfamethoxazole**	Oral	Treatment: 4-6mg **TMP/kg** Q12	160mg	**Doses expressed in Trimethoprim (TMP) component** **Use in children >/= 2mo**
LINCOSAMIDE	**Clindamycin**	Oral	Mild: 5-7mg/kg TDS Mod/severe: 7-13mg/kg q8H	600mg	May cause pseudomembraneous colitis
ANTIVIRALS					
Acyclovir		Oral	VZV (within 24 hrs of rash): 20mg/kg QDS for 5 days HSV gingivostomatitis: 20mg/kg QDS	800mg	
Oseltamivir (Tamiflu)		Oral	I Influenza Treatment: **Infants ≤ 8mo:** 3mg/kg BD **Infant ≥9mo:** 3-3.5mg/kg BD **Children and Adolescent:** ≤15kg: 30mg BD >15-23kg: 45mg BD >23-40kg: 60mg BD >40kg: 75mg BD	75mg	BD dosing for treatment; OD dosing for prophylaxis Treatment duration: 5 days ***Needs dose adjustment in renal-impairment**

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
ANTIHELMINTIC				
Albendazole	PO	Pinworm/Hookworm/Round worm/Whipworm infection: 1-2yo: 200mg single dose (premeal) >2yo: 400mg single dose (premeal) Max: 400mg single dose	400mg single dose	May use up to 3-5 consecutive days depending on indication
ANTIPYRETICS / ANALGESICS				
Paracetamol	Oral	10-15mg/kg 6H	1g	Use higher end of dose range for analgesia. Max dose 75mg/kg/day Caution in hepatic failure.
	Rectal	10-20mg/kg 6H		
Codeine/Paracetamol	Oral	1 tab Q6H	16mg/ 1000mg	**Use in children >/= 12yo** 1 tablet contains Codeine 8mg/Paracetamol 500mg
Orphenadrine/ Paracetamol (Anarex)	Oral	450mg/35mg Q8H	900mg/ 70mg	**Use in children >/= 12yo** 1 tablet contains Orphenadrine 35mg/Paracetamol 450mg
buprofen	Oral	5-10mg/kg Q6-8H	400mg 1.2g/day	**Use in children >/= 6 months** Max dose 40mg/kg/day *Use lower range for anti-pyretic effect, and higher range for anti-inflammatory effect *Caution in renal impairment, dehydration, asthma. Risk of bleeding
Diclofenac	Oral/ Rectal	0.5-1mg/kg 8-12H	50mg	**Use in children >/= 1yo** *Caution in renal impairment
Mefenamic Acid	Oral	Dysmenorrhea: 500mg once, then 250mg Q6H	500mg	**Use in children >/= 12yo**
Naproxen	Oral	Musculoskeletal pain or dysmenorrhea: 5mg/kg BD	500mg	**Use in children >/= 16yo** Dosage expressed as **naproxen base.** 250mg Naproxen base = 275mg Naproxen sodium
Etoricoxib	Oral	60-120mg once daily	120mg	**Use in children >/= 16yo**
Tramadol	Oral	1-2mg/kg 6H PRN	100mg	**Avoid use in < 12yo** Do not exceed 50mg per dose for <50kg; max daily dose 400mg ***causes nausea**

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
CONSTIPATION MEDICATIONS				
Glycerin	Rectal	1-6 mo: 1g (0.5 supp) >6mo to 2yo: 2g (1 supp) >2yo to 6yo: 3g (1.5 supp)		Each supp = 2g
Lactulose	Oral	Laxative: 0.5ml/kg 12-24H Hepatic coma: 1ml/kg 1H, then 6H		
Bisacodyl	Oral	3-12yo: 5-10mg OD >12yo: 10mg OD	10mg	**Tablets not recommended for use <3yo.** Tab: 5mg (Cannot be crushed)
	Rectal	2-12yo: 5-10mg OD >12yo: 10mg OD		
Sennoside	Oral	2-6yo: 0.5 to 1 tablet ON 6-12yo: 1-2 tablets ON	15mg	Each tablet = 7.5mg
Fleet® Enema	Rectal	≥ 2 – 12yo: 1 **Paeds enema** ≥12yo: 1 bottle **Adult enema**		**Not recommended for use < 2 yo** **Paeds enema – 2.3oz (66.6mL)** **Adult enema – 4.5oz (133mL)** Not recommended for use < 2yo Caution in renal impairment (PO4 content).
Forlax	Oral	**Constipation:** For adults/children ≥8 years old: 1-2 sachets a day		**Not to be used in <2yo** Off label use in 2-7yo Macrogol 4000 – 10g per sachet Each sachet should be dissolved in at least 120mL of water.

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
ANTI-DIARRHOEAL MEDICATIONS				
Charcoal	Oral	250mg-500mg TDS/QDS	2000mg	Nil recommendation on age limit. Suggest giving to only older children.
Lacteol fort sachet	Oral	≤2yo: ½ sachet BD >2yo: 1 sachet BD		**Avoid in immunocompromised patients**
Smecta	Oral	>2yo: 1 sachet BD- TDS		Not recommended in ≤2yo due to lead content
Racecadotril (Hidrasec)	Oral	<9kg: 10mg TDS 9-13kg: 20mg TDS 13-27kg: 30mg TDS >27kg: 60mg TDS		**Not to be used in <3 mo** Avoid using >7 days Sachet size: 10mg, 30mg
GASTRITIS MEDICATIONS				
1. Gaviscon® OR **2. MMT**	Oral	6-12yo: 5-10ml TDS prn >12yo: 10-20ml TDS prn		Give after meals/bedtime. SE: diarrhea
Famotidine	Oral	<3mo: 0.5-1mg/kg OD ≥3mo: 0.5-1mg/kg BD	40mg	
Omeprazole	Oral	0.5-1mg/kg/dose OD-BD	40mg	
ABDOMINAL PAIN MEDICATION				
Ridwind®	Oral	Infants and children < 2yo or <11 kg: 0.2ml QDS 2 – 12yo (> 11kg): 0.4ml QDS		Each 1ml = 100mg **Simethicone** Apply to feeding bottle or directly to mouth Max 2.4ml/day
Trimebutine (Debridat®)	Oral	**Tablets** 6mo – 1yo: 25mg BD 1 – 5yo: 25mg TDS 5yo-<12yo: 50mg TDS ≥12yo: 100-200mg TDS	200mg TDS (max)	Off label use 100mg tab (dose equivalent, rationalize to nearest quarter tab)

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
ANTI-EMETICS				
Cinnarizine	Oral	0.3-0.6mg/kg TDS	12.5mg	**Not to be used in <5yo** Tablet = 25mg
Domperidone	Oral	250mcg/kg TDS	10mg	Max 1 week duration
Promethazine	Oral	0.25-0.5mg/kg 6-8H	10-25mg	**Contraindicated in <2yo**
Ondansetron	SL/ Oral	0.1-0.2mg/kg 8H	8mg	**For use in children > 6months. *Only to be given in clinic setting**
GENITOURINARY TRACT MEDICATIONS				
Ural Effervescent Granules	Oral	6-12yo: 1 sachet up to TDS ≥12yo: 1-2 sachets up to QDS	2 sachets	Each 4g sachet contains sodium bicarbonate 1.76g, sodium citrate anhydrous 0.63g, citric acid anhydrous 0.72g and tartaric acid 0.89g. Contains total 650mg sodium per sachet
MUCOLYTICS / COUGH SUPPRESSANTS				
Acetylcysteine	Oral	>2yo-<6yo: 1 sachet BD 6yo-12yo: 2 sachet BD >12yo: 1 tab BD	600mg/ dose	1 sachet = 100mg N-Acetylcysteine 1 effervescent tab = 600mg **Avoid using in children </= 2yo**
Ambroxol	Oral	2-5yo: 2.5mL TDS 6-12yo: 5mL BD-TDS >12 yo: 10mL BD	60mg	**Use in children >/= 2yo** 5mL contains ambroxol 30mg
Promethazine	Oral	2-5yo: 2.5-5mg BD 5-12yo: 5-10mg BD >12yo: 10-15mg BD	25mg	Soln 5mg/5ml **Contraindicated in <2yo** **Avoid concomitant antihistamines**
Rhinathiol Promethazine	Oral	2-5yo: 2.5mL-5mL QDS 6-12yo: 7.5-12.5mL TDS 12-15yo: 12.5-15mL TDS >15yo: 15-20mL TDS	20mL	**Use in children >/= 2yo** Contains alcohol (3%) 5mL contains carbocisteine 100mg and promethazine HCl 2.5mg

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
ANTIHISTAMINES / DECONGESTANTS				
Chlorpheniramine	Oral	0.1mg/kg 6-8H	4 mg	**Avoid use of ≥ 1 antihistamine**
Cetirizine (Zyrtec)	Oral	6mo-2yo: 2.5mg OD 2-5yo: 2.5-5mg OD >5yo: 5-10mg OD	10 mg	**Avoid use of ≥ 1 antihistamine** Solution: 1mg/ml Tablet: 10mg
Loratadine	Oral	2-5yo: 5mg OD ≥6yo: 10mg OD	10mg	**Avoid use of ≥ 1 antihistamine** Solution: 1mg/ml
Fexofenadine (Telfast)	Oral	6mo - <2yo & 10.5kg: 15mg BD 2-11yo: 30mg BD ≥12yo: 60mg BD OR 180mg OD	180mg/ day	Suspension: 30mg/5mL Tab: 180mg
Telfast D® (pseudoephedrine/ fexofenadine)	Oral	≥12yo: 1 *tab* BD	1 tab	**Use in children >/= 12yo** 1 *tab*: pseudoephedrine 120mg/ Fexofenadine 60mg
Hydroxyzine (Atarax)	Oral	0.5mg/kg/dose TDS - QDS <6yo: max 12.5mg/dose >6yo: max 25mg/dose	25mg	**Use in children ≥6 mo** **Avoid use of ≥ 1 antihistamine**
Oxymetazoline (Iliadin)	Nasal	***<1mo: use Normal Saline drops instead** 1mo-1yo: 0.01% 1 drop TDS 1-6yo: 0.025% 1 drop TDS >6yo: 0.05% 1 drop TDS		**Duration ≤ 5 days** to avoid rebound congestion
Xylometazoline (Otrivin)	Nasal	***<2yo: use Normal Saline drops instead** 2-6yo: 0.05% 1 drop BD >6yo: 0.05% 2 drop BD-TDS		**Duration ≤ 5 days** to avoid rebound congestion

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
ASTHMA MEDICATIONS				
Salbutamol	MDI	2-6 puffs 4-6H	10 puffs	Via spacer and mask Each puff delivers 100mcg
	Neb	<5yo: 0.5ml 0.5% salbutamol: 2.5ml 0.9% NaCl ≥5yo: 1ml 0.5% salbutamol : 2ml 0.9% NaCl	1ml	**Minimum salbutamol volume 0.5mL**
Ipratropium Bromide	MDI	<6yo: 2-4 puffs 6-8H ≥6yo: 4-8 puffs 6-8H		
	Neb	1ml with salbutamol and/or 0.9% NaCl 6-8 hourly	1ml	Nebuliser solution 500mcg/2mL (0.025%)
Prednisolone	Oral	1-2mg/kg OD for 3-5 days	40mg	No need taper dose if <14 days
Montelukast	Oral	6mo-<6yo: 4mg once daily 6yo-<15yo: 5mg once daily ≥15yo: 10mg once daily	10mg	Granule: To consume within 15 minutes after opening
Hydrocortisone	IV	Asthma: 4mg/kg 6H	200mg	
CROUP / ANAPHYLAXIS MEDICATIONS				
Dexamethasone	Oral	Mild: 0.15mg/kg/dose Moderate – Severe: 0.6mg/kg/dose	10mg	High dose 0.6mg/kg used as single dose (can be given IV/IM/PO) Up to 3 doses MAX, 12hrs apart
Budesonide	Neb	2mg as a single dose, OR 1mg for 2 doses separated by 30min interval	2mg	May repeat every 12H until clinical improvement
Adrenaline	Neb	0.5ml/kg of 1:1000 (1mg/ml, 0.1%) Repeat after 30min if necessary	4ml = 4mg	**Severe croup; watch for rebound**. Dilute with NaCl 0.9% or give neat. Effects: 2-3hrs

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	Adult dose	REMARKS
Adrenaline/Epinephrine	SC/IM	0.01ml/kg of 1:1000 (1mg/mL, 0.1%)	0.5ml	***NOT to be given intravenously**
	IM	Epipen: 7.5kg-25kg: Epipen Jr 0.15mg >25kg: Epipen 0.3mg	0.3mg	Scan QR code for instructional video
SEIZURE MEDICATIONS				
Lorazepam	IV	0.05-0.1mg/kg Repeat after 10min if seizing	4 mg	Up to max 2 doses of benzodiazepines (rectal/IV)
Diazepam	Rectal	<10kg: 2.5mg >10kg-20kg: 5mg >20kg: 10mg	20mg	One rectal tube: 5mg/2.5mL

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

Topical Medications

DRUG	ROUTE	PAEDIATRIC DOSE	REMARKS
EAR CONDITIONS: OTITIS MEDIA			
Ciprofloxacin 2mg, Hydrocortisone 10mg/mL Ear Drop	Topical	≥ 1yo: 3 drops BD	Duration: 7 days Consider using 1 drop instead in smaller children with smaller ear canal volume
NOSE CONDITIONS: ALLERGIC RHINITIS			
Fluticasone Furoate Nasal Spray (Avamys)	Intranasal	1-2 spray each nostril once a day	**Use in children >/= 2yo** Each puff delivers 27.5mcg
Mometasone Nasal Spray	Intranasal	1-2 spray each nostril once a day	**Use in children >/= 2yo** Each puff delivers 50mcg May increase to BD in acute rhinosinusitis in adults
EYE CONDITIONS: INFECTIVE CONJUNCTIVITIS			
Tobramycin 0.3% (Tobrex Eyedrop)	Topical	1 drop BD-QDS Severe infections 1 drop 4H (Caution about dose-related conjunctival toxicity)	**Max 2 drops every hour in severe infection** until improvement *Can be used in neonates, but practice is based on labelled indications
Chloramphenicol 0.5% Eyedrop	Topical	1 drop TDS-QDS and continue for 48 hours after healing	**Up to 1 drop 2-hourly in severe infections** **Suitable for infants**
Fusidic Acid 1% Eyedrop (Fucithalmic)	Topical	1 drop BD	
Chlortetracycline 1% eye ointment	Topical	1 application TDS	Apply thin strip (approximately 1 cm) of ointment to the conjunctiva
EYE CONDITIONS: ALLERGIC CONJUNCTIVITIS			
Olopatadine HCl 0.1% (Patanol Eyedrop)	Topical	≥ 3yo: 1 drop BD-QDS	
REGULAR DIAPER RASH/ DERMATITIS			
Zinc Oxide 40% (Desitin **ointment)**	Topical	1 application PRN	

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	REMARKS
FUNGAL DIAPER RASH			
Miconazole nitrate 2% cream	Topical	1 application BD or 1 application daily when with miconazole powder	Duration: 2-6 weeks. **Continue for 1 week after symptoms disappear**
Daktacort cream (Hydrocortisone 1%, Miconazole nitrate 2%)	Topical	1 application once to twice daily	**Steroid and antifungal combination.** May use for 3-4 days until redness improves then switch to anti-fungal monotherapy
TOPICAL ANALGESIA			
Diclofenac sodium 1% Gel	Topical	Apply TDS-QDS	**Use in children >/= 1yo**
Ketoprofen 30mg plaster (Kefentech)	Topical	Apply 1 plaster BD	Avoid pasting patch on the same area to prevent skin irritation
MOISTURIZERS			
QV cream	Topical	Apply liberally PRN	Humectant/moisturizer
Emulsifying ointment	Topical	Apply liberally PRN	Emollient
ECZEMA / ATOPIC DERMATITIS			
Hydrocortisone 1% cream	Topical	Apply thinly on affected areas daily to BD	**For mild eczema** **0.5% strength available for babies/ young infants**
Potency of steroid creams			
Mild (Class VI, VII)	Hydrocortisone 0.5% cream, Hydrocortisone 1% cream, Betamethasone valerate 0.05% cream, Desonide 0.05% lotion, Desonide cream 0.05% (Desowen), Betamethasone valerate 0.025% cream.		
Moderate (Class IV, V)	Mometasone furoate 0.1% **cream/lotion**, Hydrocortisone aceponate 0.127% cream, Betamethasone valerate 0.1% **cream (≥ 1yo only)**, Fluticasone propionate 0.05% **cream**		
Potent (Class II, III)	Mometasone furoate 0.1% **ointment**, Betamethasone dipropionate 0.05% **cream/ointment**		
Very potent (Class I)	Clobetasol propionate 0.05% cream/ointment **(≥ 1yo only)**		
ASTRINGENT WASH			
Calamine lotion	Topical	Apply to affected areas OD-QDS	**Anti-pruritic,** May reapply after bath

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	REMARKS
ORAL ULCER (NON-INFECTIVE)/SORE THROAT			
Soragel (Cetalkonium Cl 0.01%, choline salicylate 8.714%)	Topical	1 application PRN	**Use in infants >/= 4 months** (As stated on the packaging) May be used for teething
Oracort E (Triamcinolone Acetonide 0.1%, Lignocaine HCl 3%)	Dental paste	1 application TDS PRN	Generally avoided in young children Contains Lidocaine HCl 3%, Triamcinolone acetonide 0.1%
Benzydamine (Difflam)	Oral lozenge	1 loz q1-2h, max 12 loz/day	**Use in children >/= 6yo** One lozenge contains Benzydamine 3mg (sugar-free)
Dequalinium	Oral lozenge	1 loz q2-3h, max 8 loz/day	**Use in children >/= 12yo** 1 lozenge contains dequalinium 0.25mg
ORAL ULCER (INFECTIVE)			
Acyclovir 5% cream	Topical	Apply 5 times a day	**For herpes labialis** Duration: 5-10 days
ORAL CARE / MUCOSITIS			
Chlorhexidine gluconate 0.2% mouthwash	Oral	5-10mL BD-TDS PRN	For **secondary infection** of mucosal ulceration and as adjunct to **control gingivitis**. **Prevention of oral candidiasis** in immunocompromised patients.
Nystatin suspension	Oral	<1mo: 1ml QDS 1-12mo: 2mL QDS; apply cotton/finger ≥12mo: 5mL TDS-QDS	**Oral thrush/ candidiasis** 100,000 units/mL Duration: 7 days & continue treatment for at least 2 days after symptoms disappear May swish and swallow
SCABIES			
Malathion 0.5% lotion	Topical	≥ 6mo: Apply over whole body (including scalp, neck, face and ears). Wash off after 24hrs. Repeat application after 7 days	Reapply if hands are washed with soap within 24hrs
Permethrin 5% Lotion	Topical	≥ 2mo: Apply over whole body including face, neck, scalp and ears; wash off after 8–14hrs. Repeat application after 7 days	Reapply if hands washed with soap within 8hrs of application

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

DRUG	ROUTE	PAEDIATRIC DOSE	REMARKS
HEAD LICE			
Malathion 0.5% lotion	Topical	≥ 6yo: Apply sufficient amount on dry hair to thoroughly wet the hair and scalp and allow the hair to dry naturally. Wash after 8-12hrs. Repeat second application after 7-9 days if needed	Use a fine-toothed (nit) comb to remove dead lice and eggs.
FIRST DEGREE/ SUPERFICIAL BURNS			
Chlorhexidine gluconate 1% cream	Topical	Clean affected area and apply generously.	**Antiseptic**
SUPERFICIAL ABRASIONS AND WOUNDS			
Chlortetracycline 1% eye ointment	Topical	1 application PRN	**For facial region**
Tetracycline 3% ointment	Topical	1 application PRN	**Avoid application to facial region for patients <4yo**
Fusidic acid 2%	Topical	1 application BD to TDS	**For infected abrasions** Duration: 7 days
Fucidic acid 2%, Betamethasone val 0.1% cream	Topical	1 application BD	**For infected abrasions** ≥1 yo only
Mupirocin 2% Ointment	Topical	1 application TDS	**For infected abrasions**
Clotrimazole 1%, Beclomethasone dipropionate 0.025%, Gentamicin sulfate 0.1%	Topical	1 application BD-TDS	**For infected abrasions** ≥1 yo only Contains Betamethasone dipropionate 0.025% which is a potent steroid
Octenisan™ Wash Lotion	Topical	Apply as body wash during showering or wiped over body during sponge bath x 5 days	
Potassium permanganate solution (1:1000)	Topical	1 application PRN Dilute 1:8 parts with water	May be used to compress or wash affected areas. Dab dry after use, not necessary to wash off.

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.

Appendix

Suitable quantities for prescribing

Suitable quantities of dermatological preparations to be prescribed for specific areas of the body		
Area of body	Creams and Ointments	Lotions
Face	15-30 g	100 ml
Both hands	25-50 g	200 ml
Scalp	50-100 g	200 ml
Both arms or both legs	100-200 g	200 ml
Trunk	400 g	500 ml
Groins and genitalia	15-25 g	100 ml
These amounts are usually suitable for children 12-18 years for twice daily application for 1 week; smaller quantities will be required for children under 12 years. These recommendations **do not apply** to corticosteroid preparations.		

Source: BNF for children 2021-2022

Disclaimer: Every effort has been made to ensure the accuracy of the information in this guide. The authors will not be responsible for any errors arising or related to its use.