

Mahvish Samar and Xinzhong Zhu*

*Multiplicative perturbation analysis for the
generalized Cholesky block downdating problem*

JOURNAL OF MATHEMATICAL INEQUALITIES

AIMS AND SCOPE

Journal of Mathematical Inequalities (**JMI**, J. Math. Inequal.) presents carefully selected original research articles from all areas of pure and applied mathematics, provided they are concerned with mathematical inequalities. **JMI** will also periodically publish invited survey articles with interesting results treating the theory of inequalities, as well as relevant book reviews.

JMI is published quarterly, in March, June, September, and December.

SUBMISSION OF MANUSCRIPTS

Authors are requested to submit their articles electronically as $\text{T}_\text{E}\text{X}$ or $\text{L}_\text{A}\text{T}_\text{E}\text{X}$ files, with enclosed Adobe Acrobat PDF format through the provided web interface on journal web page.

The author who submitted the article for publication will be denoted as a corresponding author. He/She manages all communication and correspondence with the **JMI** regarding the article, makes any revisions, and reviews and releases the author proofs. Authors may indicate a member of the Editorial Board whom they consider appropriate for the article. However, assignment to that particular editor is not assured.

COPYRIGHT

The acceptance of the article automatically implies the copyright transfer to the **JMI**. Manuscripts are accepted for review with the understanding that the same work has not been published (except in the form of an abstract), that it is not under consideration for publication elsewhere, that it will not be submitted to another journal while under review for the **JMI**, and that its submission for publication has been approved by all of the authors.

OPEN ACCESS

Journal of Mathematical Inequalities is published as open access journal. All interested readers are allowed to view, download, print, and redistribute any article without paying for subscription. Offprints of each article may be ordered from **JMI** prior to publication.

PREPARATION OF MANUSCRIPTS

Manuscripts should be written in English, using prepared journal LaTeX style.

The first page should contain the article title, authors' names (complete first names and family names), complete affiliations (name of institution, address, city, state, and zip code), e-mail addresses, proposed running head (not more than 40 characters), a short abstract (not more than 150 words), a list of key words and phrases, and the AMS 2010 Mathematics Subject Classification primary (and secondary) codes. Avoid abbreviations, mathematical symbols and formulas, diagrams, and reference to the text of the article.

Figures should be prepared in a digital form suitable for direct reproduction, at resolution of 300 dpi or higher, and in EPS, TIFF or JPEG format.

Bibliographic references should be listed alphabetically at the end of the article, each numbered by an Arabic number between square brackets. The following information should be provided for references to journals: names of authors, full title of the article, abbreviated name of the journal, volume, year of publication, and range of page numbers. For standard abbreviations of journal names, the authors should consult the latest Abbreviations of Names and Series reviewed in Mathematical Reviews.

SOURCE FILE, PROOFS

Upon acceptance of the article, the authors will be asked to send the related LaTeX source file to the Editorial Office, in accordance to the journal style. In order to accelerate the publication process, the AMS LaTeX package is strongly preferred. PDF proofs will be sent by e-mail to the corresponding author.

FORTHCOMING PAPERS

Papers accepted and prepared for publication will appear in the forthcoming section of Journal Web page. They are identical in form as final printed papers, except volume, issue and page numbers.

JMI is published by Publishing House **ELEMENT**, Zagreb, Croatia.

All correspondence and subscription orders should be addressed to the Editorial Office:

www.ele-math.com
e-mail: jmi@ele-math.com

Journal of Mathematical Inequalities
Editorial Office
Menceticeva 2, 10000 Zagreb, Croatia
Fax: +385 1 6008799

MULTIPLICATIVE PERTURBATION ANALYSIS FOR THE GENERALIZED CHOLESKY BLOCK DOWNDATING PROBLEM

MAHVISH SAMAR* AND XINZHONG ZHU

(Communicated by T. Burić)

Abstract. This article is devoted to the multiplicative perturbation analysis of the generalized Cholesky block downdating problem. The strong rigorous multiplicative perturbation bounds are first presented by bringing together the modified matrix-vector equation approach with the technique of Lyapunov majorant function and the Banach fixed point theorem. Then, the weak rigorous multiplicative bounds are developed by using the matrix-equation approach. Numerical results demonstrate that these bounds are constantly tighter than the additive perturbation bounds.

1. Introduction

Suppose that $A \in \mathbb{R}_m^{m \times m}$ is symmetric positive definite, $B \in \mathbb{R}_n^{n \times m}$, and $C \in \mathbb{R}^{n \times n}$ are symmetric positive and semi-definite, then the symmetric quasi-definite matrix $K \in \mathbb{R}^{(m+n) \times (m+n)}$ can be expressed as

$$K = \begin{bmatrix} A & B^T \\ B & -C \end{bmatrix}. \quad (1.1)$$

The matrix K always has the generalized Cholesky factorization

$$K = LJ_{m+n}L^T, \quad (1.2)$$

where

$$L = \begin{bmatrix} L_{11} & 0 \\ L_{21} & L_{22} \end{bmatrix}, \quad J_{m+n} = \begin{bmatrix} I_m & 0 \\ 0 & -I_n \end{bmatrix},$$

$L_{11} \in \mathbb{R}_m^{m \times m}$, $L_{22} \in \mathbb{R}_n^{n \times n}$ are lower triangular and $L_{21} \in \mathbb{R}_n^{n \times m}$. From (1.2), it can be simply verify that

$$A = L_{11}L_{11}^T, \quad B = L_{21}L_{11}^T, \quad C + L_{21}L_{21}^T = L_{22}L_{22}^T.$$

Mathematics subject classification (2020): 15A45, 15A23.

Keywords and phrases: Generalized Cholesky block downdating problem, multiplicative perturbation, rigorous perturbation bounds, Lyapunov majorant function, Banach fixed point theorem, matrix-equation approach.

This work was supported by the Zhejiang Normal University Postdoctoral Research Fund (Grant No. ZC304022938), the Natural Science Foundation of China (Project No. 61976196) and the Zhejiang Provincial Natural Science Foundation of China under Grant No. LZ22F030003.

* Corresponding author.

If the diagonal elements of the lower triangular matrices, L_{11} and L_{22} are positive, the factorization (1.2) is unique and L is known as the generalized Cholesky factor [1].

In this paper, we consider the generalized Cholesky block downdating problem (GCBD)

$$LJ_{m+n}L^T - YY^T = VJ_{m+n}V^T. \quad (1.3)$$

Given that K is the same as in (1.1) and $Y = (Y_m^T, Y_n^T) \in \mathbb{R}^{m+n \times k}$, find a lower triangular matrix

$$V = \begin{bmatrix} V_{11} & 0 \\ V_{21} & V_{22} \end{bmatrix} \in \mathbb{R}^{(m+n) \times (m+n)},$$

where $V_{11} \in \mathbb{R}_+^{m \times m}$, $V_{22} \in \mathbb{R}_+^{n \times n}$ are lower triangular with positive elements and $V_{21} \in \mathbb{R}_+^{n \times m}$. From [2, Corollary 1], it is simply to show that when $\|L^{-1}Y\|_2 < 1$ the GCBD problem (1.3) is always exists and the matrix V is known as the GCBD factor. Moreover, in this case

$$A - Y_m Y_m^T = V_{11} V_{11}^T, \quad B - Y_n Y_n^T = V_{21} V_{11}^T, \quad C + V_{21} V_{21}^T + Y_n Y_n^T = V_{22} V_{22}^T.$$

This problem is reduced to the Cholesky block downdating problem if we choose $K = A$, i.e., B and C are nonexistent. The Cholesky block downdating problem has acquired remarkable consideration, and its special case, i.e., single downdating ($Y \in \mathbb{R}^{m \times 1}$), has been extensively studied in the literature; see [3, 4, 5, 6, 7, 8, 9, 10] for details. Additive perturbation and multiplicative perturbation are the two distinctive types of perturbation models of matrix factorizations. The additive perturbation analysis for the GCBD problem has been considered in [11, 12].

Obviously, an additive perturbation can be obtained from the multiplicative perturbation, but the derived additive perturbation bounds will destroy the unique structures of multiplicative perturbations and lose their tendency [13]. Since the matrix scaling technique is often employed to provide better-conditioned problems [13], the multiplicative perturbation has received significant attention, and has some elements of interest compared with the additive perturbation; see [13, 14, 15, 16, 17] and references therein. Until now, there has been no work on the multiplicative perturbation bounds for the GCBD problem. So, it is interesting to introduce the multiplicative perturbation bounds for the GCBD problem.

Particularly, for (1.3), we assume two types of multiplicative perturbation matrices, $W = I_{m+n} + N$ on L : (1) $W = I_{m+n} + N$ is a general matrix; (2) $W = I_{m+n} + N$ is a lower triangular matrix. The rest of the paper is organized as follows: In Section 3, we will develop the strong rigorous multiplicative perturbation bounds by using the modified matrix-vector equation approach [13, 18, 19, 20], the Lyapunov majorant function (e.g., [11, Chapter 5]), and the Banach fixed point theorem (e.g., [11, Appendix 5]). Moreover, we will use the matrix-equation approach [21] to derive the weak rigorous multiplicative bounds in Section 4. These bounds will be less expensive to compute as compared to the strong, rigorous multiplicative perturbation bounds. Section 2 provides some useful notation and preliminary knowledge. Finally, we provide some numerical experiments to verify the theoretical results, and we show the numerical comparison between the multiplicative rigorous perturbation bounds and the additive rigorous perturbation bounds [11, 12] for the GCBD problem in Section 5.

2. Preliminaries

Some notations can be endorsed in [21] to make the presentation apparent. We still illustrate them here to make easier for readers.

The Frobenius norm and spectral norm for a given matrix $Z = (z_{ij}) \in \mathbb{R}^{m \times n}$ are denoted by $\|Z\|_F$ and $\|Z\|_2$, respectively. The following inequalities hold for these two matrix norms; see [22] for details.

$$\|QRS\|_2 \leq \|Q\|_2 \|R\|_2 \|S\|_2, \quad \|QRS\|_F \leq \|Q\|_2 \|R\|_F \|S\|_2, \quad (2.1)$$

whenever the matrix product QRS is well-defined.

For any matrix $Z = [z_1, z_2, \dots, z_n] = (z_{ij}) \in \mathbb{R}^{n \times n}$, denote the vector of the last i elements of z_j by $z_j^{(i)}$ and define

$$\text{lvec}(Z) := \begin{bmatrix} z_1^{(n)} \\ z_2^{(n-1)} \\ \vdots \\ z_n^{(1)} \end{bmatrix} \in \mathbb{R}^{v_1}, \quad \text{vec}(Z) := \begin{bmatrix} z_1 \\ z_2 \\ \vdots \\ z_n \end{bmatrix} \in \mathbb{R}^{n^2}, \quad \text{slt}(Z) := \begin{bmatrix} 0 & 0 & 0 & \cdots & 0 \\ z_{21} & 0 & 0 & \cdots & 0 \\ z_{31} & z_{32} & 0 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ z_{n1} & z_{n2} & \cdots & z_{n,n-1} & 0 \end{bmatrix},$$

$$\text{low}(Z) := \begin{bmatrix} \frac{1}{2}z_{11} & 0 & \cdots & 0 \\ z_{21} & \frac{1}{2}z_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ z_{n1} & z_{n2} & \cdots & \frac{1}{2}z_{nn} \end{bmatrix}, \quad \text{lt}(Z) := \begin{bmatrix} z_{11} & 0 & \cdots & 0 \\ z_{21} & z_{22} & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ z_{n1} & z_{n2} & \cdots & z_{nn} \end{bmatrix},$$

$\text{sut}(Z) = \text{slt}(Z^T)^T$, and $\text{diag}(Z) = \text{diag}(z_{11}, z_{22}, \dots, z_{nn})$, where $v_1 = \frac{n(n+1)}{2}$. Using the structures of these operators, we have

$$\text{lvec}(Z) = \mathfrak{K}_{\text{lvec}} \text{vec}(Z), \quad \text{vec}(\text{lt}(Z)) = \mathfrak{K}_{\text{lt}} \text{vec}(Z), \quad \text{vec}(\text{low}(Z)) = \mathfrak{K}_{\text{low}} \text{vec}(Z), \quad (2.2)$$

where

$$\begin{aligned} \mathfrak{K}_{\text{lvec}} &= \text{diag}(G_1, G_2, \dots, G_n) \in \mathbb{R}^{v_1 \times n^2}, \quad G_i = [0_{n-(i-1) \times (i-1)}, I_{n-(i-1)}] \in \mathbb{R}^{n-(i-1) \times n}, \\ \mathfrak{K}_{\text{lt}} &= \text{diag}(\hat{G}_1, \hat{G}_2, \dots, \hat{G}_n) \in \mathbb{R}^{n^2 \times n^2}, \quad \hat{G}_i = \text{diag}(0_{(i-1) \times (i-1)}, I_{n-(i-1)}) \in \mathbb{R}^{n \times n}, \\ \mathfrak{K}_{\text{low}} &= \text{diag}(\tilde{G}_1, \tilde{G}_2, \dots, \tilde{G}_n) \in \mathbb{R}^{n^2 \times n^2}, \quad \tilde{G}_i = \text{diag}(0_{(i-1) \times (i-1)}, 1/2, I_{n-i}) \in \mathbb{R}^{n \times n}. \end{aligned}$$

Moreover,

$$\mathfrak{K}_{\text{lvec}} \mathfrak{K}_{\text{lvec}}^T = I_{v_1}, \quad \mathfrak{K}_{\text{lvec}}^T \mathfrak{K}_{\text{lvec}} = \mathfrak{K}_{\text{lt}}. \quad (2.3)$$

Let $\text{lvec}^\dagger : \mathbb{R}^{v_1} \rightarrow \mathbb{R}^{n \times n}$ be the right inverse of the operator 'lvec' such that $\text{lvec} \cdot \text{lvec}^\dagger = 1_{v_1 \times v_1}$ and $\text{lvec}^\dagger \cdot \text{lvec} = \text{lt}$. Then the matrix of the operator 'lvec' is $\mathfrak{K}_{\text{lvec}}^T$. That is, $\text{lvec}^\dagger(Z) = \mathfrak{K}_{\text{lvec}}^T \text{vec}(Z)$.

Let $\mathbb{D}_n \in \mathbb{R}^{n \times n}$ be the set of diagonal matrices with positive diagonal elements. Then, for any $D_n = \text{diag}(\sigma_1, \sigma_2, \dots, \sigma_n) \in \mathbb{D}_n$, it follows that

$$\text{low}(ZD_n) = \text{low}(Z)D_n, \quad \text{low}(D_nZ) = D_n\text{low}(Z). \quad (2.4)$$

Moreover, from [23, Lemma 5.1], we have

$$\|\text{low}(Z) + D_n\text{low}(Z^T)D_n^{-1}\|_F \leq \sqrt{1 + \varsigma_{D_n}^2} \|Z\|_F, \quad (2.5)$$

where $\varsigma_{D_n} = \max_{1 \leq i < j \leq n} \{\sigma_j / \sigma_i\}$. From [21], we have

$$\|\text{low}(Z)\|_F \leq \|Z\|_F. \quad (2.6)$$

If Z is symmetric, then

$$\|\text{low}(Z)\|_F \leq \frac{1}{\sqrt{2}} \|Z\|_F. \quad (2.7)$$

Also, we have

$$\|\text{low}(Z + Z^T)\|_F \leq \sqrt{2} \|Z\|_F. \quad (2.8)$$

The Kronecker product between $Z = (Z_{ij}) \in \mathbb{R}^{m \times n}$ and $X \in \mathbb{R}^{p \times q}$ is defined as $Z \otimes X = [z_{ij}X] \in \mathbb{R}^{mp \times nq}$. Some useful results of the Kronecker product are listed below [24]:

$$\text{vec}(ZCX) = (X^T \otimes Z)\text{vec}(C), \quad (2.9)$$

$$\Pi \text{vec}(Z) = \text{vec}(Z^T), \quad (2.10)$$

$$\|X \otimes Z\|_2 = \|X\|_2 \|Z\|_2, \quad (2.11)$$

$$(X \otimes Z)(B \otimes C) = (XB \otimes ZC), \quad (2.12)$$

$$(X \otimes Z)^{-1} = X^{-1} \otimes Z^{-1}, \text{ if } X \text{ and } Z \text{ are nonsingular}, \quad (2.13)$$

where B and C are of suitable orders.

3. Strong rigorous multiplicative perturbation bounds

Let us consider that the matrices L , Y and V in (1.3) are perturbed as

$$L \rightarrow LW, \quad Y \rightarrow YU, \quad V \rightarrow \Delta V,$$

where $W = I_{m+n} + N \in \mathbb{R}^{(m+n) \times (m+n)}$, $U = I_{m+n} + M \in \mathbb{R}^{(m+n) \times (m+n)}$ and $\Delta V \in \mathbb{R}^{(m+n) \times (m+n)}$ is a lower triangular matrix with positive diagonal elements. Therefore, the perturbed form of (1.3) is

$$\begin{aligned} (V + \Delta V)J_{m+n}(V + \Delta V)^T &= (I_{m+n} + N)LJ_{m+n}L^T(I_{m+n} + N)^T \\ &\quad - (I_{m+n} + M)YY^T(I_{m+n} + M)^T. \end{aligned} \quad (3.1)$$

Extending (3.1) and using (1.3), we have

$$VJ_{m+n}(\Delta V)^T + (\Delta V)J_{m+n}V^T = N(LJ_{m+n}L^T) + (LJ_{m+n}L^T)N^T - M(Y Y^T) - (Y Y^T)M^T + N(LJ_{m+n}L^T)N^T - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T. \quad (3.2)$$

Premultiplying (3.2) by V^{-1} and right postmultiplying it by V^{-T} lead to

$$\begin{aligned} & J_{m+n}(\Delta V)^T V^{-T} + V^{-1}(\Delta V)J_{m+n} \\ &= V^{-1}(N(LJ_{m+n}L^T) + (LJ_{m+n}L^T)N^T)V^{-T} - V^{-1}(M(Y Y^T) + (Y Y^T)M^T)V^{-T} \\ & \quad + V^{-1}(N(LJ_{m+n}L^T)N^T - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)V^{-T}. \end{aligned} \quad (3.3)$$

As performed in [8, 9, 11, 14], from (3.3), we have

$$\begin{aligned} V^{-1}\Delta VJ_{m+n} &= \text{low}(V^{-1}(N(LJ_{m+n}L^T) + (LJ_{m+n}L^T)N^T)V^{-T}) \\ & \quad - \text{low}(V^{-1}(M(Y Y^T) + (Y Y^T)M^T)V^{-T}) \\ & \quad + \text{low}(V^{-1}(N(LJ_{m+n}L^T)N^T - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)V^{-T}). \end{aligned} \quad (3.4)$$

Applying the operator ‘vec’ to (3.4), and noting (2.2), (2.9) and (2.10), we get

$$\begin{aligned} & (J_{m+n}^T \otimes V^{-1}) \text{vec}(\Delta V) \\ &= \mathfrak{K}_{\text{low}}((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T))\Pi) \text{vec}(N) \\ & \quad - \mathfrak{K}_{\text{low}}((V^{-1}(Y Y^T)^T \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(Y Y^T))\Pi) \text{vec}(M) \\ & \quad + \mathfrak{K}_{\text{low}}(V^{-1} \otimes V^{-1}) \text{vec}(N(LJ_{m+n}L^T)N^T - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T). \end{aligned}$$

As performed in [8, 9, 11, 14], we can obtain

$$\begin{aligned} \text{vec}(\Delta V) &= (J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) \\ & \quad + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T))\Pi) \text{vec}(N) \\ & \quad - (J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(Y Y^T)^T \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(Y Y^T))\Pi) \text{vec}(M) \\ & \quad + (J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}(V^{-1} \otimes V^{-1}) \\ & \quad \times \text{vec}(N(LJ_{m+n}L^T)N^T - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T), \end{aligned} \quad (3.5)$$

and show that (3.5) is equivalent to

$$\begin{aligned} \text{lvec}(\Delta V) &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) \\ & \quad + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T))\Pi) \text{vec}(N) \\ & \quad - \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(Y Y^T)^T \otimes V^{-1}) \\ & \quad + (V^{-1} \otimes V^{-1}(Y Y^T))\Pi) \text{vec}(M) \\ & \quad + \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}(V^{-1} \otimes V^{-1}) \text{vec}(N(LJ_{m+n}L^T)N^T \\ & \quad - M(Y Y^T)M^T - \Delta VJ_{m+n}(\Delta V)^T). \end{aligned} \quad (3.6)$$

As a matter of convenience, suppose

$$\begin{aligned} Q_{LM} &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T)) \Pi) \\ Q_Y &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(YY^T)^T \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(YY^T)) \Pi) \\ R_V &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}(V^{-1} \otimes V^{-1}). \end{aligned} \quad (3.7)$$

Then, (3.6) becomes

$$\begin{aligned} \text{lvec}(\Delta V) &= (Q_{LM} \text{vec}(N) - Q_Y \text{vec}(M) \\ &\quad + R_V \text{vec}(N(LJ_{m+n}L^T)N^T - M(YY^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)). \end{aligned} \quad (3.8)$$

Thus, applying the operator ‘lvec[†]’ to (3.8) yields

$$\begin{aligned} \Delta V &= \text{lvec}^{\dagger}(Q_{LM} \text{vec}(N) - Q_Y \text{vec}(M) \\ &\quad + R_V \text{vec}(N(LJ_{m+n}L^T)N^T - M(YY^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)). \end{aligned} \quad (3.9)$$

We will derive the rigorous multiplicative perturbation bound for ΔV using the method of the Lyapunov majorant function and the Banach fixed point theorem based on the operator equation (3.10) as shown in [8, 9, 11, 14]. The equation (3.9) can be written as an operator equation for ΔV :

$$\begin{aligned} \Delta V &= \Phi(\Delta V, N, M) \\ &= \text{lvec}^{\dagger}(Q_{LM} \text{vec}(N) - Q_Y \text{vec}(M) \\ &\quad + R_V \text{vec}(N(LJ_{m+n}L^T)N^T - M(YY^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)). \end{aligned} \quad (3.10)$$

Suppose that $H \in \mathbb{R}^{(m+n) \times (m+n)}$ is lower triangular with positive diagonal elements and has the same structure as that of ΔV , $\|H\|_F \leq \eta$ for some $\eta \geq 0$, $\|N\|_F = \sigma_1$ and $\|M\|_F = \sigma_2$. Then it follows from the definition of the operator ‘lvec[†]’ and (2.1) that

$$\|\Phi(H, N, M)\|_F \leq \|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2 + \eta^2). \quad (3.11)$$

Using (3.11), we get the Lyapunov majorant function of the operator equation (3.10)

$$q(\eta, \sigma_1, \sigma_2) = \|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2 + \eta^2)$$

and the Lyapunov majorant equation

$$\begin{aligned} q(\eta, \sigma_1, \sigma_2) &= \eta, \quad \text{i.e.} \\ \|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2 + \eta^2) &= \eta. \end{aligned} \quad (3.12)$$

Suppose that σ_1 ,

$$\begin{aligned} \sigma_2 \in \Omega &= \{ \sigma_1, \sigma_2 \geq 0 : 1 - 4\|R_V\|_2 (\|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 \\ &\quad + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2)) \geq 0 \}. \end{aligned}$$

Then, the equation (3.12) has two nonnegative roots: $\eta_1(\sigma_1, \sigma_2) \leq \eta_2(\sigma_1, \sigma_2)$ with

$$\eta_1(\sigma_1, \sigma_2) = f(\sigma_1, \sigma_2) = \frac{2(\|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2))}{1 + \sqrt{1 - 4\|R_V\|_2 (\|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2))}}.$$

Let the set $A(\sigma_1, \sigma_2)$ be $A(\sigma_1, \sigma_2) = \{H \in \mathbb{R}^{(m+n) \times (m+n)} : H \text{ has the same structure as that of } \Delta V \text{ and } \|H\|_F \leq \eta\}$, which is closed and convex. Furthermore, we can simply verify that the operator $\Phi(\cdot, N, M)$ maps the set $A(\sigma_1, \sigma_2)$ into itself for $H, \tilde{H} \in A(\sigma_1, \sigma_2)$,

$$\|\Phi(H, N, M) - \Phi(\tilde{H}, N, M)\|_F \leq q'_\eta(f(\sigma_1, \sigma_2), \sigma_1, \sigma_2) \|H - \tilde{H}\|_F.$$

Since the derivative of the function $q(\eta, \sigma_1, \sigma_2)$ relative to η at $f(\sigma_1, \sigma_2)$ satisfies

$$q'_\eta(f(\sigma_1, \sigma_2), \sigma_1, \sigma_2) = 1 - \sqrt{1 - 4\|R_V\|_2 (\|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2))} < 1,$$

when $\sigma_1, \sigma_2 \in \Omega_1 = \{\sigma_1, \sigma_2 \geq 0 : 1 - 4\|R_V\|_2 (\|Q_{LM}\|_2 \sigma_1 + \|Q_Y\|_2 \sigma_2 + \|R_V\|_2 (\|L\|_2^2 \sigma_1^2 + \|Y\|_2^2 \sigma_2^2)) > 0\}$. Then the operator $\Phi(\cdot, N, M)$ is contractive on the set $A(\sigma_1, \sigma_2)$ for $\sigma_1, \sigma_2 \in \Omega_1$. Thus, from the Banach fixed point theorem, we have that the operator Equation (3.10), i.e. the matrix equation (3.2), has a unique solution in the set $A(\sigma_1, \sigma_2)$. As a result, $\|\Delta V\|_F \leq f(\sigma_1, \sigma_2)$ for $\sigma_1, \sigma_2 \in \Omega_1$. We summarize these results in the following theorem.

THEOREM 3.1. *Given a lower triangular matrix $L \in \mathbb{R}^{(m+n) \times (m+n)}$ with positive diagonal elements and a matrix $Y \in \mathbb{R}^{(m+n) \times k}$ such that the generalized Cholesky factorization $VJ_{m+n}V^T = LJ_{m+n}L^T - YY^T$ holds. Suppose $W = I_{m+n} + N \in \mathbb{R}^{(m+n) \times (m+n)}$ and $U = I_{m+n} + M \in \mathbb{R}^{(m+n) \times (m+n)}$. If*

$$\|R_V\|_2 (\|Q_{LM}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2)) < \frac{1}{4}, \quad (3.13)$$

then the following generalized Cholesky factorization holds:

$$\begin{aligned} & (V + \Delta V)J_{m+n}(V + \Delta V)^T \\ &= LW(LW)^T - YU(YU)^T, \\ &= (I_{m+n} + N)LJ_{m+n}L^T(I_{m+n} + N)^T - (I_{m+n} + M)YY^T(I_{m+n} + M)^T, \end{aligned}$$

and

$$\begin{aligned} & \|\Delta V\|_F \\ & \leq \frac{2(\|Q_{LM}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2))}{1 + \sqrt{1 - 4\|R_V\|_2 (\|Q_{LM}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2))}} \end{aligned} \quad (3.14)$$

$$\leq 2(\|Q_{LM}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2)). \quad (3.15)$$

Proof. It is absolutely not difficult to check that the condition (3.13) is the same as the one in Ω_1 . So, (3.14) and thus (3.15) hold. \square

REMARK 3.2. The resulting first order multiplicative perturbation bound can be obtained from (3.15) by neglecting high-order terms

$$\begin{aligned}\|\Delta V\|_F &\leq \|Q_{LM}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F \\ &= \|Q_{LM}\|_2 \|W - I_{m+n}\|_F + \|Q_Y\|_2 \|U - I_{m+n}\|_F.\end{aligned}\quad (3.16)$$

We used $W = I_{m+n} + N$ as a general matrix in the previous analysis. Next, we choose $W = I_{m+n} + N$ as a lower triangular matrix. Thus, from (3.10), (2.2) and (2.3), it follows that

$$\begin{aligned}\text{lvec}(\Delta V) &= (Q_{LT} \text{lvec}(N) - Q_Y \text{vec}(\Delta Y) \\ &\quad + R_V \text{vec}(N(LJ_{m+n}L^T)N^T - M(YY^T)M^T - \Delta VJ_{m+n}(\Delta V)^T)).\end{aligned}\quad (3.17)$$

where

$$\begin{aligned}Q_{LT} &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) \\ &\quad + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T)) \Pi) \mathfrak{K}_{\text{lvec}}^T.\end{aligned}\quad (3.18)$$

Based on the results of Theorem 3.1, we have the following theorem.

THEOREM 3.3. *Given a lower triangular matrix $L \in \mathbb{R}^{(m+n) \times (m+n)}$ with positive diagonal elements and a matrix $Y \in \mathbb{R}^{(m+n) \times k}$ such that the generalized Cholesky factorization $VJ_{m+n}V^T = LJ_{m+n}L^T - YY^T$ holds. Suppose $W = I_{m+n} + N \in \mathbb{R}^{(m+n) \times (m+n)}$ and $U = I_{m+n} + M \in \mathbb{R}^{(m+n) \times (m+n)}$. If*

$$\|R_V\|_2 (\|Q_{LT}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2)) < \frac{1}{4}, \quad (3.19)$$

then the following generalized Cholesky factorization holds:

$$\begin{aligned}(V + \Delta V)J_{m+n}(V + \Delta V)^T \\ &= LW(LW)^T - YU(YU)^T, \\ &= (I_{m+n} + N)LJ_{m+n}L^T(I_{m+n} + N)^T - (I_{m+n} + M)YY^T(I_{m+n} + M)^T,\end{aligned}$$

and

$$\begin{aligned}\|\Delta V\|_F \\ &\leq \frac{2(\|Q_{LT}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2))}{1 + \sqrt{1 - 4\|R_V\|_2 (\|Q_{LT}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2))}},\end{aligned}\quad (3.20)$$

$$\leq 2(\|Q_{LT}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F + \|R_V\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2)). \quad (3.21)$$

REMARK 3.4. By ignoring the high-order terms, we can investigate a first-order multiplicative perturbation bound from (3.21):

$$\begin{aligned}\|\Delta V\|_F &\leq \|Q_{LT}\|_2 \|N\|_F + \|Q_Y\|_2 \|M\|_F \\ &= \|Q_{LT}\|_2 \|W - I_{m+n}\|_F + \|Q_Y\|_2 \|U - I_{m+n}\|_F.\end{aligned}\quad (3.22)$$

4. Weak rigorous multiplicative perturbation bounds

To obtain the explicit expression of weak rigorous multiplicative perturbation bounds for the GCBD problem, we will adopt the matrix-equation approach originated by Chang [21] in this section.

THEOREM 4.1. *Given a lower triangular matrix $L \in \mathbb{R}^{(m+n) \times (m+n)}$ with positive diagonal elements and a matrix $Y \in \mathbb{R}^{(m+n) \times k}$ such that the generalized Cholesky factorization $VJ_{m+n}V^T = LJ_{m+n}L^T - YY^T$ holds. Suppose $S = (S_m^T, S_n^T)^T \in \mathbb{R}^{(m+n) \times k}$ and $H \in \mathbb{R}^{(m+n) \times (m \times n)}$ be lower triangular of the following form*

$$H = \begin{bmatrix} H_{11} & 0 \\ H_{21} & H_{22} \end{bmatrix},$$

where $H_{11} \in \mathbb{R}^{m \times m}$ and $H_{22} \in \mathbb{R}^{n \times n}$ are lower triangular, $H_{21} \in \mathbb{R}^{n \times m}$. Define $N = \varepsilon H$ and $M = \varepsilon S$ for some $\varepsilon \geq 0$. If

$$\|N\|_2 < 1, \quad \frac{\|L^{-1}\|_2 (\|Y\|_2 + \|YM\|_2)}{1 - \|N\|_2} < 1, \quad (4.1)$$

then $(L(I_{m+n} + N))J_{m+n}(L(I_{m+n} + N))^T - (Y(I_{m+n} + M))(Y(I_{m+n} + M))^T$ has the generalized Cholesky factorization

$$\begin{aligned}(V + \Delta V)J_{m+n}(V + \Delta V)^T \\ = (I_{m+n} + N)LJ_{m+n}L^T(I_{m+n} + N)^T - (I_{m+n} + M)YY^T(I_{m+n} + M)^T.\end{aligned}\quad (4.2)$$

Proof. For any $|t| \leq \varepsilon$ from the first condition of (4.1), it follows that

$$\|tH\|_2 \leq \|\varepsilon H\|_2 = \|N\|_2 < 1. \quad (4.3)$$

Then $L(I_{m+n} + tH)$ is nonsingular. Thus, we have

$$\begin{aligned}(L(I_{m+n} + tH))J_{m+n}(L(I_{m+n} + tH))^T - (Y(I_{m+n} + tS))(Y(I_{m+n} + tS))^T \\ = (L(I_{m+n} + tH))(J_{m+n} - ZZ^T)(L(I_{m+n} + tH))^T,\end{aligned}$$

where $Z = (L(I_{m+n} + tH))^{-1}(Y(I_{m+n} + tS)) = (I_{m+n} + tH)^{-1}L^{-1}(Y(I_{m+n} + tS))$. From (4.3) and noting (2.1), we get

$$\|(I_{m+n} + tH)^{-1}\|_2 \leq \frac{1}{1 - \|tH\|_2}.$$

Furthermore,

$$\|Z\|_2 \leq \frac{\|L^{-1}\|_2 (\|Y\|_2 + \|tSY\|_2)}{1 - \|tH\|_2} \leq \mu < 1. \quad (4.4)$$

Using Zeyl's theorem on eigenvalues of Hermitian matrices (e.g., [25], pp. 181), for the i -th eigenvalue of $J_{m+n} - ZZ^T$,

$$\lambda_i(J_{m+n}) + \lambda_{\min}(-ZZ^T) \leq \lambda_i(J_{m+n} - ZZ^T) \leq \lambda_i(J_{m+n}) + \lambda_{\max}(-ZZ^T).$$

Note that (4.4) implies $\lambda_{\min}(-ZZ^T) \geq -\mu^2$. Assume the eigenvalues of a matrix be ordered in non-decreasing order. Thus, for $0 < i \leq n$,

$$-1 - \mu^2 \leq \lambda_i(J_{m+n} - ZZ^T) \leq -1,$$

and for $n < i \leq m+n$,

$$1 - \mu^2 \leq \lambda_i(J_{m+n} - ZZ^T) \leq 1.$$

That is, $J_{m+n} - ZZ^T$ is nonsingular and has m positive eigenvalues and n negative eigenvalues, so is $(L(I_{m+n} + tH))J_{m+n}(L(I_{m+n} + tH))^T - (Y(I_{m+n} + tS))(Y(I_{m+n} + tS))^T$. In addition, we obtain

$$(L(I_{m+n} + tH))J_{m+n}(L(I_{m+n} + tH))^T - (Y(I_{m+n} + tS))(Y(I_{m+n} + tS))^T = \begin{bmatrix} G_{11} & G_{21}^T \\ G_{21} & -G_{22} \end{bmatrix},$$

where

$$\begin{aligned} G_{11} &= (L_{11}(I_m + tH_{11}))(L_{11}(I_m + tH_{11}))^T - (Y_m(I_m + tS_m))(Y_m(I_m + tS_m))^T \\ G_{22} &= (L_{22}(I_n + tH_{22}))(L_{22}(I_n + tH_{22}))^T - (L_{21}(I_n + tH_{21}))(L_{21}(I_n + tH_{21}))^T \\ &\quad + (Y_n(I_n + tS_n))(Y_n(I_n + tS_n))^T. \end{aligned}$$

Note that

$$tH = \begin{bmatrix} tH_{11} & 0 \\ * & * \end{bmatrix}. \quad (4.5)$$

Then $\|tH_{11}\|_2 \leq \|tH\|_2 < 1$, which implies that $(L_{11}(I_m + tH_{11}))$ is nonsingular. Thus, G_{11} can be rewritten as

$$G_{11} = (L_{11}(I_m + tH_{11}))(I_m - Z_m Z_m^T)(L_{11}(I_m + tH_{11}))^T,$$

where $Z_m = (L_{11}(I_m + tH_{11}))^{-1}(Y_m(I_m + tS_m))$. Similar to (4.5), we can note that Z_m is the principal submatrix with order m of Z . Then, using (4.4), $\|Z_m\|_2 \leq \|Z\|_2 \leq \mu < 1$. Accordingly, $I_m - Z_m Z_m^T$ is symmetric positive definite, so is G_{11} . Furthermore, G_{22} is clearly symmetric positive semi-definite. Thus, for each $|t| \leq \varepsilon$, $(L(I_{m+n} + tH))J_{m+n}(L(I_{m+n} + tH))^T - (Y(I_{m+n} + tS))(Y(I_{m+n} + tS))^T$ has the generalized Cholesky factorization

$$\begin{aligned} V(t)J_{m+n}V^T(t) &= (L(I_{m+n} + tH))J_{m+n}(L(I_{m+n} + tH))^T \\ &\quad - (Y(I_{m+n} + tS))(Y(I_{m+n} + tS))^T. \end{aligned}$$

Note that $V(0) = V$ and $V(\varepsilon) = V + \Delta V$. Then (4.2) holds. \square

THEOREM 4.2. *Given a lower triangular matrix $L \in \mathbb{R}^{(m+n) \times (m+n)}$ with positive diagonal elements and a matrix $Y \in \mathbb{R}^{(m+n) \times k}$ such that the generalized Cholesky factorization $VJ_{m+n}V^T = LJ_{m+n}L^T - YY^T$ holds. Suppose $W = I_{m+n} + N \in \mathbb{R}^{(m+n) \times (m+n)}$ and $U = I_{m+n} + M \in \mathbb{R}^{(m+n) \times (m+n)}$. If*

$$\|N\|_2 < 1, \quad \frac{\|L^{-1}\|_2 (\|Y\|_2 + \|YM\|_2)}{1 - \|N\|_2} < 1, \quad (4.6)$$

and

$$\begin{aligned} & \left(\| (LJ_{m+n}L^T) V^{-T} \|_2 \|V^{-1}N\|_F + \| (YY^T) V^{-T} \|_2 \|V^{-1}M\|_F \right. \\ & \quad \left. + \|V^{-1}NL\|_F^2 + \|V^{-1}MY\|_F^2 \right) < \frac{1}{2}, \end{aligned} \quad (4.7)$$

then the generalized Cholesky factorization (4.2) always exists and

$$\begin{aligned} \|\Delta V\|_F & \leq (2 + \sqrt{2}) \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\sqrt{1 + \zeta_D^2 \kappa(VD_{m+n}^{-1})} \right) \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F + \|Y\|_2^2 \|M\|_F) \\ & \quad + (2 + \sqrt{2}) \inf_{D_{m+n} \in \mathbb{D}_{m+n}} (\kappa(VD_{m+n}^{-1})) \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2). \end{aligned} \quad (4.8)$$

Proof. Let $W(t) = I_{m+n} + tN$ and $U(t) = I_{m+n} + tM$, for any $|t| \leq \varepsilon$. Using (1.3) leads to

$$\begin{aligned} & (L(I_{m+n} + tN))J_{m+n}(L(I_{m+n} + tN))^T - (Y(I_{m+n} + tM))(Y(I_{m+n} + tM))^T \\ & = (L(I_{m+n} + tN)) \left(J_{m+n} - (L(I_{m+n} + tN))^{-1} \left((Y(I_{m+n} + tM))(Y(I_{m+n} + tM))^T \right) \right. \\ & \quad \left. \times (L(I_{m+n} + tN))^{-T} \right) (L(I_{m+n} + tN))^T, \end{aligned}$$

and

$$\left\| (L(I_{m+n} + tN))^{-1} (Y(I_{m+n} + tM)) \right\|_2 \leq \frac{\|L^{-1}\|_2 (\|Y\|_2 + \|YM\|_2)}{1 - \|N\|_2},$$

and considering the results of Theorem 4.1, it follows that (4.6) holds,

$$(L(I_{m+n} + tN))J_{m+n}(L(I_{m+n} + tN))^T - (Y(I_{m+n} + tM))(Y(I_{m+n} + tM))^T$$

is positive definite and has the unique generalized Cholesky factorization, i.e.

$$\begin{aligned} & (V + \Delta V(t))J_{m+n}(V + \Delta V(t))^T \\ & = (L(I_{m+n} + tN))J_{m+n}(L(I_{m+n} + tN))^T - (Y(I_{m+n} + tM))(Y(I_{m+n} + tM))^T, \end{aligned} \quad (4.9)$$

which, with $\Delta V(0) = 0$ and $\Delta V(\varepsilon) = \Delta V$, implies (4.2). Using (4.9) and (1.3) leads to

$$\begin{aligned} & (tN(LJ_{m+n}L^T) + t(LJ_{m+n}L^T)N^T + t^2N(LJ_{m+n}L^T)N^T) - (tM(YY^T) \\ & \quad + tM(YY^T)M^T + t^2M(YY^T)M^T) \\ & = VJ_{m+n}(\Delta V(t))^T + \Delta V(t)J_{m+n}V^T + \Delta V(t)J_{m+n}(\Delta V(t))^T. \end{aligned}$$

Pre-multiplying the above equation by V^{-1} and post-multiplying by V^{-T} lead to

$$\begin{aligned}
 & J_{m+n}(\Delta V(t))^T V^{-T} + V^{-1} \Delta V(t) J_{m+n} \\
 &= t V^{-1} N (L J_{m+n} L^T) V^{-T} + t V^{-1} (L J_{m+n} L^T) N^T V^{-T} + t^2 V^{-1} N (L J_{m+n} L^T) N^T V^{-T} \\
 &\quad - \left(t V^{-1} M (Y Y^T) V^{-T} + t V^{-1} (Y Y^T) M^T V^{-T} + t^2 V^{-1} M (Y Y^T) M^T V^{-T} \right) \\
 &\quad - V^{-1} \Delta V(t) J_{m+n} (\Delta V(t))^T V^{-T}.
 \end{aligned} \tag{4.10}$$

Since $V^{-1} \Delta V(t)$ is a lower triangular matrix. Using the symbol ‘low’ and noting (4.10), we get

$$\begin{aligned}
 V^{-1} \Delta V(t) J_{m+n} &= \text{low} \left(t V^{-1} N (L J_{m+n} L^T) V^{-T} + t V^{-1} (L J_{m+n} L^T) N^T V^{-T} \right) \\
 &\quad + \text{low} \left(t^2 V^{-1} N (L J_{m+n} L^T) N^T V^{-T} \right) \\
 &\quad - \text{low} \left(t V^{-1} M (Y Y^T) V^{-T} + t V^{-1} (Y Y^T) M^T V^{-T} \right) \\
 &\quad - \text{low} \left(t^2 V^{-1} M (Y Y^T) M^T V^{-T} \right) \\
 &\quad - \text{low} \left(V^{-1} \Delta V(t) J_{m+n} (\Delta V(t))^T V^{-T} \right).
 \end{aligned} \tag{4.11}$$

Applying the Frobenius norm to (4.11) and considering (2.1), (2.7) and (2.8) yields

$$\begin{aligned}
 & \|V^{-1} \Delta V(t) J_{m+n}\|_F \\
 &\leq \sqrt{2} \left(\|t V^{-1} N (L J_{m+n} L^T) V^{-T}\|_F + \|t V^{-1} M (Y Y^T) V^{-T}\|_F \right) \\
 &\quad + \frac{1}{\sqrt{2}} \| (V^{-1} \Delta V(t) J_{m+n} (\Delta V(t))^T V^{-T}) \|_F \\
 &\quad + \frac{1}{\sqrt{2}} \left(\|t^2 V^{-1} N (L J_{m+n} L^T) N^T V^{-T}\|_F + \|t^2 V^{-1} M (Y Y^T) M^T V^{-T}\|_F \right) \\
 &\leq \sqrt{2} \left(\|t V^{-1} N\|_F \| (L J_{m+n} L^T) V^{-T} \|_2 + \|t V^{-1} M\|_2 \| (Y Y^T) V^{-T} \|_F \right) \\
 &\quad + \frac{1}{\sqrt{2}} \left(\|V^{-1} \Delta V(t)\|_F^2 + \|t V^{-1} N L\|_F^2 + \|t V^{-1} M Y\|_F^2 \right).
 \end{aligned} \tag{4.12}$$

Assume $p(t) = \|V^{-1} \Delta V(t) J_{m+n}\|_F$ and

$$\begin{aligned}
 q(t) &= 2 \left(\|t V^{-1} N\|_F \| (L J_{m+n} L^T) V^{-T} \|_2 + \|t V^{-1} M\|_2 \| (Y Y^T) V^{-T} \|_F \right) \\
 &\quad + \left(\|t V^{-1} N L\|_F^2 + \|t V^{-1} M Y\|_F^2 \right).
 \end{aligned}$$

Then, the above equation can be rewritten as

$$p(t)^2 - \sqrt{2} p(t) + q(t) \geq 0.$$

Equation (4.7), clearly shows that $q(t) < 1/2$, for $|t| \leq \varepsilon$. Therefore, $p(t) \leq p_1(t)$ or $p(t) \geq p_2(t)$, where

$$p_1(t) = \frac{1}{\sqrt{2}}(1 - \sqrt{1 - 2q(t)}) < p_2(t) = \frac{1}{\sqrt{2}}(1 + \sqrt{1 - 2q(t)}). \quad (4.13)$$

Note that $p(t)$ is continuous and $p(0) = 0 = p_1(0) < p_2(0) = \sqrt{2}$. Then $p(t) \leq p_1(t)$, for any $|t| \leq \varepsilon$. Consequently, $p(\varepsilon) \leq p_1(\varepsilon)$, i.e.

$$\|V^{-1}\Delta V(t)J_{m+n}\|_F \leq \frac{1}{\sqrt{2}}(1 - \sqrt{1 - 2q(t)}) < \frac{1}{\sqrt{2}}. \quad (4.14)$$

Putting $t = \varepsilon$ in (4.11)

$$\begin{aligned} V^{-1}\Delta VJ_{m+n} = & \text{low} \left(V^{-1}N(LJ_{m+n}L^T)V^{-T} + V^{-1}(LJ_{m+n}L^T)N^TV^{-T} \right) \\ & + \text{low} \left(V^{-1}N(LJ_{m+n}L^T)N^TV^{-T} \right) - \text{low} \left(V^{-1}M(YY^T)M^TV^{-T} \right) \\ & - \text{low} \left(V^{-1}M(YY^T)V^{-T} + V^{-1}(YY^T)M^TV^{-T} \right) \\ & - \text{low} \left(V^{-1}\Delta VJ_{m+n}(\Delta V)^TV^{-T} \right). \end{aligned} \quad (4.15)$$

Right multiplying by $D_{m+n} \in \mathbb{D}_{m+n}$, and noting $V = D_{m+n}\check{V}$ and (2.4) yield

$$\begin{aligned} \check{V}^{-1}\Delta VJ_{m+n} = & \text{low} \left(\check{V}^{-1}N(LJ_{m+n}L^T)V^{-T} + D_{m+n}(\check{V}^{-1}N(LJ_{m+n}L^T)V^{-T})^TD_{m+n}^{-1} \right) \\ & + \text{low} \left(\check{V}^{-1}N(LJ_{m+n}L^T)N^TV^{-T} \right) - \text{low} \left(\check{V}^{-1}M(YY^T)M^TV^{-T} \right) \\ & - \text{low} \left(\check{V}^{-1}M(YY^T)V^{-T} + D_{m+n}(\check{V}^{-1}M(YY^T)V^{-T})^TD_{m+n}^{-1} \right) \\ & - \text{low} \left(\check{V}^{-1}\Delta VJ_{m+n}(\Delta V)^TV^{-T} \right). \end{aligned} \quad (4.16)$$

Implementing the Frobenius norm on (4.16) and utilizing (2.5), (2.6), and (2.1) lead to

$$\begin{aligned} \|\check{V}^{-1}\Delta VJ_{m+n}\|_F & \leq \sqrt{1 + \zeta_{D_{m+n}}^2} (\|\check{V}^{-1}N(LJ_{m+n}L^T)V^{-T}\|_F \\ & \quad + \|\check{V}^{-1}M(YY^T)V^{-T}\|_F) \\ & \quad + \|\check{V}^{-1}N(LJ_{m+n}L^T)N^TV^{-T}\|_F + \|\check{V}^{-1}M(YY^T)M^TV^{-T}\|_F \\ & \quad + \|\check{V}^{-1}\Delta VJ_{m+n}(\Delta V)^TV^{-T}\|_F \\ & \leq \sqrt{1 + \zeta_{D_{m+n}}^2} \|\check{V}^{-1}\|_2 \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F + \|Y\|_2^2 \|M\|_F) \\ & \quad + \|\check{V}^{-1}\Delta V\|_2 \|V^{-1}\Delta V\|_F \\ & \quad + \|\check{V}^{-1}\|_2 \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2). \end{aligned} \quad (4.17)$$

Considering (4.14), (2.1), (4.6) and (4.7), we obtain

$$\begin{aligned} \|\check{V}^{-1}\Delta V J_{m+n}\|_F &\leq (2+\sqrt{2})\sqrt{1+\zeta_D^2}\|\check{V}^{-1}\|_2\|V^{-1}\|_2(\|L\|_2^2\|N\|_F + \|Y\|_2^2\|M\|_F) \\ &\quad + (2+\sqrt{2})\|\check{V}^{-1}\|_2\|V^{-1}\|_2(\|L\|_2^2\|N\|_F^2 + \|Y\|_2^2\|M\|_F^2), \end{aligned} \quad (4.18)$$

which, along with the fact that $\|J_{m+n}\|_2 = 1$ and the supporting proof

$$\|\Delta V\|_F \leq \|\check{V}\check{V}^{-1}\Delta V J_{m+n}\|_F \leq \|\check{V}\|_2\|\check{V}^{-1}\Delta V\|_F\|J_{m+n}\|_2, \quad \text{by (2.1)} \quad (4.19)$$

shows the bound (4.8). \square

To check the efficiency of bounds, let us prove that the bound (3.15) is significantly sharper than (4.8). Taking into consideration [26, Corollary 3.4], for any $D_{m+n} \in \mathbb{D}_{m+n}$ and $X \in \mathbb{R}^{(m+n) \times (m+n)}$, using (2.11) and (2.12), we obtain

$$\begin{aligned} \|Q_{LM}\|_2 &= \left\| \mathfrak{K}_{\text{Ivec}}(J_{m+n}^{-T} \otimes V)(I_{m+n} \otimes D_{m+n}^{-1})(I_{m+n} \otimes D_{m+n}) \right. \\ &\quad \times \left. \mathfrak{K}_{\text{low}}\left((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T))\Pi\right) \right\|_2 \\ &= \left\| \mathfrak{K}_{\text{Ivec}}(J_{m+n} \otimes VD_{m+n}^{-1}) \mathfrak{K}_{\text{low}}\left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T))\Pi\right) \right\|_2 \quad \text{by (2.12)} \\ &\leq \|VD_{m+n}^{-1}\|_2 \left\| \mathfrak{K}_{\text{low}}\left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T))\Pi\right) \right\|_2 \quad \text{by (2.11)} \\ &= \|VD_{m+n}^{-1}\|_2 \max_{\|\text{vec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}}\left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T))\Pi\right) \text{vec}(X) \right\|_2. \end{aligned} \quad (4.20)$$

Taking into account (2.9), (2.2), (2.4), (2.5) and (2.1) yields

$$\begin{aligned} &\max_{\|\text{vec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}}\left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T))\Pi\right) \text{vec}(X) \right\|_2 \\ &= \max_{\|\text{vec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \text{vec}\left(D_{m+n}V^{-1}X(LJ_{m+n}L^T)V^{-T} \right. \right. \\ &\quad \left. \left. + D_{m+n}V^{-1}(LJ_{m+n}L^T)X^T V^{-T}\right) \right\|_2 \quad \text{by (2.9)} \end{aligned}$$

$$\begin{aligned}
&= \max_{\|\text{vec}(X)\|_2=1} \left\| \text{vec} \left(\text{low} \left(D_{m+n} V^{-1} X (LJ_{m+n} L^T) V^{-T} \right. \right. \right. \\
&\quad \left. \left. \left. + D_{m+n} (D_{m+n} V^{-1} X (LJ_{m+n} L^T) V^{-T})^T D_{m+n}^{-1} \right) \right) \right\|_2 \quad \text{by (2.2)} \\
&= \max_{\|X\|_F=1} \left\| \text{low} \left(D_{m+n} V^{-1} X (LJ_{m+n} L^T) V^{-T} \right) \right. \\
&\quad \left. + D_{m+n} \text{low} \left(D_{m+n} V^{-1} X (LJ_{m+n} L^T) V^{-T} \right)^T D_{m+n}^{-1} \right\|_F \quad \text{by (2.4)} \\
&\leq \max_{\|X\|_F=1} \sqrt{1 + \zeta_{D_{m+n}}^2} \|D_{m+n} V^{-1} X (LJ_{m+n} L^T) V^{-T}\|_F \quad \text{by (2.5)} \\
&\leq \sqrt{1 + \zeta_{D_{m+n}}^2} \|D_{m+n} V^{-1}\|_2 \|(LJ_{m+n} L^T) V^{-T}\|_2. \quad \text{by (2.1)} \tag{4.21}
\end{aligned}$$

Hence, putting (4.21) into (4.20) gives

$$\|Q_{LM}\|_2 \leq \left(\inf_{D_{m+n} \in \mathbb{D}_{m+n}} \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \right) \|L\|_2^2 \|V^{-1}\|_2. \tag{4.22}$$

Therefore, we can demonstrate that,

$$\|Q_Y\|_2 \leq \left(\inf_{D_{m+n} \in \mathbb{D}_{m+n}} \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \right) \|Y\|_2^2 \|V^{-1}\|_2, \tag{4.23}$$

$$\|R_V\|_2 \leq \left(\inf_{D_{m+n} \in \mathbb{D}_{m+n}} \kappa(V D_{m+n}^{-1}) \right) \|V^{-1}\|_2, \tag{4.24}$$

(4.24) together with the fact $\kappa(V D_{m+n}^{-1}) \geq 1$ and (4.22) indicates that the bound (3.15) is absolutely tighter than (4.8).

REMARK 4.3. By ignoring the high-order terms, we can obtain a first-order multiplicative bound from (4.8):

$$\|\Delta V\|_F \leq \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \right) \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F + \|Y\|_2^2 \|M\|_F). \tag{4.25}$$

Obviously, from (4.22) and (4.23), we can check that the bound in Remark 3.2 is always tighter than the (4.25).

THEOREM 4.4. *With the same assumptions as in Theorem 4.1, we have*

$$\begin{aligned}
\|\Delta V\|_F &\leq (2 + \sqrt{2}) \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \right) \\
&\quad \times \left(\|\text{sut}((LJ_{m+n} L^T) V^{-T})\|_2 \|N\|_F + \|V^{-1}\|_2 \|Y\|_2^2 \|M\|_F \right) \\
&\quad + (2 + \sqrt{2}) \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\kappa(V D_{m+n}^{-1}) \right) \\
&\quad \times (\|\text{diag}((LJ_{m+n} L^T) V^{-T})\|_2 \|N\|_F + \|V^{-1}\|_2 (\|L\|_2^2 \|N\|_F^2 + \|Y\|_2^2 \|M\|_F^2)). \tag{4.26}
\end{aligned}$$

Proof. Applying the symbols 'sut', 'diag' and 'slt', $\text{low}(V^{-1}N(LJ_{m+n}L^T)V^{-T} + V^{-1}(LJ_{m+n}L^T)N^TV^{-T})$ can be rewritten as

$$\text{low}(V^{-1}N(LJ_{m+n}L^T)V^{-T} + V^{-1}(LJ_{m+n}L^T)N^TV^{-T}) \quad (4.27)$$

$$= V^{-1}N \text{diag}((LJ_{m+n}L^T)V^{-T}) + \text{low}(V^{-1}N \text{sut}((LJ_{m+n}L^T)V^{-T}) \\ + \text{slt}(V^{-1}(LJ_{m+n}L^T))N^TV^{-T}). \quad (4.28)$$

Substituting (4.27) into (4.15) and then pre-multiplying it by $D_{m+n} \in \mathbb{D}_{m+n}$ and using $V = D_{m+n}\check{V}$, we have

$$\begin{aligned} & \check{V}^{-1}\Delta VJ_{m+n} \\ &= V^{-1}N \text{diag}((LJ_{m+n}L^T)V^{-T}) \\ &+ \text{low}\left(V^{-1}N \text{sut}((LJ_{m+n}L^T)V^{-T}) + D_{m+n}(\text{slt}(V^{-1}(LJ_{m+n}L^T))N^TV^{-T})D_{m+n}^{-1}\right) \\ &+ \text{low}\left(\check{V}^{-1}N(LJ_{m+n}L^T)N^TV^{-T}\right) - \text{low}\left(\check{V}^{-1}M(Y Y^T)M^TV^{-T}\right) \\ &- \text{low}\left(\check{V}^{-1}M(Y Y^T)V^{-T} + D_{m+n}(\check{V}^{-1}M(Y Y^T)V^{-T})^TD_{m+n}^{-1}\right) \\ &- \text{low}\left(\check{V}^{-1}\Delta VJ_{m+n}(\Delta V)^TV^{-T}\right). \end{aligned} \quad (4.29)$$

Hence, we get the bound (4.26) by applying the Frobenius norm to (4.29) and utilizing (2.5), (2.6), (4.14), and (4.19).

Moving forward, we will show that the bound (3.21) is tighter than (4.26). By using (3.18), we get

$$\begin{aligned} \|Q_{LT}\|_2 &\leq \left\| (J_{m+n}^{-T} \otimes V)(I_{m+n} \otimes D_{m+n}^{-1})(I_{m+n} \otimes D_{m+n}) \mathfrak{K}_{\text{low}} \left((V^{-1}(LJ_{m+n}L^T) \otimes V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes V^{-1}(LJ_{m+n}L^T)) \Pi \right) \mathfrak{K}_{\text{lvec}}^T \right\| \\ &= \left\| (J_{m+n}^{-T} \otimes VD_{m+n}^{-1}) \mathfrak{K}_{\text{low}} \left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T)) \Pi \right) \mathfrak{K}_{\text{lvec}}^T \right\| \quad \text{by (2.12)} \\ &\leq \|VD_{m+n}^{-1}\|_2 \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \left((V^{-1}(LJ_{m+n}L^T) \otimes D_{m+n}V^{-1}) \right. \right. \\ &\quad \left. \left. + (V^{-1} \otimes D_{m+n}V^{-1}(LJ_{m+n}L^T)) \Pi \right) \mathfrak{K}_{\text{lvec}}^T \text{lvec}(X) \right\|_2. \end{aligned} \quad (4.30)$$

We assume X is a lower triangular matrix. Obviously, we only take the lower triangular part of X . As a result of using (2.2), (2.3), and (2.9), we get

$$\begin{aligned}
& \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \left((V^{-1} (LJ_{m+n} L^T) \otimes D_{m+n} V^{-1}) \right. \right. \\
& \quad \left. \left. + (V^{-1} \otimes D_{m+n} V^{-1} (LJ_{m+n} L^T)) \Pi \right) \mathfrak{K}_{\text{lvec}}^T \mathfrak{K}_{\text{lvec}} \text{vec}(X) \right\|_2 \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \left((V^{-1} (LJ_{m+n} L^T) \otimes D_{m+n} V^{-1}) \right. \right. \\
& \quad \left. \left. + (V^{-1} \otimes D_{m+n} V^{-1} (LJ_{m+n} L^T)) \Pi \right) \mathfrak{K}_{\text{lt}} \text{vec}(X) \right\|_2 \quad \text{by (2.3)} \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \left((V^{-1} (LJ_{m+n} L^T) \otimes D_{m+n} V^{-1}) \right. \right. \\
& \quad \left. \left. + (V^{-1} \otimes D_{m+n} V^{-1} (LJ_{m+n} L^T)) \Pi \right) \text{vec}(\text{lt}(X)) \right\|_2 \quad \text{by (2.2)} \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \left((V^{-1} (LJ_{m+n} L^T) \otimes D_{m+n} V^{-1}) \right. \right. \\
& \quad \left. \left. + (V^{-1} \otimes D_{m+n} V^{-1} (LJ_{m+n} L^T)) \Pi \right) \text{vec}(X) \right\|_2 \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \mathfrak{K}_{\text{low}} \text{vec} \left(D_{m+n} V^{-1} X (V^{-1} (LJ_{m+n} L^T))^T \right. \right. \\
& \quad \left. \left. + D_{m+n} V^{-1} (LJ_{m+n} L^T) X^T V^{-T} \right) \right\|_2 \quad \text{by (2.9)} \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \text{vec} \left(\text{low} \left(D_{m+n} V^{-1} X (V^{-1} (LJ_{m+n} L^T))^T \right. \right. \right. \\
& \quad \left. \left. + D_{m+n} V^{-1} (LJ_{m+n} L^T) X^T V^{-T} \right) \right) \right\|_2 \quad \text{by (2.2)} \\
&= \max_{\|\text{lvec}(X)\|_2=1} \left\| \text{vec} \left(\text{low} \left(D_{m+n} V^{-1} X \left(\text{sut} (V^{-1} (LJ_{m+n} L^T))^T \right. \right. \right. \right. \\
& \quad \left. \left. + D_{m+n} \left(\text{slt} (V^{-1} (LJ_{m+n} L^T)) + \text{diag} (V^{-1} (LJ_{m+n} L^T))^T \right) \right. \right. \\
& \quad \left. \left. + \text{diag} (V^{-1} (LJ_{m+n} L^T)) \right) X^T V^{-T} \right) \right) \right\|_2. \tag{4.31}
\end{aligned}$$

For any lower triangular matrix K , we have

$$\text{low}(K) + \text{low}(K^T) = K.$$

Therefore, if we set $K \equiv D_{m+n} V^{-1} X \text{diag}((LJ_{m+n} L^T) V^{-T})$, then

$$\begin{aligned}
& D_{m+n} V^{-1} X \text{diag}((LJ_{m+n} L^T) V^{-T}) \\
&= \text{low} \left(D_{m+n} V^{-1} X \text{diag} (V^{-1} (LJ_{m+n} L^T))^T + D_{m+n} \text{diag} (V^{-1} (LJ_{m+n} L^T)) X^T V^{-T} \right),
\end{aligned}$$

which together with (4.31) and (4.30) implies

$$\begin{aligned}
\|Q_{LT}\|_2 &\leq \|VD_{m+n}^{-1}\|_2 \max_{\|\text{lvec}(X)\|_2=1} \left\| \text{vec} \left(D_{m+n} V^{-1} X \text{diag}((LJ_{m+n}L^T)V^{-T}) \right. \right. \\
&\quad \left. \left. + \text{low} \left(D_{m+n} V^{-1} X \left(\text{sut}(V^{-1}(LJ_{m+n}L^T))^T \right) \right. \right. \right. \\
&\quad \left. \left. \left. + D_{m+n} \left(\text{slt}(V^{-1}(LJ_{m+n}L^T)) \right) X^T V^{-T} \right) \right) \right\|_2 \\
&\leq \|VD_{m+n}^{-1}\|_2 \max_{\|X\|_F=1} \|D_{m+n} V^{-1} X \text{diag}((LJ_{m+n}L^T)V^{-T})\|_F \\
&\quad + \|VD_{m+n}^{-1}\|_2 \max_{\|X\|_F=1} \left\| \text{low} \left(D_{m+n} V^{-1} X \left(\text{sut}(V^{-1}(LJ_{m+n}L^T))^T \right) \right. \right. \\
&\quad \left. \left. + D_{m+n} \left(\text{slt}(V^{-1}(LJ_{m+n}L^T)) \right) X^T V^{-T} \right) \right\|_F.
\end{aligned}$$

With the help of (2.5), we get

$$\begin{aligned}
\|Q_{LT}\|_2 &\leq \kappa(VD_{m+n}^{-1}) \left\| \text{diag}((LJ_{m+n}L^T)V^{-T}) \right\|_2 \\
&\quad + \max_{\|X\|_F=1} \left\| \text{low} \left(D_{m+n} V^{-1} X (\text{sut}((LJ_{m+n}L^T)V^{-T})) \right) \right. \\
&\quad \left. + D_{m+n} \left(\text{low}(D_{m+n} V^{-1} X (\text{sut}((LJ_{m+n}L^T)V^{-T})))^T D_{m+n}^{-1} \right) \right\|_F \\
&\leq \kappa(VD_{m+n}^{-1}) \left\| \text{diag}((LJ_{m+n}L^T)V^{-T}) \right\|_2 \\
&\quad + \sqrt{1 + \zeta_{D_{m+n}}^2} \|VD_{m+n}^{-1}\|_2 \max_{\|X\|_F=1} \left\| \left(D_{m+n} V^{-1} X (\text{sut}((LJ_{m+n}L^T)V^{-T})) \right) \right\|_F \\
&\leq \kappa(VD_{m+n}^{-1}) \left\| \text{diag}((LJ_{m+n}L^T)V^{-T}) \right\|_2 \\
&\quad + \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(VD_{m+n}^{-1}) \left\| (\text{sut}(LJ_{m+n}L^T)V^{-T}) \right\|_2. \tag{4.32}
\end{aligned}$$

This result together with the fact $\kappa(VD_{m+n}^{-1}) \geq 1$ and (4.23) illustrates that the bound (3.21) is significantly smaller than (4.26). \square

REMARK 4.5. By neglecting the high-order terms, we can derive a first-order multiplicative bound from (4.26):

$$\begin{aligned}
\|\Delta V\|_F &\leq \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(VD_{m+n}^{-1}) \right) \left(\|\text{sut}((LJ_{m+n}L^T)V^{-T})\|_2 \|N\|_F \right. \\
&\quad \left. + \|V^{-1}\|_2 \|Y\|_2^2 \|M\|_F \right) \\
&\quad + \inf_{D_{m+n} \in \mathbb{D}_{m+n}} \left(\kappa(VD_{m+n}^{-1}) \right) \left(\|\text{diag}((LJ_{m+n}L^T)V^{-T})\|_2 \|N\|_F \right). \tag{4.33}
\end{aligned}$$

From (4.32) and (4.23), it is easy to check that the bound in Remark 3.4 is tighter than (4.33), when $W = I_{m+n} + N$ is assumed to be a lower triangular matrix.

5. Numerical results

In this section, we provide three numerical examples to illustrate the results derived in Sections 3–4. We use an algorithm from [1] to obtain the generalized Cholesky factors L and V in (1.3). All numerical experiments are performed by using Matlab 2018a.

EXAMPLE 5.1. In first example, we compare the strong and weak multiplicative perturbation bounds. Let $A = [a_{ij}] = P_m + I_m \in \mathbb{R}^{m \times m}$ where $P_m = [p_{ij}]$ is the Pascal matrix, (i.e., $p_{1j} = p_{i1} = 1, p_{ij} = p_{i(j-1)} + p_{(i-1)j}$), $B = [b_{ij}] = 0.7 * [\max(i, j)] \in \mathbb{R}^{n \times m}$ and $C = (c_{ij}) \in \mathbb{R}^{n \times n}$ is the Lehmar matrix, (i.e., $c_{ij} = i/j$ for $j \geq i$), $J_{m+n} = \text{diag}(I_m, -I_n)$ and $Y = P\Lambda P^T$, where $P \in \mathbb{R}^{(m+n) \times (m+n)}$ is an orthogonal matrix taken from the QR factorization, $\Lambda = \text{diag}(d, d, \dots, d) \in \mathbb{R}^{(m+n) \times (m+n)}$ with $d = 0.02$. From [11], the scaled matrix D_{m+n} is denoted as follows: suppose $\zeta_1 = \sqrt{\sum_{j=1}^{m+n} f_{1j}^2}$, $\zeta_i = \sqrt{\sum_{j=i}^{m+n} f_{ij}^2}$ if $\sqrt{\sum_{j=1}^{m+n} f_{ij}^2} \leq \zeta_{i-1}$ otherwise $\zeta_i = \zeta_{i-1}$, for $i = 2, \dots, m+n$, $F = (f_{ij}) = V^T$.

Table 1: Results for various values of n , m and k

n, m, k	5, 5, 10	10, 10, 20	13, 12, 25	18, 12, 30	20, 20, 40	40, 10, 50
$\ Q_{LM}\ _2$	4.1503	13.0342	26.3812	39.3403	52.5054	87.4032
$t_{\ Q_{LM}\ _2}$	0.011385	0.014349	0.039375	0.497504	2.567943	9.750072
γ_{11}	32.9031	190.1043	530.2156	6.5214e+02	2.5475e+03	5.6403e+03
$t_{\gamma_{11}}$	0.014312	0.015310	0.016385	0.018066	0.015910	0.019883
$\ Q_Y\ _2$	0.1832	0.2072	0.3702	0.5361	1.0945	1.7304
$t_{\ Q_Y\ _2}$	0.014291	0.021533	0.028610	0.169868	3.543310	13.918751
γ_{12}	0.4732	3.1903	9.0704	39.1204	67.0123	124.2617
$t_{\gamma_{12}}$	0.011736	0.014821	0.015570	0.016765	0.015092	0.019160
$\ Q_{LT}\ _2$	2.0145	3.1026	3.5017	3.7102	3.9143	4.9813
$t_{\ Q_{LT}\ _2}$	0.015054	0.019183	0.029361	0.182677	0.577617	3.769436
γ_{13}	8.0537	21.7241	33.2653	59.4293	83.3427	162.3076
$t_{\gamma_{13}}$	0.015381	0.017907	0.017463	0.015684	0.018650	0.019702
$\ R_V\ _2$	11.3821	26.4376	51.0436	132.3054	174.3076	294.2643
$t_{\ R_V\ _2}$	0.016202	0.015268	0.022487	0.014396	0.015743	0.017608
γ_{14}	23.0543	146.0354	397.0548	3.4256e+02	7.5473e+02	3.7354e+03
$t_{\gamma_{14}}$	0.014631	0.017183	0.016836	0.016179	0.015437	0.019326
$\ R_Y\ _2$	0.2537	0.3741	0.7451	1.5803	3.5071	4.9644
$t_{\ R_Y\ _2}$	0.015494	0.015406	0.013707	0.016474	0.014801	0.015409
γ_{15}	0.4322	1.3605	6.8065	31.7402	59.4223	95.7733
$t_{\gamma_{15}}$	0.014988	0.015179	0.014046	0.019076	0.016228	0.019694

Moreover, in Table 1, we denote

$$\gamma_{11} = \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(VD_{m+n}^{-1}) \|L\|_2^2 \|V^{-1}\|_2, \quad \gamma_{12} = \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(VD_{m+n}^{-1}) \|Y\|_2^2 \|V^{-1}\|_2,$$

$$\gamma_{13} = \kappa(VD_{m+n}^{-1}) \left(\|\text{diag}((LJ_{m+n}L^T)V^{-T})\|_2 + \sqrt{1 + \zeta_{D_{m+n}}^2} \|\text{slt}((LJ_{m+n}L^T)V^{-T})\|_2 \right),$$

$\gamma_{14} = \kappa(VD_{m+n}^{-1}) \|L\|_2^2 \|V^{-1}\|_2, \quad \gamma_{15} = \kappa(VD_{m+n}^{-1}) \|Y\|_2^2 \|V^{-1}\|_2,$

and $t_{(\cdot)}$ denote the time cost in seconds for computing the bounds (3.15), (3.21), (4.8), (4.26).

From Table 1, we can see that the strong rigorous multiplicative perturbation bounds (3.15) and (3.22), the rows marked by $\|Q_{LM}\|_2, \|Q_Y\|_2, \|Q_{LT}\|_2, \|R_V\|_2 \|L\|_2^2$ and $\|R_V\|_2 \|Y\|_2^2$ are always tighter than the weak rigorous multiplicative perturbation bounds (4.8) and (4.26), the rows marked by $\gamma_{11}, \gamma_{12}, \gamma_{13}, \gamma_{14}$ and γ_{15} . In addition, we can also observe that it is indeed more expensive to estimate the bounds (3.15) and (3.22); compare the rows marked by $t_{\gamma_{11}}, t_{\gamma_{12}}, t_{\gamma_{13}}, t_{\gamma_{14}}$ and $t_{\gamma_{15}}$.

EXAMPLE 5.2. The test matrix K_1 and K_2 are set to be

$$K_1 = \begin{bmatrix} 70 & 70 & 70 & 3 & 3 \\ 70 & 90 & 100 & 7 & 9 \\ 70 & 100 & 200 & 13 & 17 \\ 3 & 7 & 13 & -20 & -56 \\ 3 & 9 & 17 & -56 & -80 \end{bmatrix}, \quad K_2 = \begin{bmatrix} 2.4957 & -0.9922 & 0.9008 & -1.0195 & -0.3655 \\ -0.9922 & 2.4957 & -0.9922 & 0.9008 & -1.0195 \\ 0.9008 & -0.9922 & 2.4957 & -0.9922 & 0.9008 \\ -1.0195 & 0.9008 & -0.9922 & -2.4957 & 0.9922 \\ -0.3655 & -1.0195 & 0.9008 & 0.9922 & -2.4957 \end{bmatrix},$$

here $Y = \mu [0.240 \ -0.899 \ 0.899 \ 1.560 \ -2.390]^T$ and $J_{3+2} = \text{diag}(I_3, -I_2)$.

Table 2: Results for various values of μ

ω	Test matrices	μ_1	μ_2	μ_3	μ_4	μ_5	μ_6
$\ Q_{LM}\ _2$	K_1	1.2903	1.2903	1.2903	1.2904	1.2904	1.2904
	K_2	1.5214	1.5214	1.5214	1.5215	1.5215	1.5215
γ_{11}	K_1	1.2643e+02	1.2643e+02	1.2872e+02	1.7642e+02	1.7642e+02	1.7642e+02
	K_2	1.4453e+02	1.4453e+02	1.4467e+02	1.9362e+02	1.9362e+02	1.9362e+02
$\ Q_Y\ _2$	K_1	2.2632	2.2580	2.2531	0.0232	3.3227e-04	4.2827e-06
	K_2	2.5214	2.5112	2.5034	0.3204	5.3097e-04	7.3437e-06
γ_{12}	K_1	3.9126	3.9042	2.9959	1.4420	4.8028e-02	4.7028e-04
	K_2	5.8632	5.8436	5.6209	1.8062	9.9201e-02	8.7403e-03
$\ Q_{LT}\ _2$	K_1	1.1357	1.1357	1.1357	1.1357	1.1357	1.1357
	K_2	1.4792	1.4792	1.4792	1.4792	1.4792	1.4792
γ_{13}	K_1	76.1082	76.3042	76.6201	77.3321	77.4508	77.6902
	K_2	84.1376	84.4201	84.4853	85.0675	85.3065	85.3457
$\ R_V\ _2$	K_1	7.3214e+01	7.3214e+01	7.5326e+01	7.5422e+01	7.6402e+01	7.6402e+01
	K_2	9.3053e+01	9.3053e+01	9.4467e+01	9.4761e+01	9.9502e+01	9.9502e+01
γ_{14}	K_1	1.1643e+02	1.1643e+02	1.1872e+02	1.6642e+02	1.6642e+02	1.6642e+02
	K_2	1.3453e+02	1.3453e+02	1.3467e+02	1.8362e+02	1.8362e+02	1.8362e+02
$\ R_Y\ _2$	K_1	0.7861	0.7835	0.6328	0.3116	1.2092e-03	1.0453e-05
	K_2	0.9861	0.9835	0.8328	0.5116	1.1654e-02	1.1921e-04
γ_{15}	K_1	3.6126	3.6042	2.9059	1.0420	3.2028e-02	3.2028e-04
	K_2	5.5632	5.5436	5.3209	1.2062	5.9201e-02	5.7403e-03

Table 2 provides the numerical results for various values of μ ’s, as follows:

$\mu_1 = 1.004015006005433e-2, \quad \mu_2 = 1.003021021209640e-2,$
 $\mu_3 = 9.036225416303058e-3, \quad \mu_4 = \mu_3e-1, \quad \mu_5 = \mu_3e-3, \quad \mu_6 = \mu_3e-5.$

From Table 2, we can find that the strong multiplicative perturbation bounds (3.15) and (3.21) are always less than the weak multiplicative perturbation bounds (4.8) and (4.26), where $\gamma_{11}, \gamma_{12}, \gamma_{13}, \gamma_{14}$ and γ_{15} are given in the previous example. Furthermore, Tables 1 and 2 show that when the multiplicative perturbation matrix $W = I_{m+n} + N$ is set to be a lower triangular matrix, the strong and weak multiplicative perturbation bounds for the GCBD problem are smaller than the corresponding ones when the matrix $W = I_{m+n} + N$ is set to be a general matrix.

EXAMPLE 5.3. Let

$$L = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 & 0 & 0 \\ -e & 1 & 0 & 0 & 0 & 0 & 0 \\ -e & -e & 1 & 0 & 0 & 0 & 0 \\ -e & -e & -e & 1 & 0 & 0 & 0 \\ -e & -e & -e & -e & 1 & 0 & 0 \\ -e & -e & -e & -e & -e & 1 & 0 \\ -e & -e & -e & -e & -e & -e & 1 \end{bmatrix} \text{diag}\left(1, \delta, \delta^2, \delta^3, \delta^4, \delta^5, \delta^6\right),$$

$$Y = \mu \begin{bmatrix} 0.240 \\ -0.899 \\ 0.899 \\ 1.360 \\ -2.190 \\ 1.560 \\ 2.301 \end{bmatrix}, \quad J_{4+3} = \text{diag}(I_4, -I_3),$$

where $e = 0.98$ and $\delta = \sqrt{1 - e^2}$. From [11, 12], we have the following strong and weak additive rigorous perturbation bounds respectively:

$$\begin{aligned} G_{LG} &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1} \otimes V^{-1} L J_{m+n}) \Pi + (V^{-1} L J_{m+n}^T \otimes V^{-1})), \\ G_Y &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1} \otimes V^{-1} Y) \Pi + (V^{-1} Y \otimes V^{-1})), \\ H_V &= \mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}(V^{-1} \otimes V^{-1}), \\ G_{LT} &= (\mathfrak{K}_{\text{lvec}}(J_{m+n}^{-T} \otimes V) \mathfrak{K}_{\text{low}}((V^{-1} \otimes V^{-1} L J_{m+n}) \Pi + (V^{-1} L J_{m+n}^T \otimes V^{-1}))) \mathfrak{K}_{\text{lvec}}^T, \end{aligned}$$

and

$$\begin{aligned} \beta_{11} &= \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \|V^{-1} L\|_2, \quad \beta_{12} = \sqrt{1 + \zeta_{D_{m+n}}^2} \kappa(V D_{m+n}^{-1}) \|V^{-1} Y\|_2, \\ \beta_{13} &= \kappa(V D_{m+n}^{-1}) (\|\text{diag}(V^{-1} L)\|_2 + \sqrt{1 + \zeta_{D_{m+n}}^2} \|\text{slt}(V^{-1} L)\|_2), \\ \beta_{14} &= \kappa(V D_{m+n}^{-1}) \|V^{-1}\|_2. \end{aligned}$$

In this example, we compare the additive and multiplicative rigorous perturbation bounds for different values of μ 's, which are given above. From Table 3, we can clearly observe that the strong and weak multiplicative bounds $\|Q_{LM}\|_2$, $\|Q_Y\|_2$, $\|Q_{LT}\|_2$, $\|R_V\|_2$, and γ_{11} , γ_{12} , γ_{13} , γ_{14} are tighter than the strong and weak additive perturbation

Table 3: Comparison of additive and multiplicative rigorous perturbation bounds

ω	μ_1	μ_2	μ_3	μ_4	μ_5	μ_6
$\ Q_{LM}\ _2$	2.2997	2.2996	2.2983	2.2981	2.2981	2.2981
$\ G_{LG}\ _2$	12.6871	12.6871	12.6871	12.6871	12.6871	12.6871
$\ Q_Y\ _2$	21.4490	21.0468	9.0424	0.0592	5.4072e-06	5.9131e-10
$\ G_Y\ _2$	43.2302	43.6391	26.3015	1.8391	6.3261e-02	7.2604e-03
$\ Q_{LT}\ _2$	1.3628	1.3628	1.3628	1.3628	1.3628	1.3628
$\ G_{LT}\ _2$	2.5943	2.5943	2.5943	2.5943	2.5943	2.5943
$\ R_V\ _2$	46.3491	46.3491	48.3607	48.3607	57.5203	57.7287
$\ H_V\ _2$	89.6343	89.6343	92.7343	92.9807	98.8408	98.8626
γ_{11}	8.0674e+02	8.2665e+02	8.5674e+02	8.5832e+02	8.7609e+02	8.9612e+02
β_{11}	1.4553e+03	1.4597e+03	1.6138e+03	1.6234e+03	1.9721e+03	1.9860e+03
γ_{12}	46.5435	45.6645	19.4237	1.1456	1.2537e-03	1.2537e-06
β_{12}	1.4806e+03	1.4540e+03	686.5176	44.3787	0.4431	0.1434
γ_{13}	35.0578	35.0389	35.0104	17.1122	17.1307	17.1307
β_{13}	106.3003	104.3310	47.1390	37.2814	37.1612	37.1612
γ_{14}	5.3607e+02	5.3607e+02	5.4241e+02	5.4241e+02	5.7437e+02	5.8327e+02
β_{14}	1.2062e+03	1.2062e+03	1.2938e+03	1.2938e+03	1.6072e+03	1.6849e+03

bounds $\|G_{LG}\|_2, \|G_Y\|_2, \|G_{LT}\|_2, \|H_V\|_2$, and $\beta_{11}, \beta_{12}, \beta_{13}, \beta_{14}$ respectively. The multiplicative perturbation bounds are useful to estimate the tighter bounds since the matrix is significantly ill-conditioned.

Acknowledgements. We acknowledge the principal editor and anonymous referees for their useful comments and constructive suggestions which helped considerably to improve the quality of the paper.

REFERENCES

[1] J. X. ZHAO, *The generalized Cholesky factorization method for saddle point problems*, Appl. Math. Comput. 92 (1998) 49–58.

[2] X. M. FANG, *The generalized Cholesky factorization and perturbation of the real symmetric matrices*, J. Zhaoqing. Univ. 29 (2008) 14–16.

[3] Å. BJÖRCK, H. PARK, L. ELDÉN, *Accurate downdating of least squares solutions*, SIAM J. Matrix Anal. Appl. 15 (1994) 549–568.

[4] A. W. BOJANCZYK, R. P. BRENT, V. P. DOOREN, F. R. DE HOOG, *A note on downdating the Cholesky factorization*, SIAM J. Sci. Statist. Comput. 8 (1987) 210–221.

[5] X. W. CHANG, C. C. PAIGE, *Perturbation analyses for the Cholesky downdating problem*, SIAM J. Matrix Anal. Appl. 19 (1998) 429–443.

[6] L. ELDÉN, H. PARK, *Perturbation analysis for block downdating of a Cholesky decomposition* Numerische Mathematik. 68 (1994) 457–467.

[7] C. PAN, *A perturbation analysis of the problem of downdating a Cholesky factorization*, Linear Algebra Appl. 183 (1993) 103–115.

[8] G. STEWART, *The effects of rounding error on an algorithm for downdating a Cholesky factorization*, J. Inst. Math. Appl. 23 (1979) 203–213.

[9] J. SUN, *Perturbation analysis of the Cholesky downdating and QR updating problems*, SIAM J. Matrix Anal. Appl. 16 (1995) 760–775.

[10] A. FAROOQ, M. SAMAR, H. LI, C. MU, *Sensitivity analysis for the block Cholesky downdating problem*, Int. J. Comput. Math. 97 (2020) 1234–1253.

[11] H. LI, H. YANG, H. SHAO, *Perturbation analysis for block downdating of the generalized cholesky factorization*, Appl. Math. Comput. 218 (2012) 9451–9461.

- [12] A. FAROOQ, M. SAMAR, *Sensitivity analysis for the generalized Cholesky block downdating problem*, Linear and Multilinear Algebra 70 (2022), 997–1022.
- [13] X. W. CHANG, R. C. LI, *Multiplicative perturbation analysis for QR factorizations*, Num. Algebra Control Optim. 1 (2011) 301–316.
- [14] A. FAROOQ, M. SAMAR, H. LI AND C. MU, *Improved rigorous multiplicative perturbation bounds for the generalized Cholesky factorization and the Cholesky-like factorization*, Math. Inequal. Appl. 22 (2019) 133–149.
- [15] A. FAROOQ, M. SAMAR, *Multiplicative perturbation bounds for the block Cholesky downdating problem*, Int. J. Comput. Math. 97 (2020) 2421–2435.
- [16] H.-Y. LI, Y. YANG, *Rigorous multiplicative perturbation bounds for the generalized Cholesky factorization and the Cholesky-like factorization*, J. Math. Inequal. 8 (2014) 925–937.
- [17] Y. YANG, H. LI, *Multiplicative perturbation bounds for the SR factorization*, J. Math. Res. Appl. 34 (2014) 423–434.
- [18] H.-Y. LI, Y. WEI, *Improved rigorous perturbation bounds for the LU and QR factorizations*, Numer. Linear Algebra Appl. 22 (2015) 1115–1130.
- [19] M. SAMAR, A. FAROOQ, H. LI, C. MU, *Sensitivity analysis for the generalized Cholesky factorization*, App. Math. Comput. 362 (2019) 124556.
- [20] H.-Y. LI, Y. M. WEI, Y. YANG, *New rigorous perturbation bounds for the Cholesky-like factorization of skew-symmetric matrix*, Linear Algebra Appl. 491 (2016) 83–100.
- [21] X.-W. CHANG, *Perturbation analysis of Some Matrix factorization*, PhD thesis, School of Computer Science, McGill University, Montreal, 1997.
- [22] G. W. STEWART, J.-G. SUN, *Matrix Perturbation Theory*, Academic Press, Boston, 1990.
- [23] X.-W. CHANG, C. C. PAIGE, G. W. STEWART, *Perturbation analyses for the QR factorization*, SIAM J. Matrix Anal. Appl. 18 (1997) 775–791.
- [24] R. A. HORN, C. R. JOHNSON, *Topics in matrix analysis*, Cambridge University Press, 1991.
- [25] R. A. HORN, C. R. JOHNSON, *Matrix Analysis*, Cambridge University Press, Cambridge, UK, 1985.
- [26] W. LI, Z. XIE, S. VONG, *Sensitivity analysis for the symplectic QR factorization*, J. Franklin. Inst. 353 (2016) 1186–1205.

(Received April 19, 2022)

Mahvish Samar
College of Mathematics and Computer Science
Zhejiang Normal University
Jinhua 321004, China
e-mail: mahvishsamar@hotmail.com

Xinzhong Zhu
College of Mathematics and Computer Science
Zhejiang Normal University
Jinhua 321004, China

