

### Homework 9

Due by April 13, 2021

1. A small barber shop, operated by a single barber, has room for only two costumers. Potential costumers arrive at a Poisson rate of 3 per hour, and the successive serving times are independent exponential random variables of mean  $1/4$  hour.
  - (a) What is the average number of customers in the shop?
  - (b) What is the proportion of potential costumers that enters the shop?
  - (c) If the barber could work 3 times as fast, how much more businesss would he do (in average)? (Hint: Compare potential costumers that enters the shop.)
2. A call center has two agents, John and Mary. It has four phone lines. An incoming call that finds all lines are occupied receives a busy signal and is rejected. An accepted call finds all agents busy waits in the queue. While waiting (listening to music), the caller may abandon without service. When a caller's waiting time exceeds her patience time, the caller abandons without service. The patience times are iid exponentially distributed with mean 4 minutes. John's processing times of calls are iid exponentially distributed with mean 2 minutes. Mary's processing times of calls are iid exponentially distributed with mean 1 minute. The arrival process is Poisson with rate  $\lambda = 1$  call per minute. An arrival to an empty system always is always processed by John.
  - (a) Describe a CTMC so it can be used to compute the probability in the next part. You need to specify all input data for this CTMC including the state space.
  - (b) Using Matlab/Python to compute the probability that there are three calls in system at time  $t = 2$  minutes given that the system is empty at  $t = 0$ .
  - (c) Find the expected number of calls in system at time  $t = 2$  given the system is empty at  $t = 0$ .
  - (d) Long-run average utilization of each agent.
  - (e) Long-run average queue size.
  - (f) The rate at which calls are leaving agents.
3. Consider a call center that is staffed by  $K$  agents with three phone lines. Call arrivals follow a Poisson process with rate 2 per minute. An arriving call that finds all lines busy is lost. An arriving call that finds all agents busy will wait in a phone line until it enters into service. Call processing times are exponentially distributed with mean 1 minute.

- (a) Find the throughput and average waiting time when  $K = 1$ .
  - (b) Find the throughput and average waiting time when  $K = 2$ .
  - (c) Find the throughput and average waiting time when  $K = 3$ .
4. For problem 2, please find
- (a) long-run fraction of calls gets busy signal.
  - (b) average waiting time among those calls that enter into system.
  - (c) fraction of calls that enter into system, but later abandon without service.
5. Consider a CTMC on state space  $S$  with generator  $G$ . Prove that a distribution  $\pi$  on  $S$  is a *stationary distribution* of the CTMC if and only if for any “cut” (partition)  $(A, A^c)$  with  $A \subset S$ :

$$\sum_{i \in A} \sum_{j \in A^c} \pi(i) G_{ij} = \sum_{i \in A^c} \sum_{j \in A} \pi(i) G_{ij}.$$