

CSC 4020 Fundamental of Machine Learning: Linear Regression

Baoyuan Wu
School of Data Science, CUHK-SZ

January 25/27, 2021

Outline

- 1 Some illustrations and review of last week
- 2 Linear Regression: A Deterministic Perspective
- 3 Linear Regression: A Probabilistic Perspective
 - Probabilistic modeling
 - Robust linear regression
 - Ridge regression
 - Lasso regression
- 4 Generalized Linear Regression

Some illustrations

- **Slides:** On each Saturday, TA will upload the slides without notes of the next week to the BB system, and the slides with notes and the zoom videos will be uploaded after the class.

Some illustrations

- **Slides:** On each Saturday, TA will upload the slides without notes of the next week to the BB system, and the slides with notes and the zoom videos will be uploaded after the class.
- **Participation and quiz:** we will have 5 random quizzes with 2 scores for each, and 10 scores in total. There will be one or two questions in each quiz, and a few minutes will be left at the end of the class to finish the quiz. Anyone submits the quiz to TA will earn 1 score for participation, and gives the correct answer to the question(s) will earn another 1 score. The policy of answering questions will be replaced by this new policy. To ensure fairness, in the first quiz, students who had answered questions in last week will automatically earn 2 scores if he/she submits the quiz.

Some illustrations

- **Slides:** On each Saturday, TA will upload the slides without notes of the next week to the BB system, and the slides with notes and the zoom videos will be uploaded after the class.
- **Participation and quiz:** we will have 5 random quizzes with 2 scores for each, and 10 scores in total. There will be one or two questions in each quiz, and a few minutes will be left at the end of the class to finish the quiz. Anyone submits the quiz to TA will earn 1 score for participation, and gives the correct answer to the question(s) will earn another 1 score. The policy of answering questions will be replaced by this new policy. To ensure fairness, in the first quiz, students who had answered questions in last week will automatically earn 2 scores if he/she submits the quiz.
- **About the contents:** this is a fundamental course of machine learning, and our students are from different grades and have different backgrounds. Thus, the most important and basic contents will be mainly covered in this course. But, I will provide some reading materials if one is interested in some special topics, and welcome to discuss with me after the class or in office hour. Any suggestion about some interesting/cutting-edge machine learning topics are welcomed, and I will try to cover some.

Some illustrations

- **Slides:** On each Saturday, TA will upload the slides without notes of the next week to the BB system, and the slides with notes and the zoom videos will be uploaded after the class.
- **Participation and quiz:** we will have 5 random quizzes with 2 scores for each, and 10 scores in total. There will be one or two questions in each quiz, and a few minutes will be left at the end of the class to finish the quiz. Anyone submits the quiz to TA will earn 1 score for participation, and gives the correct answer to the question(s) will earn another 1 score. The policy of answering questions will be replaced by this new policy. To ensure fairness, in the first quiz, students who had answered questions in last week will automatically earn 2 scores if he/she submits the quiz.
- **About the contents:** this is a fundamental course of machine learning, and our students are from different grades and have different backgrounds. Thus, the most important and basic contents will be mainly covered in this course. But, I will provide some reading materials if one is interested in some special topics, and welcome to discuss with me after the class or in office hour. Any suggestion about some interesting/cutting-edge machine learning topics are welcomed, and I will try to cover some.
- Welcome to the office hour at **Wednesday 10:30–11:30am in DY 411.**

Review of last week

- **Probability theory:**

- Discrete probability distributions: Bernoulli, Binomial, Beta
- Continuous probability distributions: Gaussian, Student t , Laplace


Review of last week

- **Probability theory:**

- Discrete probability distributions: Bernoulli, Binomial, Beta
- Continuous probability distributions: Gaussian, Student t , Laplace

- **Information theory:**


- Information
- Entropy, marginal/conditional/joint entropy, relative entropy (KL divergence, mutual information)




Properties of KL divergence

- $D(p_X(x)||q_X(x)) \geq 0$ with equality if and only if $p_X(x) = q_X(x)$.
- $D(p_X(x)||q_X(x)) \neq D(q_X(x)||p_X(x))$

$$q^* = \operatorname{argmin}_q D_{\text{KL}}(p||q)$$



$$q^* = \operatorname{argmin}_q D_{\text{KL}}(q||p)$$



One constraint with respect to q is missing at last time, i.e., it is the single mode distribution! More detailed derivations could be found at <https://dibyaghosh.com/blog/probability/kldivergence.html>.


Linear regression

- Here we start from a simple example of one dimensional input variable, and the training dataset $D = \{(x_i, y_i)\}_{i=1}^m$ can be plotted on the $x - y$ plane.

Linear regression

- Here we start from a simple example of one dimensional input variable, and the training dataset $D = \{(x_i, y_i)\}_{i=1}^m$ can be plotted on the $x - y$ plane.
- m indicates the number of training samples; x denotes the input variable/feature; y denotes the output variable.

Size in feet ² (x)	Price in 1000's (y)
2104	460
1416	232
1514	315
852	178
...	...



Linear hypothesis function

- Our goal is to find a linear hypothesis function to well fit the training data D , *i.e.*,

$$h_{\theta}(x) = \underbrace{\theta_0}_{\text{bias}} + \underbrace{\theta_1 \phi(x)}_{\text{weight}} = [\theta_0, \theta_1][1; \phi(x)] = \underbrace{\hat{\phi}(x)^{\top}}_{\text{feature vector}} \underbrace{\theta}_{\text{weight vector}} \quad (1)$$

Linear hypothesis function

- Our goal is to find a linear hypothesis function to well fit the training data D , i.e.,

$$h_{\theta}(x) = \theta_0 + \theta_1 \phi(x) = [\theta_0, \theta_1][1; \phi(x)] = \hat{\phi}(x)^{\top} \theta \quad (1)$$

where $\phi(x)$ is called **basis expansion**, which is specified as different forms, such as $\phi(x) = x$ or $\phi(x) = [x^3; x^2; x]$. In the following, we will use $\phi(x) = x$ as example, while other expansions will be introduced later.

$$\begin{matrix} & x^3 \\ [\theta_0, \theta_2, \theta_3] & x^2 \\ & x^1 \end{matrix}$$

Linear hypothesis function

- Our goal is to find a linear hypothesis function to well fit the training data D , *i.e.*,

$$h_{\theta}(x) = \theta_0 + \theta_1 \phi(x) = [\theta_0, \theta_1][1; \phi(x)] = \hat{\phi}(x)^{\top} \theta \quad (1)$$

where $\phi(x)$ is called **basis expansion**, which is specified as different forms, such as $\phi(x) = x$ or $\phi(x) = [x^3; x^2; x]$. In the following, we will use $\phi(x) = x$ as example, while other expansions will be introduced later.

- Given θ_0, θ_1 , $h_{\theta}(x)$ is the function of x .

Linear hypothesis function

- Our goal is to find a linear hypothesis function to well fit the training data D , *i.e.*,

$$h_{\theta}(x) = \theta_0 + \theta_1 \phi(x) = [\theta_0, \theta_1][1; \phi(x)] = \hat{\phi}(x)^{\top} \theta \quad (1)$$

where $\phi(x)$ is called **basis expansion**, which is specified as different forms, such as $\phi(x) = x$ or $\phi(x) = [x^3; x^2; x]$. In the following, we will use $\phi(x) = x$ as example, while other expansions will be introduced later.

- Given θ_0, θ_1 , $h_{\theta}(x)$ is the function of x .
- Given x , $h_{\theta}(x)$ is a **linear function** of $\theta = [\theta_0; \theta_1]$. This is why it is called **linear regression**.

$$\underline{x^{\top}} \underline{\theta}$$


Linear hypothesis function

- Our goal is to find a linear hypothesis function to well fit the training data D , i.e.,

$$h_{\theta}(x) = \theta_0 + \theta_1 \phi(x) = [\theta_0, \theta_1][1; \phi(x)] = \hat{\phi}(x)^{\top} \theta \quad (1)$$

where $\phi(x)$ is called **basis expansion**, which is specified as different forms, such as $\phi(x) = x$ or $\phi(x) = [x^3; x^2; x]$. In the following, we will use $\phi(x) = x$ as example, while other expansions will be introduced later.

- Given θ_0, θ_1 , $h_{\theta}(x)$ is the function of x .
- Given x , $h_{\theta}(x)$ is a **linear function** of $\theta = [\theta_0; \theta_1]$. This is why it is called **linear regression**.
- Then, given D , how to learn θ ?



Cost function

- We design the following **cost function** to minimize the difference between the prediction $h_{\theta}(x_i)$ and the ground-truth value y_i , i.e.,

$$\min_{\theta} J(\theta) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x_i) - y_i)^2 \quad (2)$$

$$= \frac{1}{2} \sum_{i=1}^m (\theta_0 + \theta_1 x_i - y_i)^2, \quad (3)$$

$$= \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^{\top} \theta - y_i)^2 \quad (4)$$

which is called residual sum of squares (RSS) or sum of squared errors (SSE).

Cost function

- We design the following **cost function** to minimize the difference between the prediction $h_{\theta}(x_i)$ and the ground-truth value y_i , i.e.,

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m (h_{\theta}(x_i) - y_i)^2 \quad (2)$$

$$= \frac{1}{2} \sum_{i=1}^m (\theta_0 + \theta_1 x_i - y_i)^2, \quad (3)$$

$$= \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^{\top} \theta - y_i)^2 \quad (4)$$

which is called **residual sum of squares** (RSS) or sum of squared errors (SSE).

- $J(\theta)$ is a **convex** or **non-convex** function? What is the shape of it?



- The linear regression is formulated to the following optimization problem

$$\boldsymbol{\theta}^* = \arg \min_{\boldsymbol{\theta}} J(\boldsymbol{\theta}) = \frac{1}{2} \sum_{i=1}^m \underbrace{(\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2}_{\text{red underline}}. \quad (5)$$

Gradient descent


- The linear regression is formulated to the following optimization problem

$$\theta^* = \arg \min_{\theta} J(\theta) = \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i)^2. \quad (5)$$

- θ can be updated by **gradient descent algorithm**,

$$\theta \leftarrow \theta - \alpha \frac{\partial J(\theta)}{\partial \theta}, \quad \frac{\partial J(\theta)}{\partial \theta} = \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i) \bar{x}_i \quad (6)$$

where α is called step-size or learning rate.



Gradient descent

- The linear regression is formulated to the following optimization problem

$$\boldsymbol{\theta}^* = \arg \min_{\boldsymbol{\theta}} J(\boldsymbol{\theta}) = \frac{1}{2} \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2. \quad (5)$$

- $\boldsymbol{\theta}$ can be updated by **gradient descent algorithm**,

$$\boldsymbol{\theta} \leftarrow \boldsymbol{\theta} - \alpha \frac{\partial J(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}}, \quad \frac{\partial J(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i) \bar{\mathbf{x}}_i \quad (6)$$

where α is called step-size or learning rate.

- Does gradient descent always converge to the optimal solution?

Gradient descent

- The linear regression is formulated to the following optimization problem

$$\theta^* = \arg \min_{\theta} J(\theta) = \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i)^2. \quad (5)$$

- θ can be updated by **gradient descent algorithm**,

$$\theta \leftarrow \theta - \alpha \frac{\partial J(\theta)}{\partial \theta}, \quad \frac{\partial J(\theta)}{\partial \theta} = \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i) \bar{x}_i \quad (6)$$

where α is called step-size or learning rate.

- Does gradient descent always converge to the optimal solution? (Plot the trajectory of gradient descent on curve or contours)

$m d^2$

Analytical solution

- If we set the gradient to 0, then we can get the following solution

$$\frac{\partial J(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i) \bar{\mathbf{x}}_i = \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta} - \mathbf{X}^\top \mathbf{y} = 0 \quad (7)$$

$$\Rightarrow \boldsymbol{\theta}^* = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}, \quad (8)$$

$$\mathbf{X} = \begin{bmatrix} \mathbf{x}_1^\top \\ \mathbf{x}_2^\top \\ \vdots \\ \mathbf{x}_m^\top \end{bmatrix}$$

Analytical solution

- If we set the gradient to 0, then we can get the following solution

$$\frac{\partial J(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i) \bar{\mathbf{x}}_i = \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta} - \mathbf{X}^\top \mathbf{y} = 0 \quad (7)$$

$$\Rightarrow \underline{\boldsymbol{\theta}^* = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}}, \quad (8)$$

which are called **normal equation** and **ordinary least squares** (OLS) solution, respectively. $\mathbf{X} = [\bar{\mathbf{x}}_1^\top; \bar{\mathbf{x}}_2^\top; \dots; \bar{\mathbf{x}}_m^\top] \in \mathbb{R}^{m \times d}$.

Analytical solution

- If we set the gradient to 0, then we can get the following solution

$$\frac{\partial J(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i) \bar{\mathbf{x}}_i = \mathbf{X}^\top \mathbf{X} \boldsymbol{\theta} - \mathbf{X}^\top \mathbf{y} = 0 \quad (7)$$

$$\Rightarrow \boldsymbol{\theta}^* = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}, \quad (8)$$

which are called **normal equation** and **ordinary least squares** (OLS) solution, respectively. $\mathbf{X} = [\bar{\mathbf{x}}_1^\top; \bar{\mathbf{x}}_2^\top; \dots; \bar{\mathbf{x}}_m^\top] \in \mathbb{R}^{m \times d}$.

- Since there is a closed-form solution, why do we need gradient descent algorithm?

Geometric interpretation

- Since $\theta^* = (\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}$, then the predictions of \mathbf{X} can be obtained by

$$\hat{\mathbf{y}} = \mathbf{X}\theta^* = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}, \quad (9)$$

which corresponds to the orthogonal projection of \mathbf{y} onto the column space of \mathbf{X} .

$$\mathbf{X} = [\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_d]$$

$$\alpha_1 \mathbf{x}_1 + \alpha_2 \mathbf{x}_2 + \dots + \alpha_d \mathbf{x}_d$$


Geometric interpretation

- Since $\theta^* = (X^\top X)^{-1} X^\top y$, then the predictions of X can be obtained by

$$\hat{\mathbf{y}} = \mathbf{X}\boldsymbol{\theta}^* = \mathbf{X}(\mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}, \quad (9)$$

which corresponds to the **orthogonal projection** of \mathbf{y} onto the column space of \mathbf{X} .

$$X = \begin{pmatrix} 1 & 2 \\ 1 & -2 \\ 1 & 2 \end{pmatrix}, \quad y = \begin{pmatrix} 8.89 \\ 0.61 \\ 1.77 \end{pmatrix}$$



Normal equation vs. gradient descent

	Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_0	x_1	x_2	x_3	x_4	y
1	2104	5	1	45	460
1	1416	3	2	40	232
1	1534	3	2	30	315
1	852	2	1	36	178

$X = \begin{bmatrix} 1 & 2104 & 5 & 1 & 45 \\ 1 & 1416 & 3 & 2 & 40 \\ 1 & 1534 & 3 & 2 & 30 \\ 1 & 852 & 2 & 1 & 36 \end{bmatrix}$
 $m \times (n+1)$

$y = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$
 m -dimensional vector

$\theta = (X^T X)^{-1} X^T y$

Normal equation vs. gradient descent

	Size (feet ²)	Number of bedrooms	Number of floors	Age of home (years)	Price (\$1000)
x_0	x_1	x_2	x_3	x_4	y
1	2104	5	1	45	460
1	1416	3	2	40	232
1	1534	3	2	30	315
1	852	2	1	36	178

$X = \begin{bmatrix} 1 & 2104 & 5 & 1 & 45 \\ 1 & 1416 & 3 & 2 & 40 \\ 1 & 1534 & 3 & 2 & 30 \\ 1 & 852 & 2 & 1 & 36 \end{bmatrix}$
 $y = \begin{bmatrix} 460 \\ 232 \\ 315 \\ 178 \end{bmatrix}$

$\theta = (X^T X)^{-1} X^T y$

$\theta \leftarrow \theta - \alpha \nabla_{\theta}$

$(n+1) \times (n+1)$

Gradient Descent	Normal Equation
Need to choose alpha	No need to <u>choose alpha</u>
Needs many iterations	No need to iterate
$O(kn^2)$	$O(n^3)$, need to calculate inverse of $X^T X$
Works well when n is large	Slow if n is very large

Probabilistic modeling

- We assume that the relationship between the input variable/feature \mathbf{x} and the output (variable) y is

$$\underbrace{y}_{\phi(\mathbf{x})} = \underbrace{\boldsymbol{\theta}^\top}_{\mathbf{w}} \underbrace{\mathbf{x}}_{\mathbf{z}} + \underbrace{e}_{\text{noise}}, \text{ where } e \sim \mathcal{N}(0, \sigma^2), \quad (10)$$

where e is called **observation noise** or **residual error**, and it is independent with any specific input \mathbf{x} .


Probabilistic modeling

- We assume that the relationship between the input variable/feature \mathbf{x} and the output variable y is

$$y = \theta^\top \mathbf{x} + e, \text{ where } e \sim \mathcal{N}(0, \sigma^2), \quad (10)$$

where e is called **observation noise** or **residual error**, and it is independent with any specific input \mathbf{x} .

- Thus, the output y can also be seen as a random variable, and its conditional probability is formulated as


$$p(y|\mathbf{x}, \theta) = \mathcal{N}(\theta^\top \mathbf{x}, \sigma^2) \quad (11)$$

Maximum log-likelihood estimation

- The parameter θ can be learned by maximum log-likelihood estimation (MLE), given the training dataset $D = \{(\mathbf{x}_i, y_i)\}_{i=1}^m$, as follows

$$\theta_{MLE} = \arg \max_{\theta} \log \mathcal{L}(\theta|D) \quad (12)$$

$$= \sum_i^m \log p(y|\mathbf{x}, \theta) = \sum_i^m \log \mathcal{N}(\theta^\top \mathbf{x}, \sigma^2) \quad (13)$$

$$= \underbrace{-\log(\sigma^m (2\pi)^{\frac{m}{2}})}_{\text{constant}} - \frac{1}{2\sigma^2} \sum_i^m (y_i - \theta^\top \mathbf{x}_i) \quad (14)$$

Maximum log-likelihood estimation

- The parameter θ can be learned by maximum log-likelihood estimation (MLE), given the training dataset $D = \{(\mathbf{x}_i, y_i)\}_{i=1}^m$, as follows

$$\theta_{MLE} = \arg \max_{\theta} \log \mathcal{L}(\theta|D) \quad (12)$$

$$= \sum_i^m \log p(y|\mathbf{x}, \theta) = \sum_i^m \log \mathcal{N}(\theta^\top \mathbf{x}, \sigma^2) \quad (13)$$

$$= -\log(\sigma^m (2\pi)^{\frac{m}{2}}) - \frac{1}{2\sigma^2} \sum_i^m (y_i - \theta^\top \mathbf{x}_i)^2 \quad (14)$$


- Removing the constants w.r.t. θ ,

$$\theta_{MLE} = \arg \min_{\theta} \underbrace{\frac{1}{2} \sum_i^m (y_i - \theta^\top \mathbf{x}_i)^2}_{\text{cost function}}, \quad (15)$$

which is exactly same with the cost function from the deterministic perspective.

Robust linear regression


- When there is a few outliers in the training data D , which are far from most other points, then learned parameters θ_{MLE} will be significantly influenced, leading to very poor fit.



Robust linear regression

- When there is a few outliers in the training data D , which are far from most other points, then learned parameters θ_{MLE} will be significantly influenced, leading to very poor fit.
- Let's see the loss curve of the residual sum of squares (RSS),

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i)^2. \quad (16)$$



Robust linear regression

- When there is a few outliers in the training data D , which are far from most other points, then learned parameters $\boldsymbol{\theta}_{MLE}$ will be significantly influenced, leading to very poor fit.
- Let's see the loss curve of the residual sum of squares (RSS),

$$J(\boldsymbol{\theta}) = \frac{1}{2} \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2. \quad (16)$$


- The error increases quadratically along with the residual. To minimize such a large error, the linear model will be significantly changed.

Robust linear regression

- When there is a few outliers in the training data D , which are far from most other points, then learned parameters θ_{MLE} will be significantly influenced, leading to very poor fit.
- Let's see the loss curve of the residual sum of squares (RSS),

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m (\bar{x}_i^\top \theta - y_i)^2. \quad (16)$$

- The error increases quadratically along with the residual. To minimize such a large error, the linear model will be significantly changed.
- How to alleviate the significant influence of outliers?





Robust linear regression

- When there is a few outliers in the training data D , which are far from most other points, then learned parameters θ_{MLE} will be significantly influenced, leading to very poor fit.
- Let's see the loss curve of the residual sum of squares (RSS),

$$J(\theta) = \frac{1}{2} \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \theta - y_i)^2. \quad (16)$$

- The error increases quadratically along with the residual. To minimize such a large error, the linear model will be significantly changed.
- How to alleviate the significant influence of outliers?



Robust linear regression

- We adopt the ℓ_1 loss to replace the ℓ_2 loss, as follows


$$J(\boldsymbol{\theta}) = \sum_{i=1}^m |\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i|. \quad (17)$$

Robust linear regression

- We adopt the ℓ_1 loss to replace the ℓ_2 loss, as follows

$$J(\theta) = \sum_{i=1}^m |\bar{\mathbf{x}}_i^\top \theta - y_i|. \quad (17)$$

- The curves of ℓ_1 and ℓ_2 losses are shown as follows.




Robust linear regression

- We adopt the ℓ_1 loss to replace the ℓ_2 loss, as follows

$$J(\boldsymbol{\theta}) = \sum_{i=1}^m |\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i|. \quad (17)$$

- The curves of ℓ_1 and ℓ_2 losses are shown ad follows.
- When the residual is large, the ℓ_1 loss is much smaller than the ℓ_2 loss, such that the influence of outliers could be alleviated.



Robust linear regression

- Actually, the above ℓ_1 loss can also be derived from the probabilistic perspective, by assuming that

$$p(y|\mathbf{x}, \boldsymbol{\theta}, b) = \text{Lap}(y|\mathbf{x}, \boldsymbol{\theta}, b) \propto \exp\left(-\frac{1}{b}|y - \boldsymbol{\theta}^\top \mathbf{x}|\right) \quad (18)$$

Robust linear regression



- Actually, the above ℓ_1 loss can also be derived from the probabilistic perspective, by assuming that

$$p(y|\mathbf{x}, \boldsymbol{\theta}, b) = \text{Lap}(y|\mathbf{x}, \boldsymbol{\theta}, b) \propto \exp(-\frac{1}{b}|y - \boldsymbol{\theta}^\top \mathbf{x}|) \quad (18)$$

- Applying the maximum log-likelihood estimation (MLE), we will obtain

$$\boldsymbol{\theta}_{MLE} = \arg \max_{\boldsymbol{\theta}} \log \mathcal{L}(\boldsymbol{\theta}|D) = \sum_i^m \log p(y_i|\mathbf{x}_i, \boldsymbol{\theta}) \quad (19)$$

$$\equiv \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i| \quad (20)$$



Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^{\top} \boldsymbol{\theta} - y_i| \quad (21)$$

Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^{\top} \boldsymbol{\theta} - y_i| \quad (21)$$

- However, the ℓ_1 loss function is non-differentiable and non-linear. The gradient descent algorithm cannot be adopted.

Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^{\top} \boldsymbol{\theta} - y_i| \quad (21)$$

- However, the ℓ_1 loss function is non-differentiable and non-linear. The gradient descent algorithm cannot be adopted.
- We can transform it to a linear program, as follows

Robust linear regression

$$\theta_{MLE} = \arg \min_{\theta} \sum_{i=1}^m \underbrace{|x_i^{\top} \theta - y_i|}_{\Delta} \quad (21)$$

- However, the ℓ_1 loss function is non-differentiable and non-linear. The gradient descent algorithm cannot be adopted.
- We can transform it to a linear program, as follows

$$\min_{\theta, t} \sum_{i=1}^m t_i = \min_{\theta, t} \mathbf{1}^{\top} t \quad (22)$$

Handwritten notes: A red arrow points from the t_i in the objective to the t_i in the constraint. Another red arrow points from the $|x_i^{\top} \theta - y_i| \leq t_i$ constraint to the t_i in the objective.

$$\text{s.t. } -t_i \leq \underbrace{x_i^{\top} \theta - y_i}_{\Delta} \leq t_i, 1 \leq i \leq m. \quad (23)$$

Handwritten notes: A red arrow points from the θ in the constraint to the θ in the objective. Another red arrow points from the t in the constraint to the t in the objective. The constraint is written as $[x_i^{\top} \theta - y_i] \leq t$.

Handwritten notes: The linear program is written as $Az \leq b$.


Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\boldsymbol{x}_i^\top \boldsymbol{\theta} - y_i| \quad (24)$$

Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^\top \boldsymbol{\theta} - y_i| \quad (24)$$

- We can also utilize the following equation:

$$|a| = \min_{\mu > 0} \left(\frac{a^2}{2\mu} + \mu \right) \quad (25)$$



Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^\top \boldsymbol{\theta} - y_i| \quad (24)$$

- We can also utilize the following equation:

$$|a| = \min_{\mu > 0} \frac{1}{2} \left(\frac{a^2}{\mu} + \mu \right) \quad (25)$$

- Then, the ℓ_1 minimization problem can be reformulated as follows

$$\min_{\boldsymbol{\theta}} \min_{\mu > 0} \frac{1}{2} \left(\frac{(\mathbf{x}^\top \boldsymbol{\theta} - y_i)^2}{\mu} + \mu \right). \quad (26)$$


Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^\top \boldsymbol{\theta} - y_i| \quad (24)$$

- We can also utilize the following equation:

$$|a| = \min_{\mu > 0} \frac{1}{2} \left(\frac{a^2}{\mu} + \mu \right) \quad (25)$$

- Then, the ℓ_1 minimization problem can be reformulated as follows

$$\min_{\boldsymbol{\theta}} \min_{\mu > 0} \frac{1}{2} \left(\frac{(\mathbf{x}^\top \boldsymbol{\theta} - y_i)^2}{\mu} + \mu \right). \quad (26)$$

- It can be iteratively and alternatively optimized as follows:

- Given $\boldsymbol{\theta}$, $\mu_i = |\mathbf{x}_i^\top \boldsymbol{\theta} - y_i|$
 - Given μ_i , $\boldsymbol{\theta} = \min_{\boldsymbol{\theta}} (\mathbf{x}_i^\top \boldsymbol{\theta} - y_i)^2$
- Handwritten notes:* A red circle around μ_i in the first bullet. A red arrow points from the μ_i in the second bullet to the μ_i in the first bullet. A red line is drawn under the second bullet, with the handwritten text "or μ_i " written below it.

Robust linear regression

$$\boldsymbol{\theta}_{MLE} = \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m |\mathbf{x}_i^\top \boldsymbol{\theta} - y_i| \quad (24)$$

- We can also utilize the following equation:

$$|a| = \min_{\mu > 0} \frac{1}{2} \left(\frac{a^2}{\mu} + \mu \right) \quad (25)$$

- Then, the ℓ_1 minimization problem can be reformulated as follows

$$\min_{\boldsymbol{\theta}} \min_{\mu > 0} \frac{1}{2} \left(\frac{(\mathbf{x}^\top \boldsymbol{\theta} - y_i)^2}{\mu} + \mu \right). \quad (26)$$

- It can be iteratively and alternatively optimized as follows:


- Given $\boldsymbol{\theta}$, $\mu = |\mathbf{x}^\top \boldsymbol{\theta} - y_i|$
- Given μ , $\boldsymbol{\theta} = \min_{\boldsymbol{\theta}} (\mathbf{x}^\top \boldsymbol{\theta} - y_i)^2$

- It is called iteratively reweighted least squares method.

(IRLS)

Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.




Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.
- What approaches we have introduced to alleviate overfitting?


Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.
- What approaches we have introduced to alleviate overfitting? Ocam's razor or cross-validation
- Is there other more theoretical approaches? SURE!




Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.
- What approaches we have introduced to alleviate overfitting? Ocam's razor or cross-validation
- Is there other more theoretical approaches? SURE!




Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.
- What approaches we have introduced to alleviate overfitting? Ocam's razor or cross-validation
- Is there other more theoretical approaches? SURE!




Ridge regression

- As demonstrated in the first week, overfitting is an important challenge for linear regression.
- What approaches we have introduced to alleviate overfitting? Ocam's razor or cross-validation
- Is there other more theoretical approaches? SURE!




Ridge regression



$$h_{\theta}(x) = \sum_{i=1}^{14} \alpha_i x^i - \frac{1}{7} x^{10}$$

- Let's see one simple example, we use a polynomial function with 14 degree to fit $m = 21$ data points. The learned curve is very “wiggly” (see above).


Ridge regression



- Let's see one simple example, we use a polynomial function with 14 degree to fit $m = 21$ data points. The learned curve is very “wiggly” (see above).
- The parameter values of this curve are as follows

6.56, -36.934, -109.25, 543.452, 1022.561, -3046.224, -3768.013, 8524.54,
 6607.897, -12640.058, -5530.188, 9479.73, 1774, 639, -2821.526

Ridge regression



- Let's see one simple example, we use a polynomial function with 14 degree to fit $m = 21$ data points. The learned curve is very “wiggly” (see above).
- The parameter values of this curve are as follows

6.56, -36.934, -109.25, 543.452, 1022.561, -3046.224, -3768.013, 8524.54,
6607.897, -12640.058, -5530.188, 9479.73, 1774, 639, -2821.526

- There are many large positive/negative values, such that a small change of features could lead to significant change of output.

Ridge regression

- How to get smaller parameter values?

Ridge regression

- How to get smaller parameter values?
- We can assume that the parameter follow a zero-mean Gaussian prior

$$p(\theta) = \mathcal{N}(\theta | \mathbf{0}, \tau^2 \mathbf{I}) \quad (27)$$



Ridge regression

- How to get smaller parameter values?
- We can assume that the parameter follow a zero-mean Gaussian prior

$$p(\boldsymbol{\theta}) = \mathcal{N}(\boldsymbol{\theta}|\mathbf{0}, \tau^2\mathbf{I}) \quad (27)$$

- Utilizing this prior, we obtain the maximum a posteriori (MAP) estimation

$$\boldsymbol{\theta}_{MAP} = \arg \max_{\boldsymbol{\theta}} \sum_i^m \log p(y|\mathbf{x}, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta}) \quad (28)$$

$$= \sum_i^m \log \mathcal{N}(\boldsymbol{\theta}^\top \mathbf{x}, \sigma^2) + \mathcal{N}(\boldsymbol{\theta}|\mathbf{0}, \tau^2\mathbf{I}) \quad (29)$$

$$\equiv \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2 + \lambda \|\boldsymbol{\theta}\|_2^2. \quad (30)$$

loss

regularization

Ridge regression

- How to get smaller parameter values?
- We can assume that the parameter follow a zero-mean Gaussian prior

$$p(\boldsymbol{\theta}) = \mathcal{N}(\boldsymbol{\theta}|\mathbf{0}, \tau^2\mathbf{I}) \quad (27)$$

- Utilizing this prior, we obtain the maximum a posteriori (MAP) estimation

$$\boldsymbol{\theta}_{MAP} = \arg \max_{\boldsymbol{\theta}} \sum_i^m \log p(y|\mathbf{x}, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta}) \quad (28)$$

$$= \sum_i^m \log \mathcal{N}(\boldsymbol{\theta}^\top \mathbf{x}, \sigma^2) + \mathcal{N}(\boldsymbol{\theta}|\mathbf{0}, \tau^2\mathbf{I}) \quad (29)$$

$$\equiv \arg \min_{\boldsymbol{\theta}} \left(\sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2 + \lambda \|\boldsymbol{\theta}\|_2^2 \right) \quad (30)$$

- The corresponding closed-form solution is given by

$$\boldsymbol{\theta}_{MAP} = (\lambda \mathbf{I} + \mathbf{X}^\top \mathbf{X})^{-1} \mathbf{X}^\top \mathbf{y}. \quad (31)$$


Ridge regression


- The above method is also known as **ridge regression**, or penalized least squares.

Ridge regression

- The above method is also known as **ridge regression**, or **penalized least squares**.
- In general, adding a Gaussian prior to the parameters of a model to encourage them to be small is called ℓ_2 **regularization** or **weight decay**.

Ridge regression

- The above method is also known as **ridge regression**, or **penalized least squares**.
- In general, adding a Gaussian prior to the parameters of a model to encourage them to be small is called ℓ_2 **regularization** or **weight decay**.
- As shown below, when we set a larger λ , *i.e.*, more weight on the prior, the resulting curve will be smoother. 



Lasso regression

- We can replace the Gaussian prior by a Laplacian prior, *i.e.*,

$$p(\boldsymbol{\theta}) = \text{Lap}(\boldsymbol{\theta}|\mathbf{0}, b) = \frac{1}{2b} \exp\left(-\frac{|\boldsymbol{\theta}|}{b}\right), \quad (32)$$

Lasso regression

- We can replace the Gaussian prior by a Laplacian prior, *i.e.*,

$$p(\boldsymbol{\theta}) = \text{Lap}(\boldsymbol{\theta}|\mathbf{0}, b) = \frac{1}{2b} \exp\left(-\frac{|\boldsymbol{\theta}|}{b}\right), \quad (32)$$

- The combination of the Gaussian distribution of $p(y|\mathbf{x}, \boldsymbol{\theta})$ and the Laplacian prior, leading to

$$\boldsymbol{\theta}_{MAP} = \arg \max_{\boldsymbol{\theta}} \sum_i^m \log p(y|\mathbf{x}, \boldsymbol{\theta}) + \log p(\boldsymbol{\theta}) \quad (33)$$

$$= \sum_i^m \log \mathcal{N}(\boldsymbol{\theta}^\top \mathbf{x}, \sigma^2) + \text{Lap}(\boldsymbol{\theta}|\mathbf{0}, b) \quad (34)$$


$$\equiv \arg \min_{\boldsymbol{\theta}} \sum_{i=1}^m (\bar{\mathbf{x}}_i^\top \boldsymbol{\theta} - y_i)^2 + \lambda |\boldsymbol{\theta}|. \quad (35)$$

Lasso regression

- It is **Lasso regression**, and the regularization is called ℓ_1 **regularization**. It will encourage the sparse parameters.


Lasso regression

- It is **Lasso regression**, and the regularization is called ℓ_1 **regularization**. It will encourage the sparse parameters.
- As shown below, when we set a larger λ , *i.e.*, more weight on the prior, the resulting curve will be smoother.



Geometry of Ridge and Lasso regression

- Geometry of Ridge and Lasso regression. Which one is Ridge?




Summary of different linear regressions

Note that the uniform distribution will not change the mode of the likelihood.

Thus, MAP estimation with a uniform prior corresponds to MLE.

$p(y \mathbf{x}, \boldsymbol{\theta})$	$p(\boldsymbol{\theta})$	regression method
Gaussian	Uniform	Least squares
Gaussian	Gaussian	Ridge regression
Gaussian	Laplace	Lasso regression
Laplace	Uniform	Robust regression
Student	Uniform	Robust regression



Generalized linear regression

- **Linear model:**

$$\mu(\mathbf{x}|\boldsymbol{\theta}) = \boldsymbol{\theta}^\top \phi(\mathbf{x}), \quad (36)$$

$$y(\mathbf{x}|\boldsymbol{\theta}) \sim f(\mu(\mathbf{x}|\boldsymbol{\theta})), \quad (37)$$

where f denotes a distribution function.

Generalized linear regression

- **Linear model:**

$$\mu(\mathbf{x}|\boldsymbol{\theta}) = \boldsymbol{\theta}^\top \phi(\mathbf{x}), \quad (36)$$

$$y(\mathbf{x}|\boldsymbol{\theta}) \sim f(\mu(\mathbf{x}|\boldsymbol{\theta})), \quad (37)$$

where f denotes a distribution function.

- **Generalized linear model (GLM):**

$$\mu(\mathbf{x}|\boldsymbol{\theta}) = g^{-1}(\boldsymbol{\theta}^\top \phi(\mathbf{x})), \quad (38)$$

$$y(\mathbf{x}|\boldsymbol{\theta}) \sim f(\mu(\mathbf{x}|\boldsymbol{\theta})), \quad (39)$$

where g is called **link function**, which is required to be monotonically increasing differentiable.

Generalized linear regression

- **Linear model:**

$$\mu(\mathbf{x}|\boldsymbol{\theta}) = \boldsymbol{\theta}^\top \phi(\mathbf{x}), \quad (36)$$

$$y(\mathbf{x}|\boldsymbol{\theta}) \sim f(\mu(\mathbf{x}|\boldsymbol{\theta})), \quad (37)$$

where f denotes a distribution function.

- **Generalized linear model (GLM):**

$$\mu(\mathbf{x}|\boldsymbol{\theta}) = g^{-1}(\boldsymbol{\theta}^\top \phi(\mathbf{x})), \quad (38)$$

$$y(\mathbf{x}|\boldsymbol{\theta}) \sim f(\mu(\mathbf{x}|\boldsymbol{\theta})), \quad (39)$$

where g is called **link function**, which is required to be monotonically increasing differentiable.

- The standard linear model is a special case of GLM with $g(a) = a$.

Why we need generalized linear regression

- **Why we need generalized linear model?** Let's see one example.

In the early stages of a disease epidemic, the rate at which new cases occur can often increase exponentially through time. Hence, if μ_i is the expected number of new cases on day t_i , a model of the form

$$\mu_i = \gamma \exp(\delta t_i)$$

seems appropriate.

- ▶ Such a model can be turned into GLM form, by using a **log link** so that

$$\log(\mu_i) = \log(\gamma) + \delta t_i = \beta_0 + \beta_1 t_i.$$

- ▶ Since this is a count, the **Poisson distribution** (with expected value μ_i) is probably a reasonable distribution to try.

Log linear regression


- **Poisson distribution** The Poisson distribution is popular for modeling the number of times an event occurs in an interval of time or space.

Log linear regression

- **Poisson distribution** The Poisson distribution is popular for modeling the number of times an event occurs in an interval of time or space.
- A discrete random variable X is said to have a Poisson distribution with parameter $\lambda > 0$ if for $k = 0, 1, 2, \dots$, the probability mass function of X is given by

$$f(k; \lambda) = P(X = k | \lambda) = \frac{\lambda^k e^{-\lambda}}{k!}, \quad (40)$$

where e is Euler's number ($e = 2.71828\dots$), k is the number of occurrences, $k!$ is the factorial of k .



Log linear regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \textit{Poisson}(\lambda_i) = \frac{\lambda^{y_i} e^{-\lambda}}{y_i!}, \quad \ln \lambda_i = \boldsymbol{\theta}^\top \mathbf{x}_i \quad (41)$$

Log linear regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \text{Poisson}(\lambda_i) = \frac{\lambda^{y_i} e^{-\lambda}}{y_i!}, \quad \ln \lambda_i = \boldsymbol{\theta}^\top \mathbf{x}_i \quad (41)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \sum_{i=1}^m y_i \log \lambda_i - \lambda_i - \log y_i! \quad (42)$$

Log linear regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \text{Poisson}(\lambda_i) = \frac{\lambda_i^{y_i} e^{-\lambda_i}}{y_i!}, \quad \ln \lambda_i = \boldsymbol{\theta}^\top \mathbf{x}_i \quad (41)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \sum_{i=1}^m y_i \log \lambda_i - \lambda_i - \log y_i! \quad (42)$$

- We have

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (y_i \mathbf{x}_i - e^{\boldsymbol{\theta}^\top \mathbf{x}_i}) = 0 \quad \Rightarrow \quad \ln y_i = (\boldsymbol{\theta}^*)^\top \mathbf{x}_i \quad (43)$$

Log linear regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \text{Poisson}(\lambda_i) = \frac{\lambda^{y_i} e^{-\lambda}}{y_i!}, \quad \ln \lambda_i = \boldsymbol{\theta}^\top \mathbf{x}_i \quad (41)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = \sum_{i=1}^m y_i \log \lambda_i - \lambda_i - \log y_i! \quad (42)$$

- We have

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (y_i \mathbf{x}_i - e^{\boldsymbol{\theta}^\top \mathbf{x}_i}) = 0 \quad \Rightarrow \quad \ln y_i = (\boldsymbol{\theta}^*)^\top \mathbf{x}_i \quad (43)$$

- Plot the log-linear regression as below.

Logistic regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}, N) = \text{Bin}(y_i|N, \mu_i) = \binom{N}{y_i} \mu_i^{y_i} (1 - \mu_i)^{N-y_i}, \quad \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (44)$$

Logistic regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}, N) = \text{Bin}(y_i|N, \mu_i) = \binom{N}{y_i} \mu_i^{y_i} (1 - \mu_i)^{N-y_i}, \quad \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (44)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = y_i \log \mu_i + (N - y_i) \log(1 - \mu_i) \quad (45)$$

Logistic regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}, N) = \text{Bin}(y_i|N, \mu_i) = \binom{N}{y_i} \mu_i^{y_i} (1 - \mu_i)^{N-y_i}, \quad \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (44)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = y_i \log \mu_i + (N - y_i) \log(1 - \mu_i) \quad (45)$$

- We have

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (y_i - N\mu_i) \mathbf{x}_i = 0 \quad \Rightarrow \quad \frac{y_i}{N} = \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (46)$$

Logistic regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}, N) = \text{Bin}(y_i|N, \mu_i) = \binom{N}{y_i} \mu_i^{y_i} (1 - \mu_i)^{N-y_i}, \quad \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (44)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = y_i \log \mu_i + (N - y_i) \log(1 - \mu_i) \quad (45)$$

- We have

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (y_i - N\mu_i) \mathbf{x}_i = 0 \quad \Rightarrow \quad \frac{y_i}{N} = \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (46)$$

- Since the $\sigma(a) = \frac{1}{1+e^{-a}}$ is called **sigmoid function** or **logit function**, the above model is called **logit regression** or **logistic regression**.

Logistic regression

- We assume that the conditional probability follows

$$P(y_i|\mathbf{x}_i, \boldsymbol{\theta}, N) = \text{Bin}(y_i|N, \mu_i) = \binom{N}{y_i} \mu_i^{y_i} (1 - \mu_i)^{N-y_i}, \quad \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (44)$$

- The log-likelihood function is formulated as follows

$$\mathcal{L}(\boldsymbol{\theta}) = \sum_{i=1}^m \log P(y_i|\mathbf{x}_i, \boldsymbol{\theta}) = y_i \log \mu_i + (N - y_i) \log(1 - \mu_i) \quad (45)$$

- We have

$$\frac{\partial \mathcal{L}(\boldsymbol{\theta})}{\partial \boldsymbol{\theta}} = \sum_{i=1}^m (y_i - N\mu_i) \mathbf{x}_i = 0 \quad \Rightarrow \quad \frac{y_i}{N} = \mu_i = \frac{1}{1 + e^{-\boldsymbol{\theta}^\top \mathbf{x}_i}}. \quad (46)$$

- Since the $\sigma(a) = \frac{1}{1+e^{-a}}$ is called **sigmoid function** or **logit function**, the above model is called **logit regression** or **logistic regression**.
- Since $\frac{y_i}{N} \in [0, 1]$, it can be seen as the posterior probability. Thus, logistic regression is a classification model, rather than regression.

Summary

- Linear model is the linear function of the parameter θ , rather than the input feature

Summary

- Linear model is the linear function of the parameter θ , rather than the input feature
- Linear model is a special case of generalized linear model, while generalized linear model is not always linear

- Linear model is the linear function of the parameter θ , rather than the input feature
- Linear model is a special case of generalized linear model, while generalized linear model is not always linear
- Choosing different linear models is equivalent to choosing different distributions of $p(y|\mathbf{x}, \theta)$ and $p(\theta)$, according to the task and the data

Reading material

- <https://www.stat.cmu.edu/~ryantibs/advmethods/notes/glm.pdf>