



Synthesis of data products for ocean carbonate chemistry

Li-Qing Jiang^{1, 2}, Amanda Fay³, Jens Daniel Müller⁴, Lydia Keppler⁵, Dustin Carroll^{6, 7}, Siv K. Lauvset⁸, Tim DeVries⁹, Judith Hauck^{10, 11}, Christian Rödenbeck¹², Luke Gregor^{4, 13}, Nicolas Metzl¹⁴, Andrea J. Fassbender¹⁵, Jean-Pierre Gattuso^{16, 17}, Peter Landschützer¹⁸, Rik Wanninkhof¹⁹, Christopher Sabine²⁰, Simone R. Alin¹⁵, Mario Hoppema²¹, Are Olsen²², Matthew P. Humphreys²³, Kumiko Azetsu-Scott²⁴, Dorothee C. E. Bakker²⁵, Leticia Barbero^{19, 26}, Nicholas R. Bates^{27, 28}, Nicole Besemer¹⁹, Henry C. Bittig²⁹, Albert E. Boyd^{19, 26}, Daniel Broullón³⁰, Wei-Jun Cai³¹, Brendan R. Carter³², Thi-Tuyet-Trang Chau^{33, 34}, Chen-Tung Arthur Chen³⁵, Frédéric Cyr^{36, 37}, John E. Dore³⁸, Ian Enochs¹⁹, Richard A. Feely¹⁵, Hernan E. Garcia², Marion Gehlen³³, Lucas Gloege³, Melchor González-Dávila³⁹, Nicolas Gruber⁴, Yosuke Iida⁴⁰, Masao Ishii⁴¹, Esther Kennedy⁴², Alex Kozyr^{1, 2}, Nico Lange⁸, Claire Lo Monaco¹⁴, Derek P. Manzello⁴³, Galen A. McKinley³, Natalie M. Monacci⁴⁴, Xose A. Padin³⁰, Ana M. Palacio-Castro^{19, 26}, Fiz F. Pérez³⁰, Alizée Roobaert⁴⁵, J. Magdalena Santana-Casiano³⁹, Jonathan Sharp^{15, 46}, Adrienne Sutton¹⁵, Jim Swift⁴⁷, Toste Tanhua⁴⁸, Maciej Telszewski⁴⁹, Jens Terhaar^{50, 51}, Ruben van Hooidonk⁵², Anton Velo³⁰, Andrew J. Watson⁵³, Angelique E. White²⁰, Zelun Wu³¹, Hyelim Yoo^{1, 2}, Jiye Zeng⁵⁴.

¹Cooperative Institute for Satellite Earth System Studies, Earth System Science Interdisciplinary Center, University of Maryland, College Park, Maryland 20740, United States.

²NOAA/NESDIS National Centers for Environmental Information, Silver Spring, Maryland 20910, United States.

³Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York 10964, United States.

⁴Environmental Physics, Institute of Biogeochemistry and Pollutant Dynamics, ETH Zürich, Zürich, Switzerland.

⁵Vycarb Inc., Brooklyn, New York, United States.

⁶Moss Landing Marine Laboratories, San José State University, Moss Landing, California, United States.

⁷Jet Propulsion Laboratory, California Institute of Technology, Pasadena, California, United States.

⁸NORCE Norwegian Research Centre, Bjerknes Centre for Climate Research, Bergen, Norway.

⁹Department of Geography and Earth Research Institute, University of California, Santa Barbara, California, United States.

¹⁰Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany

¹¹University of Bremen, Bremen, Germany.

¹²Max Planck Institute for Biogeochemistry, Jena, Germany.

¹³Swiss Data Science Center, ETH Zurich and EPFL, Zurich, Switzerland

¹⁴Laboratoire LOCEAN/IPSL, Sorbonne Université-CNRS-IRD-MNHN, 75005 Paris, France.

¹⁵NOAA/OAR Pacific Marine Environmental Laboratory, Seattle, Washington 98115, United States.

¹⁶Sorbonne Université, CNRS, Laboratoire d'Océanographie de Villefranche, 06230 Villefranche-sur-Mer, France.

¹⁷Institute for Sustainable Development and International Relations, Paris, France.

¹⁸Flanders Marine Institute (VLIZ), Jacobsenstraat 1, 8400 Ostend, Belgium.

¹⁹NOAA/OAR Atlantic Oceanographic and Meteorological Laboratory, Miami, Florida 33149, United States.

²⁰Department of Oceanography, University of Hawai'i at Mānoa, 1000 Pope Rd., Honolulu, Hawaii 96822, United States.

²¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Bremerhaven, Germany.

²²Geophysical Institute, University of Bergen and Bjerknes Centre for Climate Research, Bergen, Norway.

²³Department of Ocean Systems, NIOZ Royal Netherlands Institute for Sea Research, Texel, the Netherlands.



- 45 ²⁴Fisheries and Oceans, Canada, Bedford Institute of Oceanography, Dartmouth, Nov Scotia, Canada.
- 46 ²⁵Centre for Ocean and Atmospheric Sciences, School of Environmental Sciences, University of East Anglia,
47 Norwich NR4 7TJ, United Kingdom.
- 48 ²⁶Cooperative Institute for Marine and Atmospheric Studies, Rosenstiel School of Marine and Atmospheric Science,
49 University of Miami, Miami, Florida 33149, United States.
- 50 ²⁷Bermuda Institute of Ocean Sciences, ASU-BIOS, 17 Biological Lane, St. Georges, GEO1, Bermuda.
- 51 ²⁸School of Ocean Futures, Arizona State University, Tempe, Arizona 85281, United States.
- 52 ²⁹Leibniz Institute for Baltic Sea Research Warnemünde (IOW), Rostock, Germany.
- 53 ³⁰Instituto de Investigaciones Mariñas, IIM - CSIC, Vigo, Spain.
- 54 ³¹School of Marine Science and Policy, University of Delaware, Newark, Delaware 19716, United States.
- 55 ³²Independent Researcher.
- 56 ³³Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL, CEA-CNRS-UVSQ, Université Paris-
57 Saclay, 91191 Gif-sur-Yvette, France.
- 58 ³⁴Laboratoire Interdisciplinaire des Sciences du Numérique (LISN), Centre de recherches INRIA, Université Paris-
59 Saclay, René Thom, F-91190, Gif-sur-Yvette, France.
- 60 ³⁵Department of Oceanography, National Sun Yat-Sen University, Kaohsiung, Taiwan.
- 61 ³⁶Northwest Atlantic Fisheries Centre, Fisheries and Oceans Canada, St. John's, Newfoundland and Labrador,
62 Canada.
- 63 ³⁷Current affiliation: Centre for Fisheries Ecosystems Research, Fisheries and Marine Institute of Memorial
64 University of Newfoundland, St. John's, Canada.
- 65 ³⁸Department of Land Resources and Environmental Sciences, Montana State University, Bozeman, Montana,
66 United States.
- 67 ³⁹Instituto de Oceanografía y Cambio Global, IOCAG, Universidad de Las Palmas de Gran Canaria, 35017 Las
68 Palmas de Gran Canaria, Spain.
- 69 ⁴⁰Atmosphere and Ocean Department, Japan Meteorological Agency, 3-6-9 Toranomon, Minato City, Tokyo, Japan.
- 70 ⁴¹Meteorological Research Institute, Japan Meteorological Agency, Tsukuba, Japan.
- 71 ⁴²University of California Davis, Bodega Marine Laboratory, Davis, California, United States.
- 72 ⁴³Coral Reef Watch, U.S. National Oceanic and Atmospheric Administration, College Park, Maryland, United
73 States.
- 74 ⁴⁴College of Fisheries and Ocean Sciences, University of Alaska Fairbanks, Fairbanks, Alaska, United States.
- 75 ⁴⁵Flanders Marine Institute (VLIZ), Jacobsenstraat 1, 8400 Ostend, Belgium.
- 76 ⁴⁶Cooperative Institute for Climate, Ocean, and Ecosystem Studies, University of Washington, Seattle, Washington,
77 United States.
- 78 ⁴⁷Scripps Institution of Oceanography, La Jolla, California, United States.
- 79 ⁴⁸GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany.
- 80 ⁴⁹International Ocean Carbon Coordination Project, Institute of Oceanology of Polish Academy of Sciences, Sopot,
81 Poland.
- 82 ⁵⁰Climate and Environmental Physics, Physics Institute, University of Bern, Bern, Switzerland.
- 83 ⁵¹Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland.
- 84 ⁵²Science for Climate Change Resilience, Boulder, Colorado 80302, United States.
- 85 ⁵³Global Systems Institute, University of Exeter, Exeter, United Kingdom.
- 86 ⁵⁴National Institute for Environmental Studies, Tsukuba, Japan.
- 87 *Correspondence to:* Li-Qing Jiang (liqing.jiang@noaa.gov)



88 **Abstract.** As the largest active carbon reservoir on Earth, the ocean is a cornerstone of the global carbon cycle,
89 playing a pivotal role in modulating ocean health and regulating climate. Understanding these crucial roles requires
90 access to a broad array of data products documenting the changing chemistry of the global ocean as a vast and
91 interconnected system. This review article provides a comprehensive overview of 60 existing ocean carbonate
92 chemistry data products, encompassing compilations of cruise datasets, derived gap-filled data products, model
93 simulations, and compilations thereof. It is intended to help researchers identify and access data products that best
94 align with their research objectives, thereby advancing our understanding of the ocean's evolving carbonate
95 chemistry.



96 1 Introduction

97 Since the onset of the Industrial Revolution in 1750, human activities, such as the burning of fossil fuels, cement
 98 production, and land-use change, have emitted ~2600 Gt carbon dioxide (CO₂) into the atmosphere, causing the
 99 atmospheric CO₂ levels to increase by ~50% (DeVries, 2022; Friedlingstein et al., 2025; Tans and Keeling, 2025).
 100 The global carbon cycle, encompassing the exchange of CO₂ among the atmosphere, oceans, terrestrial ecosystems,
 101 and geosphere, plays a critical role in regulating atmospheric CO₂ levels (Archer, 2010; DeVries, 2022; Holzer and
 102 DeVries, 2022). As the largest dynamic CO₂ reservoir, the ocean holds approximately 45 times the amount of
 103 carbon found in the atmosphere currently and actively exchanges it with the air above and sediments below. On
 104 timescales from decades to millennia, the ocean imposes a dominant control over atmospheric CO₂ levels (Revelle
 105 and Suess, 1957; Broecker, 1982; Archer et al., 2009; DeVries, 2022).

106 The ocean currently absorbs about a quarter of human-caused CO₂ emissions (Sabine et al., 2004; Gruber et al.,
 107 2019a; Carroll et al., 2022; Crisp et al., 2022; Terhaar et al., 2022a; Gruber et al., 2023; DeVries et al., 2023; Müller
 108 et al., 2023; Schimel and Carroll, 2024). The chemistry of the ocean has been shifting as a result of atmospheric CO₂
 109 uptake (Feely et al., 2023; Ma et al., 2023; Müller et al., 2023; Fassbender et al., 2023; Keppler et al., 2023; Jiang et
 110 al., 2023; Müller and Gruber, 2024). Since the beginning of the Industrial Revolution, the dissolved inorganic
 111 carbon (DIC) content in the surface ocean has risen from 1690 to 1730 Gt of Carbon, and in the subsurface ocean
 112 from 35,400 to 35,560 Gt C (Sabine et al., 2004; Müller et al., 2023). This seemingly small increase of 0.5% belies a
 113 significant depletion of the oceans' buffer capacity (DeVries, 2022).

114 As CO₂ enters seawater, a portion of it reacts with water to form carbonic acid. This is the first in a series of
 115 rapid acid-base reactions that release protons and decrease the availability of carbonate ions, which are building
 116 materials that many marine organisms, such as mollusks, crustaceans, and corals, use to construct their shells and
 117 skeletons (Gattuso and Hansson, 2011). This process, termed as “ocean acidification (OA)”, has already decreased
 118 surface ocean pH by roughly 0.11 (~30% increase in acidity) since 1750 (Orr et al., 2005; Jiang et al., 2019;
 119 Kwiatkowski et al., 2020; Jiang et al., 2023; IPCC, 2023). In the subsurface ocean, the trends of some acidification
 120 parameters, e.g., pH and hydrogen ion content ([H⁺]), can be even greater due to the increasing sensitivity of [H⁺] to
 121 changes in DIC with depth (Chen et al., 2017; Fassbender et al., 2023; Müller and Gruber, 2024). This ongoing
 122 acidification threatens critical ocean ecosystem services, including food security, fisheries, aquaculture, and the
 123 broader Blue Economy, for billions of people globally (Cooley and Doney, 2009; Doney et al., 2020).

124 In some parts of the ocean, OA is driven not only by the uptake of carbon but also by other processes (Delaigue
 125 et al., 2024), for example via alkalinity changes driven by freshening of the Arctic Ocean (Terhaar et al., 2021a) or
 126 changes in the carbon and alkalinity export from Arctic rivers (Terhaar et al., 2019; Qi et al., 2022; Bertin et al.
 127 2023). Local anthropogenic inputs through rivers or from air pollution also lead to OA (e.g. Sarma et al., 2015;
 128 Sridvi and Sarma, 2021). In addition, coastal eutrophication and hypoxia could cause enhanced OA in oxygen-
 129 depleted bottom water due to weakened seawater buffer capacity by biologically induced CO₂ (Cai et al. 2011). If
 130 anthropogenic CO₂ emissions continue without mitigation, as per the shared socioeconomic pathway (SSP5-8.5)



scenario, surface ocean pH could decrease by a further 0.3 to 0.4 by 2100, equivalent to a 100–150% increase in acidity (Kwiatkowski et al., 2020; Jiang et al., 2023). If society, however, succeeds at reducing emissions, the future acidity level becomes highly uncertain as it sensitively depends on the transient response of the Earth system and the amount of reductions of non-CO₂ radiative agents (Terhaar et al., 2023).

In summary, monitoring ocean carbonate chemistry is essential for (a) tracking the evolving ocean carbon sink, and (b) understanding ocean acidification and its ecological impacts. Additionally, monitoring ocean carbonate chemistry is crucial when considering marine carbon dioxide removal (mCDR) strategies such as ocean alkalinity enhancement (OAE), artificial upwelling, ocean fertilization and electrochemical ocean CO₂ removal (Kheshgi, 1995; Bach et al., 2019; Schimel and Carroll, 2024; Oschlies et al., 2025). The ocean's vast and interconnected nature necessitates that data from individual oceanographic cruises be meticulously preserved, subject to rigorous quality control, and uniformly formatted to promote their usability (Brett et al., 2020; Schoderer et al., 2024). Following the methodology established by Lange et al. (2023), we have curated an exhaustive catalogue of synthesis products pertaining to ocean carbonate chemistry, including cruise data compilations, gridded gap-filled data products, and other derived data products. This compilation spans both global and regional scales, providing a holistic view of the current state of ocean biogeochemistry data aggregation.

2 Methods

In this paper, data products are defined as outputs that aggregate, quality-control, and transform individual datasets from multiple sources into a unified, structured format to support research, decision-making, or operational needs for specific end users. The data products included in this study were identified through a literature review and discussions with researchers via the Ocean Acidification Information Exchange.

3 Results and Discussion

3.1 Data products for ocean carbonate chemistry

The aim of establishing this catalog of ocean carbonate chemistry data synthesis products is to enhance user access and raise awareness of available resources. The list will be regularly updated to include the latest data synthesis products. Contributors are encouraged to submit new data products through the Ocean Acidification Information Exchange (OAIE) interface: <https://www.oainfoexchange.org/members/updates/71585>. If preferred, submissions can also be directed via email to noaa.ocads@noaa.gov.

These products are organized based on end-user needs:

1. Cruise data compilations (no interpolation or extrapolation).
2. Time-series data products (no interpolation or extrapolation).
3. Derived gap-filled (interpolated) products for the surface ocean, starting with products offering a



- 162 climatological snapshot of the ocean, followed by those showing temporal changes.
- 163 4. Derived gap-filled (interpolated) products for the interior ocean, also starting with products offering a
- 164 climatological snapshot of the ocean, followed by those showing temporal changes.
- 165 5. Multi-product analyses of 3 and 4. These compilations also include hindcast model simulations of the
- 166 ocean carbon cycle and biogeochemistry.
- 167 6. Model and hybrid data products projecting ocean carbon variables into the future [Note: Here the term
- 168 “model” refers to ocean biogeochemical models (Fennel et al., 2022). If a statistical model or machine
- 169 learning model is used for gap filling, the product is not categorized as a model output product in this
- 170 compilation.]

171 Each section includes numbered descriptions of each data product in that class, as well as a summary table of the

172 data products with corresponding IDs so the user can easily jump to the associated product description. Although

173 some data products, such as Surface Ocean CO₂ Atlas (SOCAT) and Lamont-Doherty Earth Observatory (LDEO)

174 surface partial pressure of carbon dioxide ($p\text{CO}_2$) Database report only one ocean carbonate variable, i.e., fugacity of

175 carbon dioxide ($f\text{CO}_2$), they provide a foundation from which additional variables can be derived using empirical

176 algorithms. For instance, total alkalinity content (TA) can be estimated from salinity and temperature and other

177 factors (Lee et al., 2006) and by neural network approaches such as those developed by Velo et al. (2013) and

178 Broullón et al. (2019). Beyond TA, neural network algorithms have been extended to estimate dissolved inorganic

179 carbon (DIC) as demonstrated by Broullón et al. (2020), and even the full marine carbonate system (MCS) through

180 frameworks like CANYON-B/CONTENT (Bittig et al., 2018) and Empirical Seawater Property Estimation Routines

181 (ESPERs) (Carter et al., 2021). While these methods primarily employ neural networks, both Velo et al. (2013) and

182 Carter et al. (2021) provide alternative estimation approaches based on local interpolation, through their 3-

183 dimensional moving window multilinear regression algorithm (3DwMLR) and locally interpolated regression (LIR)

184 methods, respectively. Utilizing such derived data, the complete suite of ocean carbonate parameters can then be

185 calculated using computer software, such as CO2SYS (Lewis and Wallace, 1998; van Heuven et al., 2011; Orr et al.,

186 2018; Sharp et al., 2023) or its Python implementation PyCO2SYS (Humphreys et al., 2022). An in-depth

187 explanation of the methods employed for these calculations can be found in the Supplementary material of Jiang et

188 al. (2022a).

189 **3.1.1 Cruise data compilations (no interpolation or extrapolation):**

190 The data compilations described in this section standardize datasets collected from individual research vessels,

191 ships of opportunity (SOOP), and unmanned platforms, presenting them in a uniform format for easy access. These

192 datasets typically undergo both primary QC (identifying outliers and obvious errors within an individual cruise

193 dataset) and secondary QC (objectively comparing data from one cruise against another or a previously synthesized

194 dataset to quantify systematic differences in reported values).



- 195 **1) SOCAT:** The Surface Ocean CO₂ Atlas (SOCAT) represents the most extensive collection of observational
 196 ocean CO₂ data for the global surface ocean (Bakker et al., 2016). It features fugacity of carbon dioxide
 197 (*f*CO₂) measurements from both the open ocean and the coastal ocean, predominantly sourced from
 198 research vessels, SOOP, and autonomous platforms including fixed moorings and autonomous surface
 199 vehicles (ASVs). Since 2013, SOCAT has been updated annually, and assigned dataset flags indicating
 200 uncertainty and completeness of metadata. SOCAT is also available as a gridded product. To access the
 201 latest version of the SOCAT data product (with 40 million data points), visit <https://socat.info/>.

- 202 **2) LDEO Surface *p*CO₂ Database:** Dr. Taro Takahashi [Lamont Doherty Earth Observatory (LDEO),
 203 Palisades, New York] started synthesizing global surface ocean CO₂ data in 1997, compiling three decades
 204 of observations (~250,000 measurements) to create inaugural monthly global surface *p*CO₂ maps
 205 (Takahashi et al., 1997; Takahashi et al., 2002). The most recent version (V2019) expanded this dataset to
 206 approximately 14.2 million surface water *p*CO₂ measurements spanning from 1957–2019. Distinct from the
 207 SOCAT database, the LDEO database reports *p*CO₂, instead of *f*CO₂, exclusively from equilibrator-CO₂
 208 analyzer systems, with an average estimated uncertainty of ± 2.5 μatm. The database is also interpolated
 209 onto a global surface ocean 4° x 5° grid for a reference year 2000 (Takahashi et al., 2009) and 2010 (Fay et
 210 al. 2024). Access to the LDEO surface *p*CO₂ database (Version 2019) is provided by the Ocean Carbon and
 211 Acidification Data System (OCADS) with the DOI: “10.3334/CDIAC/otg.ndp088(v2015)”, accessible at:
 212 <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0160492.html>. Additionally, there is a
 213 dedicated webpage at OCADS for the LDEO Database: [https://www.ncei.noaa.gov/access/ocean-carbon-](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/LDEO_Underway_Database/)
 214 acidification-data-system/oceans/LDEO_Underway_Database/.

- 215 **3) GLODAPv2:** The Global Ocean Data Analysis Project Version 2 (GLODAPv2) aggregates
 216 biogeochemical data collected from discrete bottle samples, offering extensive global coverage from the
 217 surface to depths (Key et al., 2015; Olsen et al., 2016; Lauvset et al., 2024). While GLODAP is primarily a
 218 product for basin-scale repeat hydrography data, it also includes coastal datasets and observations from a
 219 few time-series. The GLODAPv2 data product provides rigorously quality-controlled measurements for 14
 220 essential oceanographic variables: temperature, salinity, dissolved oxygen, nitrate, silicate, phosphate,
 221 dissolved inorganic carbon (DIC), total alkalinity (TA), pH, chlorofluorocarbons (CFC-11, CFC-12, CFC-
 222 113), carbon tetrachloride (CCl₄), and sulfur hexafluoride (SF₆). These variables, excluding temperature,
 223 undergo both primary and secondary quality control procedures to detect outliers and adjust for significant
 224 measurement biases. GLODAPv2 was first published in 2016 and has been updated annually through a
 225 living data process in Earth System Science Data since 2019. For these updates, new data (including
 226 historical data not previously included in the data product) are quality controlled and adjusted to the 2016
 227 version (Olsen et al., 2019; Olsen et al., 2020; Lauvset et al., 2021; Lauvset et al., 2022; Lauvset et al.,
 228 2024). Since the global repeat hydrography programs operate with decadal repetitions, the aim is to
 229 produce a completely new version of GLODAP, where all cruise datasets will be reevaluated, every



decade. Release of the GLODAPv3 data product is planned for 2026. For more information on the secondary quality control process, refer to Tanhua et al. (2010) and Lauvset and Tanhua (2015). GLODAPv2 offers two kind of products: the collection of quality controlled data from discrete bottle samples taken at sampling location (Key et al., 2015; Olsen et al., 2016; Olsen et al., 2019; Olsen et al., 2020; Lauvset et al., 2021; Lauvset et al., 2022; Lauvset et al., 2024), and a gridded product, interpolated to a 1°x1° grid and the 33 standard depth levels of WOA (World Ocean Atlas) (Lauvset et al., 2016). All versions of the GLODAPv2 data product can be accessed at <https://glodap.info/>.

4) JOA Suite: The Java OceanAtlas (JOA) Suite offers both a user-friendly application and a library of ocean profile data curated by Jim Swift (Scripps Institution of Oceanography, La Jolla, California, USA). Similar to GLODAPv2, this data product serves as a comprehensive repository of quality-controlled discrete bottle based measurements (and limited CTD), spanning from the surface to the depths of the global ocean. Unlike GLODAPv2, no offset corrections were applied to the JOA data product. The JOA data product encompasses a range of oceanographic variables including temperature, salinity, dissolved oxygen, DIC, TA, silicate, phosphate, nitrate, nitrite, CFC-11, CFC-12, SF₆, and CTD parameters associated with the water sample data. To access the JOA application and data, visit: <https://joa.ucsd.edu/>. Currently, there is not a peer-reviewed paper or public-accessible report for this data product. Cite the data product itself as: “Swift, J. (2022), Java OceanAtlas Data, https://joa.ucsd.edu/Data_homepage”, or cite the entire JOA Suite as: “Swift, J. and Osborne, J. (2022), The Java OceanAtlas Suite, <https://joa.ucsd.edu>”.

5) WOD: In addition to the GLODAPv2 and JOA Suite, users can also access historical and recent original biogeochemical data collected from discrete bottle samples in a uniform format and units, along with their originator quality control (QC) flags, through the World Ocean Database (WOD) (Mishonov et al., 2024). Like the JOA Suite, these measured data remain unaltered. The WOD allows users to filter and subset data with specific variables, platforms, institutions, projects, regions, or time periods (Garcia et al., 2024). Users can visualize sampling locations on a “distribution plot” and access a cruise list for all selected data and variables. Users also have the option of exporting data in NetCDF or Comma-Separated Values (CSV) formats. Additionally, all data in the WOD are reproducible and traceable to their original sources archived at NOAA’s National Centers for Environmental Information (NCEI) for long-term preservation. The WOD is accessible at <https://www.ncei.noaa.gov/products/world-ocean-database>.

6) SNAPO-CO₂: Metzl et al. (2024) aggregated over 44,400 measurements of DIC and TA from a series of research cruises and ships of opportunity (SOOP) across various oceanic regions in 1993-2022, under the aegis of several French research programs, to create a product called “Service National d’Analyse des Paramètres Océaniques du CO₂ (SNAPO-CO₂)”. The majority of the samples were analyzed by the Service National d’Analyse des Paramètres Océaniques du CO₂ (SNAPO-CO₂) at the LOCEAN laboratory in Paris, France. Sampling was performed either from CTD-rosette casts (Niskin bottles) or collected from the ship’s flow-through system (intake at roughly 5m depth). DIC and TA determinations were conducted



- simultaneously through potentiometric titration in a closed-cell setup, calibrated with certified reference material to achieve an accuracy of $\pm 4 \mu\text{mol kg}^{-1}$ for both parameters, as per Edmond (1970). This methodology was also applied for real-time measurements during OISO cruises, with data from the South Indian Ocean for 1998–2018 included in this compilation. The data is split into two sets — one for the global ocean and coastal zones, and another for the Mediterranean Sea — both accessible in the same format: <https://doi.org/10.17882/95414>. Additionally, this data product is available at OCADS (DOI: “10.25921/ptyh-0y90”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0285681.html> (Metzl et al., 2023)
- 7) Arctic Ocean anthropogenic carbon estimates:** This dataset includes anthropogenic carbon estimates in the Arctic Ocean based on measurements of transient tracers, such as CFC-12 and SF₆ (Terhaar et al., 2020; Tanhua et al., 2009). Using the transient time distribution (TTD) method, anthropogenic carbon estimates were estimated at measurement locations across all basins of the Arctic Ocean between 1983 and 2005. In addition to these estimates, adjusted estimates of anthropogenic carbon at these locations are provided that account for differences in the saturation of transient tracers and anthropogenic carbon in Arctic Ocean surface waters that caused anthropogenic carbon estimates to be biased low (Terhaar et al., 2020). It is recommended to use the adjusted estimates. This dataset can be accessed at <https://doi.org/10.17882/103920> (Terhaar et al., 2024).
- 8) CODAP-NA:** Jiang et al. (2021) curated and synthesized two decades of discrete measurements of carbonate system variables, dissolved oxygen, and nutrient chemistry data from the North American continental shelves to generate the first version of Coastal Ocean Data Analysis Data Product in North America (CODAP-NA). The 2021 release encompasses 3,391 oceanographic profiles from 61 research cruises spanning the North American continental shelves from Alaska to Mexico in the west and from Canada to the Caribbean in the east. It includes 14 key variables, including temperature, salinity, DO, DIC, TA, pH, carbonate ion, *f*CO₂, silicate, phosphate, nitrate, all of which have undergone rigorous quality control. Note that certain datasets meeting the GLODAPv2 QC standards are also included in the GLODAPv2 since its 2022 release. CODAP-NA is available at OCADS (DOI: “10.25921/531n-c230”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0219960.html> (Jiang et al., 2020).
- 9) AZMP Carbon:** Gibb et al. (2023) compiled carbonate parameters data from the Canadian Atlantic Zone Monitoring Program (AZMP Carbon) since 2014. More than 100 seagoing missions are represented in this dataset. The sample strategy corresponds generally to full-depth water samples mostly collected along standardised hydrographic sections. The majority of these data were collected as part of the Atlantic Zone Monitoring Program (AZMP) of Fisheries and Oceans Canada (DFO). Implemented in 1998, the AZMP aims to characterize and understand the causes of oceanic variability at the seasonal, inter-annual and decadal scales in support of, among other things, fisheries management in the Atlantic Zone (including the Gulf of St. Lawrence, the Scotian shelf and the Newfoundland and Labrador shelf). Since 2014, a minimum



of two of the three following carbonate parameters, TA, DIC and pH, are also acquired by the program at standardised hydrographic stations across the zone (sampled up to three times a year). Each measurement is completed with corresponding temperature, salinity and, when available, nutrients and dissolved oxygen concentration data. This dataset also includes samples collected as part of ships of opportunity, fishing and other scientific trips. The entire dataset comprises 19,531 discrete samples [last updated 21 August 2024]. Among this number, 18,085 have at least two of the three carbonate system parameters (e.g., TA, DIC and pH), allowing the derivation of other parameters such as the saturation state relative to aragonite and calcite (Ω_{arg} and Ω_{cal}) and partial pressure of CO_2 ($p\text{CO}_2$, in μatm) using the CO2SYS program modified for Python (<https://github.com/mvdm7/PyCO2SYS/tree/v1.2.1>, last access: 8 January 2023; Humphreys et al., 2020). The full dataset of measured and derived parameters is available from the Federated Research Data Repository: <https://doi.org/10.20383/102.0673> (Cyr et al., 2022) and is updated annually.

10) MOCHA: Kennedy et al. (2023) curated a comprehensive coastal ocean data product called “Multistressor Observations of Coastal Hypoxia and Acidification (MOCHA)”, encompassing temperature, salinity, dissolved oxygen, ocean carbonate variables (TA, DIC, pH, $p\text{CO}_2$, $f\text{CO}_2$), nutrients, and chlorophyll measurements from the full water column along the U.S. west coast. The synthesis integrates observations from 71 different sources, including high-resolution autonomous sensors, synoptic oceanographic cruises, and shoreline samples. The MOCHA synthesis spans from the shoreline to well-beyond the continental shelf and incorporates observations from CODAP-NA, CalCOFI, and other large-scale oceanographic cruises to facilitate linking nearshore, high-resolution observations to broader oceanographic conditions. It boasts 15.9 million temperature readings, 5.0 million salinity measurements, 3.9 million dissolved oxygen records, and 2.3 million pH measurements, along with 8,368 dissolved inorganic carbon, 10,144 total alkalinity, and 505,000 $p\text{CO}_2/f\text{CO}_2$ measurements, with limited additional chlorophyll and nutrient observations. To reduce the computational load from high-resolution sensors, the synthesis is also available as a “daily aggregated” dataset, with all data sources averaged by day, location, and depth. All data in the MOCHA synthesis product has been quality controlled to a “plausible and reasonable” standard, but researchers requiring high-precision coastal data may need to apply additional QC tests. The data product is available at OCADS (DOI: “10.25921/2vve-fh39”): <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0277984.html>, while the methods and product are described in Kennedy et al. (2024).

11) ARIOS: The Acidification in the Rias and the Iberian Continental Shelf (ARIOS) project involved compiling and analyzing the historical record of carbon system measurements and associated parameters conducted by the Instituto de Investigaciones Mariñas (IIM-CSIC) in Vigo, Spain. This dataset comprises 3,343 oceanographic stations and 17,653 discrete samples, combining measurements of pH, alkalinity, and other physical (pressure, temperature and salinity) and biogeochemical parameters (dissolved oxygen, nitrate, phosphate, and silicate) off the northwestern Iberian Peninsula from June 1976 to September 2018



(Padin et al., 2020). The oceanography cruises funded by 24 projects were primarily carried out in the Ría de Vigo coastal inlet, but also in an area ranging from the Bay of Biscay to the Portuguese coast. Robust seasonal cycles and long-term trends were calculated along a longitudinal section, gathering data from the coastal and oceanic zone of the Iberian upwelling system. The pH in the surface waters of these separated regions, which were highly variable due to intense photosynthesis and the remineralization of organic matter, showed an interannual acidification ranging from 0.0012 to 0.0039 yr⁻¹ that increased towards the coastline and interior embayments. A synthesis paper is available at <https://doi.org/10.5194/essd-12-2647-2020> (Padin et al., 2020), and the data product is available at <https://doi.org/10.20350/digitalCSIC/12498> (Pérez et al., 2020).

12) Marine inorganic carbon chemistry observations in the northern Gulf of Alaska: Monacci et al. (2023) compiled a data product of discrete seawater samples collected each May and September over a 10-year period from 2008 to 2017 along the long-term hydrographic line in the Gulf of Alaska (GAK Line). Samples were collected from a sampling rosette on a profiling CTD. Data variables include profiled seawater temperature, salinity, and dissolved oxygen. Discrete sample variables include dissolved oxygen (i.e., Winkler titrations), macronutrients (nitrate, nitrite, phosphate, silicic acid), dissolved inorganic carbon (DIC), and total alkalinity (TA). The repeat hydrographic cruises were funded by the Alaska Ocean Observing System (AOOS), the Exxon Valdez Oil Spill Trustee Council (EVOS), Gulf Watch Alaska, and the North Pacific Research Board (NPRB) and were mostly conducted aboard the United States Fish and Wildlife Service (USFWS) R/V *Tiglañ*. All carbonate parameters were analysed at the Ocean Acidification Research Center (OARC) at the University of Alaska Fairbanks (UAF). This data product is available at OCADS (DOI: 10.25921/x9sg-9b08): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0277034.html> (Monacci et al., 2023), and the synthesis paper can be accessed at <https://doi.org/10.5194/essd-16-647-2024> (Monacci et al., 2024).

13) Coral Reef Carbonate Chemistry Off the Florida Keys: Palacio-Castro et al. (2023) compiled discrete seawater samples from 38 permanent stations located along 10 inshore-offshore transects at the Florida Coral Reef. These samples were collected as part of NOAA's National Coral Reef Monitoring Program (NCRMP) and the South Florida Ecosystem Restoration Research (SFER) cruises. Sampling efforts commenced in 2010, with every two months collections initiated in 2015, resulting in a total of 47 sampling cruises and 1,538 discrete seawater samples. For all samples, a minimum of two of the carbonate equilibrium parameters (TA, DIC) were measured, in addition to salinity and temperature. The aragonite saturation state (Ω_{Ar}), partial pressure of CO₂ ($p\text{CO}_2$) and pH were derived from the measured parameters using the R package seacarb (Gattuso et al., 2021a). The time series analysis provides insight into the dynamic carbonate conditions spanning the inshore to offshore gradients, encompassing four distinct regions of the Florida Coral Reef: Biscayne Bay, the Upper Keys, Middle Keys, and Lower Keys. The findings underscore significant variability in the seasonality and interannual trends of surface carbonate



chemistry across different regions and reef zones. Data is available at NCEI (DOI: “10.25921/vfz0-dg77”):
<https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.nodc:NCRMP-CO3-Atlantic>
(Manzello et al., 2018).

14) Salish cruise data package and multi-stressor data product: Alin et al. (2021) compiled data from 35 individual cruise data sets that sampled marine waters of the southern Salish Sea and northern Washington coast (USA) during 2008–2018. These data sets were collected in support of research and monitoring efforts of the University of Washington Puget Sound Regional Synthesis Model (2008–2013), Washington Ocean Acidification Center (WOAC, 2014–present), NOAA Ocean Acidification Observing Network, Northwest Association of Networked Ocean Observing Systems, and NOAA Pacific Marine Environmental Laboratory. Ongoing seasonal sampling occurred during April, July, and September for Puget Sound cruises has occurred under WOAC support since 2014 and most frequently during May and October for Sound-to-Sea cruises, which sample from Puget Sound through the Strait of Juan de Fuca to the northern Washington coast. The Salish cruise data package contains observations from a total of 715 oceanographic profiles, with > 7490 sensor measurements of temperature, salinity, and oxygen; ≥ 6070 measurements of discrete oxygen and nutrient (nitrate, phosphate, silicate, ammonium, nitrite) samples; and ≥ 4462 measurements of inorganic carbon variables (DIC and TA). Alin et al. (2023) published a follow-on data product based on the Salish cruise data package, which only included the 3971 samples with complete information for temperature, salinity, oxygen, nutrients, DIC, and TA. To facilitate applications of this data product to understanding multi-stressor ocean conditions in Pacific Northwest marine waters by various end users, Alin et al. (2023) also included the most commonly used calculated carbonate system parameters in this data product (pH_{total} , $f\text{CO}_2$, $p\text{CO}_2$, Ω_{Ar} , Ω_{Ca}). The data package is available at OCADS (DOI: “10.25921/zgk5-ep63”): https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/SalishCruise_DataPackage.html. The multi-stressor data product is available at: https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/SalishCruises_DataProduct.html. Two synthesis papers describing the data package and interpreting the data product can be found at: <https://essd.copernicus.org/articles/16/837/2024/> (Alin et al. 2024a) and <https://bg.copernicus.org/articles/21/1639/2024/> (Alin et al. 2024b).

Table 1. Ocean carbonate chemistry data products out of cruise data compilations (no gridding or gap filling).

No.	Name	Open or coastal ocean	Surface or water column	Discrete bottle or continuous	Highlights	Reference
1	SOCAT	Open + Coastal	Surface	Continuous	The largest collection of surface ocean carbon observations	Bakker et al. (2016)
2	LDEO Surface $p\text{CO}_2$ Database	Open + Coastal	Surface	Continuous	The LDEO database reports $p\text{CO}_2$ exclusively from equilibrator- CO_2 analyzer systems	Takahashi et al. 2017



3	GLODAPv2	Open ocean	Water column	Discrete bottle	Adjustments are applied by comparing data in the deep ocean (>2000 m) using a crossover and inversion method as described by Johnson et al (2001).	Lauvset et al. (2024)
4	JOA Suite	Open ocean	Water column	Discrete bottle	Similar to GLODAPv2, with no adjustments	Swift (2022)
5	WOD	Open ocean	Water column	Discrete bottle	Similar to GLODAPv2, with no adjustments	Mishonov et al. (2024)
6	SNAPCO-CO ₂	Open + Coastal	Water column	Discrete bottle and semi-continuous	A compilation of cruises from multiple French initiatives	Metzl et al. (2024)
7	Arctic Ocean anthropogenic carbon estimates	Open ocean	water column	Discrete bottle (normalized to year 2005)	Observation-based estimates of anthropogenic carbon in the Arctic Ocean	Tanhua et al. (2009); Terhaar et al. (2020)
8	CODAP-NA	Coastal ocean	Water column	Discrete bottle	Similar to GLODAPv2, but for the coastal ocean	Jiang et al. (2021)
9	AZMP Carbon	Continental Shelf and slope	Water column	Discrete bottle	A compilation of cruises from the Atlantic Zone Monitoring Program (AZMP) since 2014	Gibb et al. (2023)
10	MOCHA	Coastal ocean	Water column	Discrete bottle + Continuous	U.S. West Coast	Kennedy et al. (2024)
11	ARIOS	Coastal ocean	Water column	Discrete bottle	An OA Database for the Galician Upwelling Ecosystem off the NW Iberian Peninsula from 1976 to 2018	Padin et al., (2020)
12	Marine inorganic carbon chemistry observations in the northern Gulf of Alaska	Coastal ocean	Water column	Discrete bottle	A synthesis of twenty cruises from 2008 to 2017 on the Gulf of Alaska (GAK) Line	Monacci et al. (2023)
13	Coral Reef Carbonate Chemistry Off the Florida Keys	Coastal / regional	Water column	Discrete bottle	Temporal trends of DIC, TA, pCO ₂ , pH, Ω_{arag} , in different areas of the Florida Keys	Palacio-Castro et al. (2023)
14	Salish cruise data package and multi-stressor data product	Coastal / estuarine	Water column	Discrete bottle	A data compilation and multi-stressor (ocean acidification, hypoxia, temperature) data product based on cruises from 2002 to 2018 in the southern Salish Sea and Washington coast	Alin et al. (2021, 2023)

3.1.2 Time-series data products (no interpolation or extrapolation):

The time-series data products described in this section include observations collected at regular time intervals, over a sustained period, and at fixed locations. The data often represent changes in a particular oceanographic variable over time, such as temperature, salinity, TA and DIC. The list below includes both climate-quality time-series data products compiled at selected stations, and data products compiling time-series measurements at multiple locations. Additionally, some hydrographic sections are measured regularly enough to warrant being called a time



series, e.g., Line P (Freeland, 2007), sections in the northwest Pacific (Ishii et al., 2011a), the OVIDE lines (Mercier et al., 2024). Measurements from these sections are typically included in cruise data compilations (3.1.1) and are not listed separately here.

15) BATS: The Bermuda Atlantic Time-series Study (BATS) observations and data products extend to forty years of observations of DIC and TA and ocean acidification indicators, and constitute the longest continuous record of warming, salinification, ocean deoxygenation, and ocean acidification in the open ocean (Bates and Johnson, 2023). The sustained observations at the BATS site began in October 1988, approximately 80 km to the south-east of Bermuda (<https://bios.asu.edu/bats>). The program comprises monthly cruises with CTD, water-column biogeochemical sampling and rate measurements (e.g., primary and export production) plus additional cruises in the spring period and annual transects between the Gulf Stream and Puerto Rico. CO₂-carbonate chemistry sampling includes full-depth bottle DIC and alkalinity data (including additional surface measurements going back to 1983 collected at the Hydrostation S site). Hydrostation S is located ~25 km south-east of Bermuda (<https://bios.asu.edu/research/projects/hydrostation-s>) and began in 1954 with biweekly cruises each year. Underway *f*CO₂/*p*CO₂ data collected from the R/V *Atlantic Explorer* that supports the BATS and Hydrostation S sites constitutes part of the annual data submission to SOCAT. The BATS project page at the Biological and Chemical Oceanography Data Management Office (BCO-DMO) includes metadata and data streams (<https://demo.bco-dmo.org/project/2124>). Hydrostation S data and DOIs are also available at BCO-DMO (<https://www.bco-dmo.org/project/859583>).

16) HOT: The Hawaii Ocean Time-series (HOT) CO₂ measurement program documents 35+ years of inorganic carbon dynamics in the open waters of the central North Pacific. Since October 1988, full ocean depth profiles of DIC and TA have been analyzed, and direct measurements of pH have been made over most of this longest-running Pacific Ocean time-series study. The overall scientific mission of HOT has been to sustain observations of the variability of hydrological and ecological properties, elemental cycling, heat fluxes, and circulation of the North Pacific Subtropical Gyre (NPSG). The program is based on shipboard observations and experiments conducted on ~10 expeditions per annum to Station ALOHA (22.75°N, 158°W). HOT program background information and details of sampling strategy may be found in Karl and Lukas (1996) and Karl et al. (2001). Results from the HOT CO₂ measurement program may be found in Winn et al. (1994, 1998), Dore et al. (2003, 2009, 2014), and Knor et al. (2023). The HOT project page, metadata, data streams and data identifiers are listed at <https://www.bco-dmo.org/project/2101>. A surface ocean data product that includes CO₂SYN-calculated values of *p*CO₂, carbonate mineral saturation states and other derived quantities may be found at <https://hahana.soest.hawaii.edu/hot/hotco2/hotco2.html> and <https://doi.org/10.5281/zenodo.15060930>. A MAPCO₂ system on the Woods Hole Oceanographic Institution Hawaii Ocean Time-series Site mooring (WHOTS; <https://www.soest.hawaii.edu/whots/>) has provided a near-continuous record of surface *p*CO₂ since 2004, and is anchored by the longer high-accuracy



HOT ship-based program (see Sutton et al. 2019 and Knor et al. 2023). These synergistic measurements have contributed to global ocean carbon observation networks (e.g., the newly released SOCATv22), which have improved our ability to characterize natural and anthropogenic drivers of ocean carbon uptake and acidification.

17) ESTOC: The European Station for Time Series in the Ocean (ESTOC) began carbon dioxide monitoring in October 1995, providing a 30-year record on total alkalinity (TA), pH, and dissolved inorganic carbon (DIC). This dataset represents the longest continuous monthly record of warming, rising carbon dioxide levels, and acidification in the eastern North Atlantic (González-Dávila and Santana-Casiano, 2023). ESTOC is located 100 km north of the Canary Islands archipelago (<https://plocan.eu/en/installations/ocean-observatory>). The program includes a ship-based observation system, measuring physical, chemical, and biological parameters throughout the 3,670-meter water column. It also features a moored platform for surface meteorological and oceanic observations as well as subsurface measurements, maintained by the Canary Island Oceanic Platform (PLOCAN, <https://plocan.eu/en>) and the University of Las Palmas de Gran Canaria (<https://iocag.ulpgc.es/research/research-units/quima>). Carbon dioxide system measurements include full-depth bottle sampling for photometric pH, total alkalinity, and DIC, conducted monthly from 1995 to 2008, every two months until 2018, and semiannually in recent years due to limited ship time, timed to coincide with moored structure maintenance. ESTOC is also visited every two weeks by a volunteer observing ship, ES-SOOP-CanOA (https://meta.icos-cp.eu/resources/stations/OS_687B), part of the European Research Infrastructure ICOS (<https://www.icos-cp.eu/observations/ocean/stations>), which provides real-time surface data on carbon dioxide fluxes and ocean acidification. The program also includes the CO₂-ESTOC oceanographic buoy (<https://meta.icos-cp.eu/labeling/>). The full dataset with DOIs is accessible on Pangea (González-Dávila and Santana-Casiano, 2023).

18) Point B Time-series: The Point B time series documents the carbonate chemistry at a coastal site of the Bay of Villefranche (43.686200N 7.314800E) in Villefranche-sur-mer, France, northwestern Mediterranean Sea. Since January 2007, seawater is sampled weekly at 1 and 50 m, and analyzed for total dissolved inorganic carbon and total alkalinity (Kapsenberg et al., 2017). Salinity and temperature are extracted from CTD profiles. Parameters of the carbonate system such as pH (total hydrogen ion scale) are calculated using the R package seacarb. Data are available at Pangea: <https://doi.org/10.1594/PANGAEA.727120> (Gattuso et al., 2021b).

19) Ny-Ålesund Time-series: The Ny-Ålesund time series documents the carbonate chemistry at a coastal site of Kongsfjorden, Spitsbergen (78.930660N 11.920030E) during the period 2015–2021. It is the first high-frequency (1 hour), multi-year (6 years) dataset of salinity, temperature, dissolved inorganic carbon, total alkalinity, pCO₂, and pH in the High-Arctic Ocean (Gattuso et al., 2023a). Data are available at Pangea: <https://doi.org/10.1594/PANGAEA.957028> (Gattuso et al., 2023b).



- 473 **20) SPOTS:** The Synthesis Product for Ocean Time-Series (SPOTS) is a ship-based biogeochemical pilot,
 474 aiming at regularly providing high quality data from fixed time-series stations with consistent format and
 475 semantics (Lange et al., 2024a). The pilot includes data from 12 fixed ship-based time-series programs with
 476 a focus on the Global Ocean Observing System’s biogeochemical essential ocean variables. These stations
 477 represent unique marine environments across a variety of spatiotemporal resolutions and ranges, with data
 478 from 1983 to 2021. While implementing the FAIR Principles (Wilkinson et al., 2016) and open data, Lange
 479 et al. also improved the metadata of the time-series stations to interoperate with the IOC-UNESCO Ocean
 480 Data and Information System (ODIS). Additionally, an extensive quality assessment resulted in enhanced
 481 intra- and inter-station comparability. Altogether, SPOTS’ pilot increased the readiness of biogeochemical
 482 time-series (Lange et al., 2023) and facilitates a variety of applications that benefit from the collective
 483 value of biogeochemical time-series observations. Data are available at [https://www.bco-](https://www.bco-dmo.org/dataset/896862)
 484 [dmo.org/dataset/896862](https://www.bco-dmo.org/dataset/896862) with the DOI “10.26008/1912/bco-dmo.896862.2” (Lange et al., 2024b).
- 485 **21) Autonomous $p\text{CO}_2$ and pH time series from 40 surface buoys:** Sutton et al. (2019) established a living
 486 dataset comprising 40 individual autonomous moored surface ocean $p\text{CO}_2$ time series established between
 487 2004 and 2013, 17 of them also include autonomous pH measurements. These time series characterize a
 488 wide range of surface-ocean carbonate conditions, across a variety of environments, including 17 oceanic
 489 and 13 coastal locations, as well as 10 coral reefs. The dataset serves as a significant resource for assessing
 490 the natural variability of surface ocean carbonate chemistry across diverse regions. Data are available at
 491 OCADS (DOI “10.7289/v5db8043”):
 492 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0173932.html> (Sutton et al., 2018).
 493 Additionally, there is a dedicated webpage at OCADS for this project:
 494 <https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/Moorings/ndp097.html>.

495 **Table 2. Time-series based ocean carbonate chemistry data synthesis products.**

No.	Name	Open or coastal ocean	Surface or water column	Discrete bottle or autonomous	Highlights	Reference
15	BATS	Open	Top 4600 m	CTD profiles, discrete bottle data from hydrocast, rate measurements (primary, export and bacterial production), biomass and other biological measurements, underway measurements	One of the longest continuous record of warming, salinification, ocean deoxygenation, and ocean acidification in the open ocean	Bates and Johnson (2020, 2023)
16	HOT	Open	Top 4500 m	Discrete Bottle for shipboard measurements; sensor measurements on WHOTS mooring	35+ years of inorganic carbon dynamics in the open waters of the central North Pacific	Dore et al. (2009)



17	ESTOC	Open	Top 3570m	Discrete Bottle for shipboard measurements; sensor measurements on ESTOC mooring	The longest continuous monthly record of warming, rising carbon dioxide levels, and OA in the eastern North Atlantic	González-Davila and Santana-Casiano (2023)
18	Point B Time-series	Coastal	5 and 50 m	CTD profiles, discrete bottle data	Carbonate chemistry at a coastal site of the Bay of Villefranche, France	Kapsenberg et al. (2017)
19	Ny-Ålesund Time-series	Coastal	12 m	CTD, discrete bottle data, sensor measurements at the COSYNA/MOSES-AWIPEV underwater observatory	Carbonate chemistry at a coastal site of Kongsfjorden, Spitsbergen	Gattuso et al. (2023)
20	SPOTS	Open + Coastal	Water column	Discrete	The pilot includes biogeochemical data from 12 fixed ship-based time series programs	Lange et al. (2024a), Lange et al. (2024b)
21	Autonomous $p\text{CO}_2$ and pH time series from 40 surface buoys	Open + Coastal	Surface	Autonomous	Based on 40 moored surface $p\text{CO}_2$ time series, with 17 of them containing pH	Sutton et al. (2019)

3.1.3 Gridded and derived data products - Surface:

This section describes gridded data products that have been derived from observations through interpolation and gap-filling procedures, and depict the surface ocean. Note that this compilation focuses primarily on data products with global coverage, acknowledging that many regional gap-filled products became available in recent years and shall be include in future updates:

22) Updated Takahashi delta $f\text{CO}_2$ and flux climatology: Following on previous climatologies published by the late Taro Takahashi in 1997 and 2009, Fay et al. (2024) created a legacy climatology using his methodology and the updated SOCAT database of observations. This product provides 12 months of delta $f\text{CO}_2$ values and corresponding fluxes for a base year of 2010 at $4^\circ \times 5^\circ$ resolution subsequently subgridded to $1^\circ \times 1^\circ$ resolution and near-global coverage. This climatology represents the mean of ocean conditions over the last four decades and is distinctive relative to many other mechanistic machine learning approaches in that it interpolates in time and space using only the available $f\text{CO}_2$ data and a surface water advection scheme rather than using proxy variables for gap filling. It uses the median of observations to determine a reference year of 2010 and fluxes are provided using air-sea partial pressure differences and inputs from the SeaFlux product (Fay et al., 2021). The climatology product is available at OCADS (DOI: "10.25921/295g-sn13"): <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0282251.html>. The related manuscript is available at ESSD: <https://doi.org/10.5194/essd-16-2123-2024> (Fay et al., 2024).



- 513 **23) MPI-ULB-SOM-FFN:** Landschützer et al. (2020a) created a uniform $p\text{CO}_2$ climatology combining open
 514 and coastal oceans. It is a monthly climatological gridded global surface ocean $p\text{CO}_2$ data product without
 515 adjusting for a specific reference year. Developed on a higher-resolution $0.25^\circ \times 0.25^\circ$ global surface-ocean
 516 grid, this product is the result of combining two neural network-based $p\text{CO}_2$ products: the open ocean
 517 product described below (i.e., Landschützer et al. 2016) and the coastal product created by Laruelle et al
 518 (2017). Consequently, it represents coastal zones better. Data collected between 1998 and 2015 from the
 519 SOCAT database (Version 5) were used to create this data product. The merged climatology is available at
 520 OCADS (DOI: “10.25921/qb25-f418”):
 521 <https://www.ncei.noaa.gov/data/oceans/ncEI/ocads/metadata/0209633.html>. Additionally, there is a
 522 dedicated web page at OCADS for this project: [https://www.ncei.noaa.gov/access/ocean-carbon-](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/MPI-ULB-SOM_FFN_clim.html)
 523 acidification-data-system/oceans/MPI-ULB-SOM_FFN_clim.html.
- 524 **24) VLIZ-SOM-FFN:** Landschützer et al. (2016) employed the Self-Organizing Map-Feed-Forward Network
 525 (SOM-FFN) neural network method designed by Landschützer et al. (2013) to map the sea surface $p\text{CO}_2$
 526 from the Surface Ocean CO_2 Atlas (SOCAT; see above No. 1) (Bakker et al., 2014) to generate monthly
 527 $p\text{CO}_2$ fields on a $1^\circ \times 1^\circ$ global surface ocean grid, covering the period from 1982 to near present. It is
 528 based on the gridded $p\text{CO}_2$ measurements from SOCAT and is updated regularly. The creation of the $p\text{CO}_2$
 529 fields involve a two-step neural network approach, which has been extensively detailed and validated in
 530 previous works by Landschützer et al. (2013, 2014, 2016). In the initial step, the global ocean is clustered
 531 into biogeochemical provinces, and subsequently, the non-linear relationship between CO_2 driver variables
 532 and gridded data from SOCAT (Bakker et al. 2016) is reconstructed. Air-sea CO_2 fluxes are also computed
 533 based on the air-sea $p\text{CO}_2$ difference, utilizing a bulk gas transfer formulation as described by Landschützer
 534 et al. (2013, 2014, 2016). The product is available at OCADS (DOI: “10.7289/v5z899n6”):
 535 <https://www.ncei.noaa.gov/data/oceans/ncEI/ocads/metadata/0160558.html>. Additionally, there is a
 536 dedicated page at OCADS for this project: [https://www.ncei.noaa.gov/access/ocean-carbon-acidification-](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/SPCO2_1982_present_ETH_SOM_FFN.html)
 537 data-system/oceans/SPCO2_1982_present_ETH_SOM_FFN.html.
- 538 **25) JMA-MLR:** Iida et al. (2021) developed a monthly data product for inorganic carbon variables on a $1^\circ \times$
 539 1° global surface ocean grid for the period 1993–2018. Variables include dissolved inorganic carbon (DIC),
 540 total alkalinity (TA), CO_2 partial pressure ($p\text{CO}_2$), sea-air CO_2 flux, pH, and aragonite saturation state
 541 (Ω_{arag}). They leveraged data products such as SOCAT.v2019 (Bakker et al., 2016) and GLODAPv2.2019
 542 (Olsen et al., 2019), as well as satellite-based variables, including sea-surface dynamic height (SSDH),
 543 mixed layer depth (MLD), and chlorophyll-a. The product is updated annually using the latest SOCAT and
 544 GLODAPv2 data. The data product can be accessed at:
 545 https://www.data.jma.go.jp/kaiyou/english/co2_flux/co2_flux_data_en.html.
 546



26) OceanSODA-ETHZ:

(a) OceanSODA-ETHZv1 is a monthly gridded global surface ocean data product for multiple ocean carbon variables, including dissolved inorganic carbon (DIC), total alkalinity (TA), partial pressure of carbon dioxide ($p\text{CO}_2$), pH on total scale, and saturation states of aragonite and calcite (Ω_{arag} and Ω_{calc}) (Gregor and Gruber, 2020; Gregor and Gruber, 2021; Gregor and Gruber, 2023; Ma et al., 2023). This dataset is structured on a $1^\circ \times 1^\circ$ global surface ocean grid with monthly resolution from 1982–2022, facilitating research on OA over seasonal to decadal scales. The OceanSODA-ETHZ data product was created by extrapolating in time and space the surface ocean observations of $f\text{CO}_2$ from SOCATv2022 (Bakker et al., 2016) and TA from GLODAPv2.2022 using the newly developed Geospatial Random Cluster Ensemble Regression (GRaCER) method (Gregor, 2021). TA and $p\text{CO}_2$ were then used to calculate the remaining parameters of the marine carbonate system with the PyCO2SYS software (Humphreys et al., 2022). Phosphate and silicate from the World Ocean Atlas (2018) product were used (Boyer et al., 2018; Garcia et al., 2018a). The OceanSODA-ETHZ data product is available at OCADS (DOI: “10.25921/m5wx-ja34”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0220059.html>.

(b) OceanSODA-ETHZv2 is a surface $f\text{CO}_2$ product with a $0.25^\circ \times 0.25^\circ$ spatial resolution and an 8-day temporal resolution, providing estimates starting from 1982 (Gregor et al., 2024a; Gregor et al., 2024b). The high-resolution outputs are suitable for investigating the shorter- and finer-scale dynamics of surface $f\text{CO}_2$. Despite sharing a name with its predecessor, OceanSODA-ETHZv2 does not provide TA estimates and employs a different methodology, as described in the following steps: 1) The atmospheric trend of CO_2 is removed by subtracting marine boundary layer CO_2 concentrations from SOCAT $f\text{CO}_2$ producing a new target $\Delta^*\text{CO}_2$ to reduce the biases at the start and end of the time series. 2) An 8-day seasonal climatology of $\Delta^*\text{CO}_2$ is estimated using Gradient Boosted Decision Trees (GBDT), which is later used as a predictor. 3) The non-seasonal thermal component is removed from $\Delta^*\text{CO}_2$, resulting in a new target, $\Delta^*\text{CO}_2^{\text{nonT}}$. 4) The new target is estimated using a feed-forward neural network, with the GBDT as one of the forcing variables. 5) Steps 4 through to 1 are inverted to arrive at $f\text{CO}_2$. 6) Air-sea CO_2 fluxes are computed using ERA5 winds. Data are available at <https://doi.org/10.5281/zenodo.11206365> and are updated annually.

27) LDEO-HPD $f\text{CO}_2$ product: The LDEO Hybrid Physics Data (LDEO-HPD) estimates the temporal evolution of surface ocean $f\text{CO}_2$ and air-sea CO_2 exchange, utilizing the strengths of observations and global ocean biogeochemical models (GOBMs) (Gloege et al., 2022). GOBMs are internally consistent, mechanistic representations of the ocean circulation and carbon cycle, and have long been the standard for making spatiotemporally resolved estimates of air-sea CO_2 fluxes. However, there is often a bias between the modelled $f\text{CO}_2$ and available surface ocean measurements (Fay and McKinley 2021). The LDEO-HPD approach trains an eXtreme Gradient Boosting (XGB) algorithm to learn a non-linear relationship between model-data $f\text{CO}_2$ mismatch and observed predictor variables: SST, SSS, chlorophyll concentration, mixed layer depth). The GOBM $f\text{CO}_2$ is then corrected with the predicted model-data misfit to estimate real-world $f\text{CO}_2$ for the observation period (Gloege et al. 2022). The results in reconstructed monthly surface ocean



583 $f\text{CO}_2$ and air-sea CO_2 fluxes on a $1^\circ \times 1^\circ$ grid in the open ocean beginning in 1982. Additional information
 584 can be found at oceanarbon.ldeo.columbia.edu. The data product is available at:
 585 <https://zenodo.org/records/4760205>.

586 **28) LDEO HPD product with extended temporal coverage:** Building on the work of Gloege et al. (2022),
 587 the LDEO-HPD product as mentioned above (nr. 25) can be extended back in time to predict $f\text{CO}_2$ for all
 588 available model years. Bennington et al (2022a) find that the largest component of the GOBM corrections
 589 is climatological. The smaller corrections at other timescales suggest either that these are well captured by
 590 the GOBMs or the data are insufficient. The dominance of climatological corrections supports the
 591 extension of the LDEO-HPD $f\text{CO}_2$ product backwards in time. A climatology of model-observation misfits
 592 for the best-observed period (2000–present) is applied to the GOBMs for 1959–1981, while an inter-
 593 annually varying correction is used for 1982 onward. (Bennington et al 2022a). This results in
 594 reconstructed monthly surface ocean $f\text{CO}_2$ and air–sea CO_2 fluxes on a $1^\circ \times 1^\circ$ grid covering the open
 595 ocean, beginning in 1959. Since 2022, the LDEO-HPD Back in Time product has been included in the
 596 annual release of the Global Carbon Budget. Additional information can be found at
 597 oceanarbon.ldeo.columbia.edu. The data product can be accessed via Zenodo at
 598 <https://zenodo.org/records/13891722>.

599 **29) LDEO $f\text{CO}_2$ - Residual Method:** A frequently used approach for estimating full-coverage $f\text{CO}_2$ is to train
 600 a machine learning algorithm on sparse in situ $f\text{CO}_2$ data and associated physical and biogeochemical
 601 observations. While these associated variables have well-known relationships to $f\text{CO}_2$, it is often unclear
 602 how they mechanistically drive $f\text{CO}_2$ around the world. The LDEO $f\text{CO}_2$ -Residual method takes the basic
 603 approach and enhances connections between physical understanding and reconstructed $f\text{CO}_2$. The novel
 604 approach used here includes applying pre-processing to the $f\text{CO}_2$ data to remove the direct effect of
 605 temperature – a relationship well-documented in literature and lab experiments (Takahashi et al. 2002).
 606 This enhances the biogeochemical/physical component of $f\text{CO}_2$ in the target variable (now $f\text{CO}_2$ -Residual)
 607 and reduces the complexity that the machine learning must disentangle. The resulting algorithm has
 608 physically understandable connections between input data and the output biogeochemical/physical
 609 component of $f\text{CO}_2$ (Bennington et al. 2022b). This results in reconstructed monthly surface ocean $f\text{CO}_2$
 610 and air–sea CO_2 fluxes on a $1^\circ \times 1^\circ$ grid covering the open ocean, beginning in 1982 and extended to the
 611 most recent year of available data. Additional information can be found at oceanarbon.ldeo.columbia.edu.
 612 The data product can be accessed via Zenodo at <https://zenodo.org/records/13941548>.

613 **30) CMEMS-LSCE surface ocean carbon data products:**

614 (a) **CMEMS-LSCEv1:** Monthly surface ocean $p\text{CO}_2$ and air–sea CO_2 fluxes on a $1^\circ \times 1^\circ$ grid in both the
 615 open ocean and coastal seas from 1985–2019 were reconstructed by Chau et al., (2022). CMEMS-LSCE is
 616 short for Copernicus Marine Environment Monitoring Service - Laboratoire des Sciences du Climat et de
 617 l'Environnement. This product is generated from an ensemble-based reconstruction of $p\text{CO}_2$ maps trained



with gridded data from SOCATv2020 (Bakker et al., 2016). Sea-surface $p\text{CO}_2$ values (converted from the original $f\text{CO}_2$ values in SOCATv2020) were regressed against a set of predictors with non-linear functions, i.e., feed-forward neural network (FFNN) models. The predictors include: sea-surface height (SSH), SST, SSS, MLD, chlorophyll a (Chl-a), atmospheric CO_2 mole fraction ($x\text{CO}_2$), and geographical coordinates (longitudes and latitudes). This data product is accessible at:
<https://data.ipsl.fr/catalog/srv/eng/catalog.search#/metadata/a2f0891b-763a-49e9-af1b-78ed78b16982>.

(b) **CMEMS-LSCEv2**: CMEMS-LSCEv2 corresponds to the latest version of the CMEMS-LSCE FFNN. It uses the same ensemble-based reconstruction method for $p\text{CO}_2$ maps as CMEMS-LSCEv1. Improvements include downscaling the spatial resolution to $0.25^\circ \times 0.25^\circ$ and reproducing additional surface ocean carbonate system variables on a global grid from 1985 onwards (Chau et al., 2024a). The additional surface ocean carbonate system variables are: $p\text{CO}_2$, DIC, TA, pH, and saturation states with respect to aragonite (Ω_{ar}) and calcite (Ω_{ca}). Surface ocean $p\text{CO}_2$ is reconstructed based on an ensemble of neural network models mapping gridded observation-based data provided by SOCATv2022 (Bakker et al., 2016). Surface ocean TA is estimated with a multiple linear regression approach (Carter et al., 2016, 2017). The remaining carbonate variables are calculated from $p\text{CO}_2$ and TA using a MATLAB version of CO2SYS (Lewis and Wallace, 1998; Van Heuven et al., 2011). The CMEMS-LSCE product is updated yearly for surface ocean $p\text{CO}_2$, air-sea fluxes, and the carbonate system variables. Updates are phased with release of the SOCAT database. For surface ocean $p\text{CO}_2$ and air-sea fluxes the temporal coverage is extended to the present date with a latency of 1 month (Chau et al., 2024b). Both the multi-year reconstruction and the near-real time prediction can be accessed through the CMEMS portal:
<https://doi.org/10.48670/moi-00047>.

31) CarboScope (Jena-MLS): The Jena Mixed-Layer Scheme (within the CarboScope family of data-based estimates of carbon-cycle variability) is based on observed sea surface $p\text{CO}_2$ from the Surface Ocean CO_2 Atlas (SOCAT; see above No. 1) (Bakker et al., 2014). It provides daily global fields of $p\text{CO}_2$ and sea-air CO_2 fluxes from 1957 to the year before present, on a resolution of 2.5×2 degrees. In the original method (Rödenbeck et al, 2013), a diagnostic model of the carbon balance in the ocean mixed layer is being fitted to the $p\text{CO}_2$ data, by adjusting the ocean-interior sources and sinks of carbon of the mixed layer. The multi-decadal trend is derived from the data-based Ocean Circulation Inverse Model (OCIM) estimate provided by DeVries et al. (2022). Since a later extension described in Rödenbeck et al. (2022), the variability in the ocean-interior sources and sinks is first regressed against variability in SST and wind speed. The regression step is followed by a correction step with explicit temporal variability, to also represent data variability not yet represented by the predictors of the regression. The CarboScope product is updated yearly. The results from current and previous releases can be downloaded from <https://www.bgc-jena.mpg.de/CarboScope/>.



- 651 **32) UOEx-Watson:** This product is an estimate of the atmosphere-ocean flux of CO₂ that takes into account
 652 near-surface temperature deviations (Watson et al., 2020). Most estimates use data on surface ocean *p*CO₂
 653 without considering corrections due to temperature gradients within the uppermost few millimeters of the
 654 sea surface (“Skin temperature effects”) or small effects due to changes in temperature that occur during
 655 sampling and measurement, especially when the measurement is from a commercial vessel rather than a
 656 research ship. This product takes these effects into account by recalculating *p*CO₂ from the SOCAT data
 657 base (v2019) using co-located satellite observations of skin temperature. The result is a substantial increase
 658 in the calculated net global uptake of CO₂. In other respects, the methodology for this data product follows
 659 the two-step neural network approach described by Landschützer et al. (2013, 2014). The gridded data set
 660 of sea surface *f*CO₂ adjusted to satellite-derived subskin surface temperature, is available at
 661 <https://doi.org/10.1594/PANGAEA.905316>. Ocean-atmosphere fluxes interpolated to monthly and 1° × 1°
 662 degree spatial resolution is available at OCADS (DOI: “10.25921/2dp5-xm29”):
 663 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0301544.html>.
- 664 **33) NIES-ML3:** The National Institute for Environmental Studies (NIES-ML3) product includes monthly
 665 global surface ocean *f*CO₂ in 1982-2023 on a 1° × 1° grid. Using a leave-one-year-out (LOYO) validation
 666 method and three machine learning models, Zeng et al. (2022) found that the time variant trends of ocean
 667 CO₂ could be estimated approximately by a harmonic function fitting of the annual atmospheric CO₂. They
 668 removed the estimated trends from the ocean CO₂ and applied the LOYO to the trend-removed data to
 669 obtain the trend that could not be approximated by the fitting for trend correction. The trend-removed data
 670 by the corrected trends were used to train the models. The gap-filled CO₂ maps were constructed by adding
 671 the trends to model predictions. The product is available at NIES: <https://doi.org/10.17595/20220311.001>
 672 (Zeng 2022).
- 673 **34) CSIR-ML6:** Provides monthly 1° × 1° estimates of surface *p*CO₂ (Gregor et al., 2019a). The approach uses
 674 the conceptual two-step approach of clustering and performing regressions for each cluster as Landschützer
 675 et al. (2016). CSIR-ML6 investigates the efficacy of various machine learning (ML) methods in estimating
 676 surface *p*CO₂, namely, feed-forward neural networks (FFNN), extremely randomized trees (ERT), gradient
 677 boosting machines (GBM), and support vector regression (SVR). It is found that the ensemble of all but the
 678 ERT method resulted in the best estimate, highlighting the fact that various ML methods do not produce the
 679 same outcome, particularly when data is sparse. Further, the variance between ensemble members can
 680 inform us about regions where uncertainty may be large due to methodological differences. Despite this, all
 681 methods achieve roughly the same uncertainty – a barrier, or wall beyond which the community has yet to
 682 overcome. The data are available at OCADS (DOI: “10.25921/z682-mn47”):
 683 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0206205.html>. The product is one of the six
 684 ensemble members of the SeaFlux dataset.



- 685 **35) AOML-ET:** Wanninkhof et al. (2024, 2025) developed a monthly global ocean data product of seawater
 686 $p\text{CO}_2$ and sea-air CO_2 fluxes, referred to as AOML-ET, using an extremely randomized trees (ET) machine
 687 learning technique. These maps are created on $1^\circ \times 1^\circ$ spatial grids, providing global surface ocean
 688 coverages from 1998 to 2023. AOML-ET incorporates several predictor variables, including time, location,
 689 SST, SSS, MLD, and chlorophyll-a. The model was trained using the v2020 and v2023 releases of the
 690 SOCAT data product (No. 1). Sea-air CO_2 fluxes were calculated using the air-sea CO_2 **partial**
 691 **pressure difference** ($\Delta p\text{CO}_2$) and a bulk gas transfer formulation incorporating windspeed. The
 692 dataset contains monthly $1^\circ \times 1^\circ$ NetCDF files of the AOML-ET outputs, along with the predictor
 693 variables. The data are available at OCADS (DOI: “10.25921/0s8y-q287”):
 694 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0298989.html>.
- 695 **36) ULB-SOM-FFN-coastalv2.1:** Roobaert et al. (2024) present high-resolution ($0.25^\circ \times 0.25^\circ$ grid) monthly
 696 maps showing the distribution of sea surface $p\text{CO}_2$ across the global coastal oceans, spanning from 1982 to
 697 2020. This product (ULB-SOM-FFN-coastalv2.1) builds upon the work by Laruelle et al. (2017),
 698 incorporating a two-step methodology that utilizes Self Organizing Maps (SOM) and Feed Forward
 699 Networks (FFN). This updated product now captures temporal variability, enabling the assessment of
 700 interannual variability and long-term trends in coastal air-sea CO_2 exchange, unlike the product by Laruelle
 701 et al. (2017), which only offers a climatology for a short period (1998-2015). The enhancements include
 702 additional environmental predictors and an expanded dataset for training and validation, featuring
 703 approximately 18 million direct coastal observations from the Surface Ocean CO_2 Atlas (SOCAT)
 704 database, specifically the SOCATv2022 release (Bakker et al., 2016). The product is available at OCADS
 705 (DOI: “10.25921/4sde-p068”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0279118.html>
 706 (Roobaert et al., 2023).
- 707 **37) RFR-LME:** Sharp et al. (2024b) developed a data product delineating the temporal trends of OA indicators
 708 mapped on a $0.25^\circ \times 0.25^\circ$ spatial grid, across eleven U.S. Large Marine Ecosystems (LMEs), with
 709 monthly coverage from 1998–2023. These indicators, which include the $p\text{CO}_2$, pH, Ω_{arag} , DIC, TA, Revelle
 710 Factors, among others, were derived from SOCATv2023, along with other oceanographic properties, e.g.,
 711 SST, SSS, SSH, and MLD. The methodology combined Gaussian Mixture Models to categorize the data
 712 into environmentally similar subregions, Random Forest Regressions for the spatial and temporal
 713 extrapolation of observational $f\text{CO}_2$ data, and regressions to estimate TA (Carter et al. 2021) to provide a
 714 second carbonate system constraint. The resulting maps are available at OCADS (DOI: “10.25921/h8vw-
 715 e872”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0287551.html> (Sharp et al., 2024a),
 716 while an online portal at <https://ecowatch.noaa.gov/thematic/ocean-acidification> presents regionally
 717 averaged time series for three key indicators: the partial pressure of carbon dioxide, calcium carbonate
 718 saturation state with respect to aragonite, and seawater pH.



- 719 **38) ReCAD-NAACOM- $p\text{CO}_2$:** Wu et al. (2025) developed a reconstructed $p\text{CO}_2$ product for the North
 720 American Atlantic Coastal Ocean Margins (NAACOM), spanning from the Gulf of Mexico/Gulf of
 721 America to the Grand Banks, called the Reconstructed Coastal Acidification Database- $p\text{CO}_2$ (ReCAD-
 722 NAACOM- $p\text{CO}_2$). This product employed a two-step approach combining random forest regression and
 723 linear regression to generate monthly $p\text{CO}_2$ data at 0.25° spatial resolution from 1993–2021. The model
 724 was trained using SOCAT v2023 observations as ground-truth values, incorporating various satellite-
 725 derived and reanalysis environmental variables known to influence sea surface $p\text{CO}_2$. The ReCAD-
 726 NAACOM- $p\text{CO}_2$ product demonstrates robust performance, with a coefficient of determination of 0.83, a
 727 root-mean-square error of $18.64 \mu\text{atm}$, and an accumulative uncertainty of $23.83 \mu\text{atm}$ when compared
 728 against all SOCAT observation samples. This high accuracy was maintained across model training and
 729 validation phases. The model exhibits robust performance when validated against independent test data
 730 from a period excluded from both training and validation, underscoring the product's capability to reliably
 731 reconstruct $p\text{CO}_2$ in periods lacking direct observational data within the NAACOM. Significantly, the
 732 product enables detailed investigation of regional spatial differences, seasonal cycles, and decadal changes
 733 in $p\text{CO}_2$ across the NAACOM. The ReCAD-NAACOM- $p\text{CO}_2$ dataset is publicly accessible
 734 (<https://doi.org/10.5281/zenodo.11500974>) and will be updated regularly.
- 735 **39) Gridded surface OA indicators, and air–sea CO_2 fluxes in the northern Caribbean Sea:** This dataset
 736 contains a high-quality dataset of derived products from over a million observations of surface water partial
 737 pressure/fugacity of carbon dioxide ($p\text{CO}_{2\text{w}}/f\text{CO}_{2\text{w}}$), for the Caribbean Sea, Gulf of Mexico/Gulf of
 738 America and North-West Atlantic Ocean covering the timespan from 2002-01-01 to 2019-12-30. The
 739 derived quantities include total alkalinity (TA), acidity (pH), aragonite saturation state (Ω_{Ar}) and air-sea
 740 CO_2 flux (Wanninkhof et al., 2020). This data product is available at OCADS (DOI: “10.25921/2swk-
 741 9w56”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0207749.html> (Wanninkhof et al.,
 742 2019)
- 743 **40) Ocean acidification data in the Gulf of Mexico/Gulf of America and wider Caribbean from 2014 to**
 744 **2020:** The Acidification, Climate, and Coral Reef Ecosystems Team (ACCRETE) Lab within AOML’s
 745 Ocean Chemistry and Ecosystems Division (OCED) developed a data product for tracking ocean
 746 acidification in the Caribbean and Gulf of Mexico/Gulf of America (van Hooideonk, 2022). Utilizing
 747 satellite imagery and a data-assimilative hybrid model, the tool maps key indicators of the water’s carbonate
 748 system, including $p\text{CO}_2$, TA, pH, Ω_{arag} , and Ω_{calc} . This innovation builds upon an update to the
 749 experimental Ocean Acidification Product Suite (OAPS) developed by NOAA’s Coral Reef Watch. The
 750 data product is available at OCADS (DOI: “10.25921/tt1c-dx53”):
 751 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0245950.html>. (van Hooideonk, 2022)



752 **41) Regional $p\text{CO}_2$ climatology of the Baltic Sea:** Bittig et al. (2024) used biogeochemical model output to
753 inform the mapping of sea surface $p\text{CO}_2$ observations in the Baltic Sea and to build a mean monthly
754 climatology for the period 2003 to 2021 with 3 nautical mile resolution. In a first step, spatial patterns of
755 variability were extracted from 20 years of model surface $p\text{CO}_2$ data by an EOF analysis. These spatial
756 patterns were then used to map surface $p\text{CO}_2$ observations from the Surface Ocean CO_2 Atlas (SOCAT; see
757 above No. 1) (Bakker et al., 2014) onto the Baltic Sea domain. By using an ensemble approach with
758 varying number of EOF patterns, the spatial scales of the mapping were locally adjusted based on the
759 observation's data density. Mapped monthly fields of $p\text{CO}_2$ from 2003-2021 were combined for the product
760 into a mean monthly climatology and a spatially-resolved linear trend. The climatology product is available
761 at PANGAEA (DOI: "10.1594/PANGAEA.961119"): <https://doi.org/10.1594/PANGAEA.961119> (Bittig
762 et al., 2023).

763 **Table 3. Gridded and derived ocean carbonate chemistry data synthesis products in the surface ocean.**

No.	Name	Open or coastal ocean	Spatial resolution	Temporal resolution	Methodology	Highlights	Reference
22	Updated Takahashi delta $f\text{CO}_2$ and flux climatology	Open ocean	$1^\circ \times 1^\circ$	12-month climatology for a base year of 2010	Advection Scheme	Does not use proxy variables for extrapolation. Only produced as monthly climatology.	Fay et al. (2024)
23	MPI-ULB-SOM-FFN	Open + Coastal	$0.25^\circ \times 0.25^\circ$	Monthly climatology (January through December) without reference year	2-step machine learning: Merged product of the SOM-FFN approach applied to the open ocean (Landschützer et al 2016) and the coastal ocean (Laruelle et al 2017)	Monthly gridded $p\text{CO}_2$ without adjusting for a specific reference year, high-resolution coastal ocean coverage	Landschützer et al. (2020a)
24	VLIZ SOM-FFN	Open ocean	$1^\circ \times 1^\circ$	Monthly from January 1982 onwards	2-step machine learning: Self organizing map clustering followed by a feed forward network (SOM-FFN)	Monthly gridded $p\text{CO}_2$ from 1982 through near present	Landschützer et al. (2016)
25	JMA-MLR	Open ocean	$1^\circ \times 1^\circ$	Monthly from January 1990 onwards	Multiple linear regressions	Temporal trends of DIC, TA, $p\text{CO}_2$, air-sea CO_2 flux, pH, and Ω_{arag}	Iida et al. (2021)
26(a)	OceanSODA-ETHZv1	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1982 to 2023	Ensemble of 2-step members: K-means clustering with gradient boosting and SVR regression	Temporal trends of DIC, TA, $p\text{CO}_2$, pH, Ω_{arag} , and Ω_{calc}	Gregor and Gruber (2021)



26(b)	OceanSODA-ETHZv2	Open + Coastal ocean	$0.25^\circ \times 0.25^\circ$	8-day from 1982 to 2022	FFNN	Highlighting fine-scale and short-term variability of the ocean carbon sink	Gregor, Shutler, and Gruber (2024)
27	LDEO-HPD $f\text{CO}_2$ product	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1982	XGBoost algorithm	Temporal evolution of surface ocean $f\text{CO}_2$ and air-sea CO_2 exchange	Gloege et al. (2022)
28	LDEO HPD with extended temporal coverage	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1959	XGBoost algorithm	uses model-data misfit climatology to extend estimate back in time to 1959	Bennington et al (2022a)
29	LDEO $f\text{CO}_2$ - Residual Method	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1982	XGBoost algorithm	Removes the temperature component before ML	Bennington et al (2022b)
30(a)	CMEMS-LSCEv1	Open + Coastal	$1^\circ \times 1^\circ$	Monthly from 1985 to 2019	FFNN	seamless reconstruction from coastal to open ocean	Chau et al. (2022)
30(b)	CMEMS-LSCEv2	Open + Coastal	$0.25^\circ \times 0.25^\circ$	Monthly from 1985 to 2021	FFNN	yearly extension of time series & monthly reconstruction at low latency	Chau et al. (2024)a,b
31	CarboScope (Jena-MLS)	Open ocean	$2.5^\circ \times 2^\circ$	Daily from 1957	Multi-linear regression against long-term predictors, plus auto-regressive correction	Variability of pCO_2 and sea-air CO_2 fluxes since 1957, sensitivities to SST and wind speed variations	Rödenbeck et al. (2022)
32	UOEx-Watson	Open ocean	$1^\circ \times 1^\circ$	Monthly from January 1992	Two-step neural network approach described by Landschützer et al.	air-sea fluxes of CO_2 with adjusted skin temperature effect	Watson et al., 2020
33	NIES-ML3	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1982 to 2023	FNN, GBM, RF	The prediction of a ML method was obtained from ten trainings with different seeds. The mean of the three methods was taken as the final prediction.	Zeng (2022), Zeng et al. (2022)
34	CSIR-ML6	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1982 to 2016	Ensemble: FFNN, SVR, ERT, Gradient Boosted Trees	Various ML methods produce different results when data is sparse, but all still achieving roughly the same uncertainty.	Gregor et al. (2019a)
35	AOML-ET	Open ocean	$1^\circ \times 1^\circ$	Monthly from 1998 to 2023	Extremely randomized trees (ET) machine learning technique	Monthly global sea-air CO_2 flux maps in modern era	Wanninkhof et al. (2015)



36	ULB-SOM-FFN-coastalv2.1	Coastal ocean	$0.25^\circ \times 0.25^\circ$	Monthly from 1982 to 2020	2-step machine learning: Self organizing map clustering followed by a feed forward network (SOM-FFN)	Global temporal trends of coastal $p\text{CO}_2$ and air-sea CO_2 fluxes based on SOCATv2022 with data collected from 1982–2020	Roobaert et al. (2024)
37	RFR-LME	Coastal	$0.25^\circ \times 0.25^\circ$	Monthly from 1998 to 2023	Gaussian mixture models and random forest regressions	Temporal trends of OA indicators and estimated uncertainties across 11 U.S. Large Marine Ecosystems (LMEs), with monthly coverage from 1998–2023.	Sharp et al. (2024b)
38	ReCAD-NAACOM- $p\text{CO}_2$	Coastal	$0.25^\circ \times 0.25^\circ$	Monthly from 1993 to 2021	2-step machine learning: random forest regression followed by linear regression	Sea surface $p\text{CO}_2$ in the North American Atlantic Coastal Ocean Margins (NAACOM)	Wu et al. (2025)
39	Gridded surface OA indicators, and air-sea CO_2 fluxes in the northern Caribbean Sea	Coastal	$1^\circ \times 1^\circ$	Monthly from 2002 to 2019	Gridding of the observations of $f\text{CO}_2$, SST and SSS was performed by binning and averaging the data in ($1^\circ \times 1^\circ$ by month) cells	A 17-year record of $f\text{CO}_2$, TA, pH, Ω_{Ar} , and air-sea CO_2 flux in the Caribbean Sea	Wanninkhof et al. (2020)
40	OA data in the Gulf of Mexico/Gulf of America and wider Caribbean from 2014 to 2020	Regional	$0.088^\circ \times 0.88^\circ$	Monthly from 2014 to 2020	Utilizing satellite imagery and a data-assimilative hybrid model, the tool maps key indicators of the water's carbonate system, including $p\text{CO}_2$, TA, pH, Ω_{Ar} , and Ω_{calc} .	A new tool to monitor ocean acidification over the wider Caribbean and Gulf of Mexico/Gulf of America.	van Hooijdonk (2022)
41	Regional $p\text{CO}_2$ climatology of the Baltic Sea	Regional	$0.1^\circ \times 0.05^\circ$	12-month climatology for a base year of 2013; linear trend 2003-2021	Extrapolation using model-based patterns of variability	Does not use proxy variables for extrapolation. Spatial scales adjust locally to data density	Bittig et al. (2024)

3.1.4 Gridded and derived data products – Interior ocean:

Although cruise data compilations are valuable for making data available in a uniform format, they often are constrained by their sampling strategies and can have significant gaps in space and time. Gridded and derived data products address this limitation by making some variables available at all grid points on a standardized spatial grid and at standardized depth levels through processes such as interpolation and gap filling.



- 769 **42) Global interior ocean mapped climatology from GLODAPv2 (referenced to 2002):** Lauvset et al.
 770 (2016) generated a comprehensive set of global interior ocean climatologies, mapping key biogeochemical
 771 variables on a $1^\circ \times 1^\circ$ grid for 33 depth levels from surface to 5500 m. These climatologies cover
 772 temperature, salinity, oxygen, nitrate, phosphate, silicate, DIC, TA, pH, and the saturation states of
 773 aragonite and calcite (Ω_{arag} and Ω_{calc}). This data product was created based on the quality-controlled and
 774 internally consistent GLODAPv2.2016 (Olsen et al., 2016) using the Data-Interpolating Variational
 775 Analysis (DIVA) method (Barth et al., 2014). The conceivably confounding temporal trends in DIC, pH,
 776 Ω_{arag} and Ω_{calc} due to anthropogenic influence were removed prior to mapping by normalizing their values
 777 to a reference year of 2002 using first-order calculations of anthropogenic carbon accumulation rates. For
 778 all variables, all data from the full 1972–2013 period were used, including data that did not receive full
 779 secondary quality control. This data product is not updated each year along with the main GLODAPv2 data
 780 product. The mapped data product is available at OCADS (DOI: “10.3334/cdiac/otg.ndp093_glodapv2”):
 781 <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0286118.html> (Lauvset et al., 2023b). It can
 782 also be accessed from the GLODAP website: <https://glodap.info/>.
- 783 **43) Global aragonite saturation state climatology (referenced to 2000):** Jiang et al. (2015a) developed a
 784 climatology of the derived variable, aragonite saturation state (Ω_{arag}) across the global interior ocean, on a
 785 $1^\circ \times 1^\circ$ grid at 9 standardized depth levels from the surface down to 4000m. This was accomplished by
 786 integrating data from the first version of GLODAP (Key et al., 2004), CARINA (Key et al., 2010), and
 787 PACIFICA (Suzuki et al., 2013), along with additional recent cruise datasets up to 2012. Calculations of
 788 Ω_{arag} utilized a MATLAB version of the CO2SYS program (Orr et al., 2015), with the dissociation
 789 constants for carbonic acid of Lueker et al. [2000], potassium bisulfate (KHSO_4^-) of Dickson [1990],
 790 hydrofluoric acid (HF) of Perez and Fraga (1987), and the total borate concentration equations of Lee et al.
 791 [2010]. Temporal adjustments were made to a reference year of 2000, accounting for an annual increase of
 792 $f\text{CO}_2$ of $1.6 \mu\text{atm}$ in the surface mixed layer (SML), with a rate that decreases linearly to zero $\mu\text{atm yr}^{-1}$
 793 from the bottom of the SML to a depth of 1000 m (Sabine et al., 2008). The data product is available at
 794 OCADS (DOI: “10.1002/2015GB005198”):
 795 <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0139360.html> (Jiang et al., 2015b).
- 796 **44) MOBO-DIC (Version 2020):** Keppler et al. (2020) produced a global interior ocean DIC monthly
 797 climatology (average climatological values for January through December) on a $1^\circ \times 1^\circ$ grid at 33
 798 standardized depth levels from the surface to 2000 m. The Mapped Observation-Based Oceanic Dissolved
 799 Inorganic Carbon (MOBO-DIC) mapping method adapts and extends the SOM-FFN technique originally
 800 introduced by Landschützer et al. (2013). It starts by categorizing the ocean into clusters with comparable
 801 physical and biogeochemical characteristics using self-organizing maps (SOM). Subsequently, within each
 802 SOM-defined cluster, a feed-forward network (FFN) is employed to estimate and enforce the statistical
 803 correlation between the targeted dissolved inorganic carbon (DIC) data and the predictor data available in



globally mapped fields. The product uses data from January 2004 to December 2017, and is thus centered around the years 2010/2011. The data product is available at OCADS (DOI: “10.25921/yvzj-zx46”):
https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/ndp_104/ndp104.html.

45) Monthly global interior ocean TA climatology: Broullón et al. (2019) developed a monthly global interior ocean TA climatology using a feed-forward neural network approach. This dataset offers a spatial resolution of $1^\circ \times 1^\circ$ in the horizontal, spans 102 depth levels (ranging from 0–5500 m) in the vertical dimension, and features a temporal resolution that varies from monthly (0–1500 m) to annual (1550–5500 m). The development of this climatology was based on the analysis of TA in relation to several key predictor variables, including temperature, salinity, nutrients (phosphate, nitrate, and silicate), dissolved oxygen, and sampling position (coordinates and depth), as outlined in Velo et al. (2013). Both TA and these predictor variables were sourced from GLODAPv2 (version 2016) (Olsen et al., 2016). The global interior ocean TA climatology was constructed by leveraging the established relationships between TA and the predictor variables, as well as the monthly climatologies of temperature, salinity, and dissolved oxygen from the World Ocean Atlas 2013 (WOA13) (Locarnini et al., 2013; Zweng et al., 2013; Garcia et al., 2014), and nutrients data that were obtained through the CANYON-B neural network process, applied to the previously mentioned fields. The data product is available at OCADS (DOI: “10.25921/5p69-y471”):
<https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0222470.html>.

46) Monthly global interior ocean DIC climatology: Broullón et al. (2020) employed a feed-forward neural network approach to create a monthly global interior ocean DIC climatology, centered around the year 1995. This dataset offers a $1^\circ \times 1^\circ$ spatial resolution in the horizontal domain, encompassing 102 depth levels ranging from 0–5500 m vertically. The temporal resolution varies, ranging from monthly (0–1500 m) to annual (1550–5500 m). In contrast to their previous work on TA (Broullón et al., 2019), this analysis includes the variable “year” to account for anthropogenic DIC pool changes. It also incorporates data from the Lamont–Doherty Earth Observatory (LDEO) $p\text{CO}_2$ database (Takahashi et al., 2017) alongside GLODAPv2.2019 (Olsen et al., 2019) to establish relationships between DIC and its input variables: temperature, salinity, dissolved oxygen, as well as location, pressure, and time. The DIC climatology was derived using these relationships, along with monthly climatological data for temperature, salinity, and dissolved oxygen from the World Ocean Atlas 2013 (Locarnini et al., 2013; Zweng et al., 2013; Garcia et al., 2014), as well as phosphate, nitrate, and silicate values computed from the CANYON-B neural network fed with the aforementioned fields. The data product is available at OCADS (DOI: “10.25921/ndgj-jp24”):
<https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0222469.html>.

47) MOBO-DIC (Version 2023): Keppler et al. (2023) extended the temporal resolution of MOBO-DIC to resolve monthly fields from January 2004 to December 2019, as opposed to the average climatological values in Keppler et al. (2020). This data product is on a $1^\circ \times 1^\circ$ grid at 28 depth levels from the surface to 1500 m. The data product is available at OCADS (DOI: “10.25921/z31n-3m26”):



839 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0277099.html>.

840 **48) Metrics of acidification in the ocean interior:** Fassbender et al. (2023) generated estimates of global
 841 interior ocean changes to pH, $[H^+]$, Ω_{Ar} , pCO_2 , and the Revelle sensitivity factor driven by the
 842 accumulation of anthropogenic carbon (C_{ant}) from the preindustrial period to the year 2002, and quantified
 843 the component of those changes caused by nonlinearities in the carbonate system. For each OA metric, the
 844 dataset includes year 2002 values and quasi-preindustrial values, which were estimated by subtracting C_{ant}
 845 from the year 2002 carbonate chemistry information and recomputing each OA metric without considering
 846 any warming, circulation, or biological changes that may have occurred since the preindustrial era. Data
 847 from the upper 2000 m of the GLODAPv2.2016b mapped data product (DOI:
 848 “10.3334/cdiac/otg.ndp093_glodapv2”), described in Lauvset et al., 2016, and from the preformed
 849 properties product of Carter et al., 2021 were used to make these estimates on the GLODAPv2.2016b $1^\circ \times$
 850 1° grid for 26 depth levels from surface to 2000 m. The provided uncertainties were estimated using a
 851 1000-iteration Monte Carlo simulation. Calculation details are described in Fassbender et al. (2023). Year
 852 2002 aragonite saturation state and pH values, and their uncertainties, are reproduced from the
 853 GLODAPv2.2016b mapped data product and are provided in this dataset for user convenience with the
 854 permission of the original data producer. This data product is available at OCADS (DOI: “10.25921/rdrtr-
 855 9t74”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0290073.html>.

856 **49) Anthropogenic CO_2 from 1994 to 2007:** Gruber et al. (2019a) estimated decadal changes in the oceanic
 857 content of anthropogenic CO_2 (ΔC_{ant}) for the years 1994, 2004, and 2014. The results have been derived
 858 from the GLODAPv2.2016 product (Olsen et al., 2016), utilizing the eMLR(C*) methodology pioneered by
 859 Clement and Gruber (2018). All estimates are geospatially distributed on a horizontal grid with a resolution
 860 of $1^\circ \times 1^\circ$. Two primary files are available: one providing the complete three-dimensional distribution of
 861 ΔC_{ant} , and the other containing vertically integrated values, i.e., the column inventories. This data product is
 862 available at OCADS (DOI: “10.25921/wdn2-pt10”):
 863 <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0186034.html> (Gruber et al., 2019b) .

864 **50) Decadal trends in anthropogenic CO_2 from 1994 to 2014:** Müller et al. (2023) extended the analysis by
 865 Gruber et al. (2019a) to reconstruct decadal trends in the oceanic storage of anthropogenic CO_2 (ΔC_{ant}) in
 866 the global ocean interior from 1994 to 2014. They applied the extended multiple linear regression (eMLR)
 867 method (Clement and Gruber 2018) to ship-borne observations of DIC and other biogeochemical variables
 868 from GLODAPv2.2021 (Lauvset et al., 2021). All estimates are provided on a $1^\circ \times 1^\circ$ horizontal grid. Two
 869 principal data files are provided: one featuring the comprehensive three-dimensional distribution of ΔC_{ant} ,
 870 and the other presenting the vertically integrated quantities, i.e., the column inventories. The data product is
 871 available at OCADS (DOI: “10.25921/ppcf-w020”): [https://www.ncei.noaa.gov/data/oceans/ncei/ocads/](https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0279447.html)
 872 [metadata/0279447.html](https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0279447.html).



- 873 **51) Progression of Ocean Interior Acidification over the Industrial Era:** Building on the total
 874 anthropogenic carbon estimates for 1994 from Sabine et al. (2004) and the decadal changes between 1994
 875 and 2014 reconstructed by Müller et al. (2023), Müller et al. (2024) quantified ocean interior acidification
 876 over the industrial era. To convert the increasing anthropogenic carbon concentrations into acidification
 877 estimates, their approach relied on time-invariant climatologies of ocean interior DIC, TA, temperature,
 878 salinity, and other relevant variables to determine the background state of the marine carbonate system.
 879 Hence, their estimates resolve exclusively the acidification driven by the anthropogenic carbon
 880 accumulation. In contrast to direct observations of acidification parameters—such as those collected at time
 881 series stations—this approach does not account for changes in the natural carbon cycle or the displacement
 882 of water masses. The approach by Müller et al. (2024) is conceptually similar to that of Fassbender et al.
 883 (2023), but provides temporally resolved estimates, enabling the tracking of both the spatial distribution
 884 and temporal evolution of ocean interior acidification. The data product is available at OCADS (DOI:
 885 “10.25921/tefm-x802”): <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0298993.html>
- 886 **52) CODAP-NA climatology:** Jiang et al. (2024) developed a coastal OA indicators climatology on a $1^\circ \times 1^\circ$
 887 grid, covering North American ocean margins from surface to 500 m at 14 standardized depth levels: 0, 10,
 888 20, 30, 50, 75, 100, 125, 150, 200, 250, 300, 400, and 500 m. This product includes 10 key oceanographic
 889 parameters: fugacity of carbon dioxide ($f\text{CO}_2$), pH, total hydrogen ion content, free hydrogen ion content,
 890 carbonate ion content ($[\text{CO}_3^{2-}]$), Ω_{arag} , Ω_{calc} , DIC, TA, and Revelle Factor (RF), as well as temperature and
 891 salinity. The climatology was produced with the World Ocean Atlas (WOA) gridding technologies of the
 892 NOAA National Centers for Environmental Information (NCEI), based on the recently released Coastal
 893 Ocean Data Analysis Product in North America (CODAP-NA) (Jiang et al., 2021), along with
 894 GLODAPv2.2022 (Lauvset et al., 2022). The relevant variables were adjusted to the year of 2010 before
 895 the gridding. The first-guess fields for this analysis were calculated using ESPERs (Carter et al., 2021),
 896 based on the WOA (Version 2018) climatologies for salinity (Zweng et al., 2019), temperature (Locarnini
 897 et al., 2019) and dissolved oxygen (Garcia et al., 2018b). The data product is available in NetCDF at
 898 OCADS (DOI: “10.25921/g8pb-zy76”):
 899 <https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0270962.html>. Additionally, maps of these
 900 indicators are available in jpeg at: [https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/synthesis/nacoastal.html)
 901 [system/synthesis/nacoastal.html](https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/synthesis/nacoastal.html) (Jiang et al., 2022b).



902 **Table 4. Gridded and derived ocean carbonate chemistry data synthesis products in the subsurface ocean.**

No.	Name	Open or coastal ocean	Resolution	Temporal resolution	Methodology	Highlights	Reference
42	Global interior ocean mapped climatology from GLODAPv2	Open ocean	1° × 1°	adjusted to 2002	Data Interpolating Variational Analysis (DIVA)	Ocean interior climatology for multiple variables from surface to the bottom of the ocean (referenced to year 2002)	Lauvset et al. (2016)
43	Global aragonite saturation state climatology	Open ocean	1° × 1°	adjusted to 2000	Data Interpolating Variational Analysis (DIVA)	Ocean interior climatology for aragonite saturation state from surface to 4000 m (referenced to year 2000)	Jiang et al. (2015a)
44	MOBO-DIC (Version 2020)	Open ocean	1° × 1°	monthly climatology, centered around 2010/2011	Machine learning	Seasonal variability of DIC in the interior ocean from surface to 2000 m	Keppler et al. (2020)
45	Monthly global interior ocean TA climatology	Open ocean	1° × 1°	monthly climatology	Machine learning	Ocean interior climatology for TA from surface to bottom	Broullón et al. (2019)
46	Monthly global interior ocean DIC climatology	Open ocean	1° × 1°	monthly from 1957 to 2018	Machine learning	Ocean interior climatology for DIC from surface to bottom (referenced to year 1995)	Broullón et al. (2020)
47	MOBO-DIC (Version 2023)	Open ocean	1° × 1°	monthly from Jan 2004 to Dec 2019	Machine learning	Temporal trends and interannual variability of DIC in the interior ocean from surface to 1500 m	Keppler et al. (2023)
48	Metrics of acidification in the ocean interior	Open ocean	1° × 1°	2002 and preindustrial	Reproduced from GLODAPv2.2016b (Lauvset et al., 2016) and the preformed properties of Carter et al., (2021)	Metrics of acidification in the ocean interior (to 2000 m) and the component of those changes caused by carbonate system nonlinearities	Fassbender et al. (2023)
49	Anthropogenic CO ₂ from 1994 to 2007	Open ocean	1° × 1°	1994 and 2007	Extended multiple linear regression (eMLR)	The oceanic sink for anthropogenic CO ₂ over the period 1994 to 2007	Gruber et al. (2019a)
50	Decadal trends in anthropogenic CO ₂ From 1994 to 2014	Open ocean	1° × 1°	Decadal from 1994 to 2014	eMLR(C*) extended Multiple Linear Regression applied to the tracer C*	Temporal trends in the accumulation of anthropogenic CO ₂ in the interior ocean are resolved	Müller et al. (2023)
51	Progression of Ocean Interior Acidification over the Industrial Era	Open ocean	1° × 1°	1800 – 1994 – 2004 - 2014	Conversion of anthropogenic carbon accumulation into acidification rates	Temporal trends in the progression of acidification in the interior ocean are resolved	Müller et al. (2024)



52	CODAP-NA climatology	Coastal	$1^{\circ} \times 1^{\circ}$	adjusted to 2010	Objective analysis approach of the World Ocean Atlas	The first discrete bottle based climatology in the North American ocean margins	Jiang et al. (2024)
----	----------------------	---------	------------------------------	------------------	--	---	---------------------

903

904 3.1.5 Multi-product analyses:

905 This section includes data products that have been generated by community synthesis efforts designed to inform
 906 global carbon budgets.

907 **53) SeaFlux:** Harmonization of air–sea CO₂ fluxes from surface *p*CO₂ data products using a standardized
 908 approach (Gregor and Fay, 2021). This resource provides an ensemble of six *p*CO₂ products with air-sea
 909 CO₂ fluxes computed consistently. The six included products are: CMEMS-FFNN, CSIR-ML6, JENA-
 910 MLS, JMA-MLR, MPI-SOMFFN, and NIES-FNN. First, missing areas of *p*CO₂ estimates (mostly high-
 911 latitude and marginal seas) are filled using a linear-regression approach, thus addressing differences in
 912 spatial coverage between the mapping products. Further, also accounts for methodological inconsistencies
 913 in flux calculations. Fluxes are calculated using three wind products (CCMPv2, ERA5, and JRA55) along
 914 with the application of a scaled gas exchange coefficient for each of the wind products. Through these
 915 steps, SeaFlux presents an product ensemble of interpolated global surface ocean *p*CO₂ and air–sea carbon
 916 flux estimates for the years 1990–2019. For more details, refer to Fay et al. (2021).

917 **54) RECCAP2:** In the context of the second iteration of the project REgional Carbon Cycle Assessment and
 918 Processes (RECCAP2), the ocean carbon community compiled, quality controlled and harmonized (in the
 919 sense of providing output on the same regular grid at the same spatial and temporal resolution) 12 global
 920 ocean biogeochemical model simulations, 11 *p*CO₂ products, one ocean interior DIC products, and three
 921 data assimilation models to constrain the ocean carbon sink between 1985 and 2018. The RECCAP2
 922 synthesis effort stands as a distinct but complementary resource to the Global Carbon Budget (GCB)
 923 project (Friedlingstein et al., 2025), which primarily focuses on anthropogenically perturbed surface CO₂
 924 fluxes from a global budgeting perspective. The individual chapters of RECCAP2 were published in this
 925 special issue of Global Biogeochemical Cycles:
 926 [https://agupubs.onlinelibrary.wiley.com/doi/toc/10.1002/\(ISSN\)2169-8961.RECCAP2](https://agupubs.onlinelibrary.wiley.com/doi/toc/10.1002/(ISSN)2169-8961.RECCAP2). The data products
 927 of this assessment are available on a $1^{\circ} \times 1^{\circ}$ horizontal grid, with monthly resolution for surface ocean
 928 variables such as air-sea CO₂ fluxes, and annual resolution for interior ocean variables, such as DIC
 929 content. The data compilation, which is described in detail in DeVries et al. (2023), is available at:
 930 <https://zenodo.org/records/7990823> (Müller, 2023).

931 **55) Global Carbon Budget:** The GCB collects annually updated estimates of the ocean carbon sink from
 932 currently nine *f*CO₂-products and ten Global Ocean Biogeochemical Models for the period 1959 to the past
 933 calendar year (Friedlingstein et al., 2025, <https://globalcarbonbudget.org/gcb-2024>). In contrast to Earth
 934 System Models, the GOBMs are here forced with atmospheric reanalysis that ingested atmosphere and



ocean observations and are thus thought to be closer to the observed climate. Gridded fields are provided on a $1^\circ \times 1^\circ$ horizontal grid and monthly resolution. In addition, globally and regionally integrated air-sea CO_2 fluxes from the native model grids are provided. Globally integrated time-series are adjusted for full ocean coverage and model bias and drift and are available for each individual $f\text{CO}_2$ -product and GOBM (<https://globalcarbonbudget.org/download/1442/?tmstv=1731323337>). The model data goes well beyond surface fluxes and includes data to analyze drivers of carbon fluxes, including several 3D variables. The model data request has been updated since RECCAP2 and also provides, for example, monthly interior ocean data of DIC, alkalinity, nutrients and oxygen. The GOBM data request was also updated to have all variables available that are needed to serve as a testbed for $f\text{CO}_2$ -products (e.g., sea surface height). Gridded surface data of sea surface fugacity and air-sea CO_2 flux of all $f\text{CO}_2$ -products and GOBMs as used in the latest release of GCB (2024) are published on Zenodo (<https://zenodo.org/records/14639761>, Hauck et al., 2025). All other GOBM output is available via <https://globalcarbonbudgetdata.org/closed-access-requests.html>.

Table 5. Ocean carbonate chemistry data product synthesis and harmonizations.

No.	Name	Open or coastal ocean	Surface or water column	Spatial resolution	Temporal resolution	Methodology	Highlights	Reference
53	SeaFlux	Open ocean + coastal	Surface only	$1^\circ \times 1^\circ$	Monthly	Consistent flux calculations for 6 $p\text{CO}_2$ products to produce an ensemble estimate	Careful consideration of flux calculation provides a resource and code to the community for independent flux calculations	Gregor and Fay (2021)
54	RECCAP2	Open ocean + coastal	Surface + Water column	$1^\circ \times 1^\circ$ (open) $0.25^\circ \times 0.25^\circ$ (coastal)	Monthly	Harmonized compilation of surface $f\text{CO}_2$ products, model simulations and ocean interior products	Quality controlled data compilation with a harmonised horizontal grid and temporal resolution	Müller (2023); DeVries et al. (2023) Resplandy et al (2024)
55	Global Carbon Budget	Open ocean + coastal	Surface + Water column	$1^\circ \times 1^\circ$	Monthly	Harmonized compilation of surface $f\text{CO}_2$ products, and global ocean biogeochemical model simulations	Annually updated and quality controlled data sets. Availability of monthly 4D ocean model output.	Friedlingstein et al., 2025

3.1.6 Model based and hybrid data products and analyses:

Model-based projections of biogeochemical variables are often available from global and regional models, such as those in the Sixth Coupled Model Intercomparison Project (Dunne et al., 2024; Durack et al., 2025). This section further includes hybrid data products, which adjust model estimates towards observation-based constraints.



- 954 **56) Decadal trends in the ocean carbon sink:** The DeVries et al. (2019) analysis examines decadal trends in
 955 global and regional air-sea CO₂ fluxes from a variety of ocean biogeochemical models that contributed to
 956 the Global Carbon Budget (see No. 55). Three sets of model simulations were performed. Simulation A
 957 uses variable climate forcing (e.g., variable wind stress, heat and freshwater fluxes) and observed
 958 atmospheric CO₂ forcing. Simulation B uses constant (repeated) climate forcing and observed atmospheric
 959 CO₂, and simulation C uses both constant climate forcing and constant atmospheric CO₂ concentrations.
 960 With these simulations, the authors partitioned decadal trends in ocean CO₂ uptake into those driven by
 961 climate variability and those driven by atmospheric CO₂. They found that climate variability drove a
 962 weakening trend of the ocean carbon sink during the 1990s, and a strengthening trend during the first
 963 decade of the 2000s. The magnitude of these trends agreed with those of an OCIM that was trained to
 964 replicate tracer data from the 1990s and 2000s (DeVries et al., 2017), indicating that the decadal trends may
 965 be driven by variability in ocean circulation. This data from this analysis is accessible through figshare at
 966 <https://doi.org/10.6084/m9.figshare.8091161.v1>.
- 967 **57) ECCO-Darwin:** Carroll et al. (2022) used the Estimating the Circulation and Climate of the Ocean-Darwin
 968 (ECCO-Darwin) global-ocean biogeochemistry state estimate to generate a data-constrained DIC budget
 969 and investigate how spatiotemporal variability in advection and mixing, air-sea CO₂ flux, and the biological
 970 pump have modulated the ocean sink for 1995–2018. ECCO-Darwin assimilates ocean circulation and
 971 physical tracers, including temperature, salinity, and sea ice, derived from the Estimating the Circulation
 972 and Climate of the Ocean (ECCO) LLC270 global-ocean and sea-ice data synthesis (Zhang et al., 2018).
 973 Additionally, it assimilates biogeochemical observations encompassing the cycling of carbon, nitrogen,
 974 phosphorus (PO₄), iron (Fe), silica (SiO₂), oxygen, and alkalinity. This inclusive approach enhances the
 975 model's fidelity by aligning it with a diverse array of observations. All ECCO-Darwin model output is
 976 available on the ECCO Data Portal: <https://data.nas.nasa.gov/ecco/>. The model code and platform-
 977 independent instructions for running ECCO-Darwin simulations can be found at:
 978 https://github.com/MITgcm-contrib/ecco_darwin.
- 979 **58) Global surface ocean pH, acidity, and Revelle factor from 1770 to 2100:** Jiang et al. (2019a) produced a
 980 high-resolution (1°×1°) data product delineating regionally varying view of global surface ocean pH,
 981 acidity, and Revelle Factor (RF) from the beginning of the Industrial Revolution to the end of this century
 982 by amalgamating recent observational seawater CO₂ data from the SOCAT database (Version 6, 1991–
 983 2018, ~23 million observations) (Bakker et al., 2016), and temporal trends at individual locations of the
 984 global surface ocean from an Earth System Model, i.e., GFDL-ESM2M (Dunne et al., 2013). The
 985 calculations were conducted under historical atmospheric CO₂ levels (pre-2005) and four Representative
 986 Concentrations Pathways (post-2005) corresponding to the Intergovernmental Panel on Climate Change
 987 (IPCC)'s 5th Assessment Report, specifically RCP2.6, RCP4.5, RCP6.0, and RCP8.5. Surface ocean TA
 988 was calculated from SSS, SST using the updated locally interpolated alkalinity regression (LIARv2)



method (Carter et al., 2017). Surface ocean pH, acidity, and RF were then calculated using a MATLAB version of the CO2SYS program (Orr et al., 2015). The data product is available at OCADS (DOI: “10.25921/kgqr-9h49”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0206289.html> (Jiang et al., 2019b).

59) Global surface ocean acidification indicators from 1750 to 2100: Jiang et al. (2023) developed a comprehensive model-data fusion product that delineates the trajectory of 10 OA indicators: $f\text{CO}_2$, pH, total hydrogen ion content, free hydrogen ion content, $[\text{CO}_3^{2-}]$, Ω_{arag} , Ω_{calc} , DIC, TA, and RF, as well as temperature and salinity at all locations of the global surface ocean from 1750 to 2100. This product marks a significant breakthrough in OA forecasting by refining temporal trends with data from 14 Earth System Models (ESMs) within the Coupled Model Intercomparison Project Phase 6 (CMIP6), and by applying bias and drift corrections from three updated observational ocean carbon data products: SOCAT (Version 2022) (Bakker et al., 2016), GLODAPv2.2022 (Lauvset et al., 2022), and CODAP-NA (Jiang et al., 2021). This dataset offers 10-year averages on a $1^\circ \times 1^\circ$ global surface ocean grid, capturing trends from preindustrial times (1750), through historical conditions (1850–2010), and projects future conditions to 2100 across five Shared Socioeconomic Pathways: SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP3-7.0, and SSP5-8.5. The gridded data product is available in NetCDF at OCADS (DOI: “10.25921/9ker-bc48”): <https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0259391.html>, and global maps of these indicators are available in jpeg at: <https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/synthesis/surface-oa-indicators.html> (Jiang et al., 2022c).

60) Simulated and constrained ocean carbon sink from 1850 to 2100 for the global ocean and the Southern Ocean: These two datasets include spatially-integrated and annually averaged values for the ocean carbon sink from 1850 to 2100 for different scenarios over the 21st century for the global ocean (Terhaar et al., 2022a, 2022b) and the Southern Ocean (Terhaar et al., 2021b, 2021c). All results are based on CMIP5 and CMIP6 models. For the global ocean carbon sink, values are available for SSP1-2.6, SSP2-4.5, and SSP5-8.5. For the Southern Ocean, values are also available for SSP1-2.6, SSP2-4.5, and SSP5-8.5 and additionally also for RCP2.6, RCP4.5, and RCP8.5. In addition, to the raw simulated values, constrained estimates of the annually averaged ocean carbon sink estimates are available. These constrained estimates adjusted the simulated carbon sink estimates for biases on the ocean’s circulation and surface carbonate chemistry (see Terhaar et al., 2021b, 2022a for details). It is recommended to use the constrained estimates. The datasets are available at <https://doi.org/10.17882/103934> and <https://doi.org/10.17882/103938>.



1020 **Table 6. Model based data synthesis products and analyses for ocean carbonate chemistry.**

No.	Name	Open or coastal ocean	Surface or water column	Spatial resolution	Temporal resolution	Highlights	Reference
56	Decadal trends in the ocean carbon sink	Open ocean	Surface	Variable	Decadal	Climate variability drove weakened ocean CO ₂ uptake in the 1990s, and strengthened CO ₂ uptake in the 2000s	DeVries et al. (2019)
57	ECCO-Darwin	Open + Coastal	Water column	$\frac{1}{3}^{\circ} \times \frac{1}{3}^{\circ}$	3-hourly, daily, and month fields available	Model-data synthesis product based on the Estimating the Circulation and Climate of the Ocean (ECCO) ocean state estimate. Fully-closed, physically-consistent 3-D biogeochemical budgets.	Carroll et al. (2020, 2022, 2024)
58	Global surface ocean pH, acidity, and buffer capacity from 1770 to 2100	Open ocean	Surface	$1^{\circ} \times 1^{\circ}$	Decadal	A model-observation fusion product for pH, acidity and Revelle Factor, leveraging GFDL-ESM2M and SOCATv6	Jiang et al. (2019)
59	Global surface ocean acidification indicators from 1750 to 2100	Open ocean	Surface	$1^{\circ} \times 1^{\circ}$	Decadal	A model-observation fusion product for all major OA indicators, leveraging a consortium of 14 Earth System Models and 3 observational data products	Jiang et al. (2023)
60	Simulated and constrained ocean carbon sink from 1850 to 2100 for the global ocean and the Southern Ocean	Open ocean	Surface	Spatially integrated	Annual (1850-2100)	A constrained estimate of the ocean carbon sink based on the simulated carbon sink from CMIP5 and CMIP6 models and constrained with observations of the ocean physics and carbonate chemistry	Terhaar et al. (2021b, 2022a)

1021 3.2 Overlaps and history

1022 Many of the data products described above exhibit significant overlap in various forms. In some cases, one or more
 1023 products are used to generate new ones, while in others, the same collection-level cruise datasets underpin multiple
 1024 products. There are a few foundational data products, such as GLODAPv2 and SOCAT, which are widely utilized to
 1025 develop other data products, including their respective gridded products (e.g., Lauvset et al., 2016). For instance,
 1026 SOCAT forms the backbone of nearly all derived products listed in Table 3, serving as a key resource for product
 1027 development or validation. Some derived products, such as the JMA-MLR (No. 25) and OceanSODA-ETHZv1 (No.
 1028 26), incorporate both SOCAT and GLODAPv2 during development. Having overlaps in data and derived products
 1029 has provided opportunities for data quality control and intercomparison of different approaches to gap filling that
 1030 would not have been available otherwise. Additional overlaps between these data products are provided below:



1031 3.2.1 SOCAT and LDEO:

1032 The quality control and synthesis of global surface ocean CO₂ data began in 1997 with Dr. Taro Takahashi and his
 1033 colleagues at the Lamont-Doherty Earth Observatory (LDEO) in Palisades, New York. His pioneering work led to
 1034 the creation of the LDEO Surface *p*CO₂ Database (No. 2), which focused on high-quality data collected by his team
 1035 and from various U.S. and international expeditions. Over time, this data set expanded to include contributions from
 1036 other laboratories, resulting in a highly influential collection of *p*CO₂ data and several seminal papers on global
 1037 surface ocean CO₂ variations and air-sea CO₂ fluxes (Takahashi et al., 1997; 2002; 2009). The last update to the
 1038 LDEO database was in 2019, following Dr. Takahashi's passing, and no further updates are anticipated (Takahashi
 1039 et al., 2021).

1040 The Surface CO₂ Atlas (SOCAT) project was developed to address questions around the current and future drivers
 1041 of CO₂ fluxes raised at the 2007 Surface Ocean CO₂ Variability and Vulnerability (SOCOVV) workshop in Paris,
 1042 France (Metzl et al., 2007). SOCAT was developed to synthesize all of the publicly available, discoverable, and
 1043 citable surface CO₂ data. Following the GLODAP model, there was a strong emphasis on an open and transparent
 1044 secondary quality control process to ensure the highest data quality. The first data release came in 2011 (Pfeil et al.,
 1045 2013; Sabine et al, 2013) and included contributions from numerous laboratories, as well as the freely available CO₂
 1046 data from the LDEO database. As of 2024, SOCAT contains ~40 million data points, with new observations added
 1047 annually. All data are rigorously standardized, and recalculated as *f*CO₂. SOCAT represents an ongoing global
 1048 community effort, with participants from all continents contributing data and to the quality control process. Initially
 1049 new versions were released every other year, but automation allowed annual public releases since version 4.

1050 3.2.2 GLODAPv2 and Java OceanAtlas (JOA) Suite:

1051 Starting in the late 1980s, the World Ocean Circulation Experiment (WOCE), Joint Global Ocean Flux Study
 1052 (JGOFS), and the NOAA Ocean-Atmosphere Exchange Study (OACES) collaborated in a multinational effort to
 1053 conduct a decadal global hydrographic survey of unparalleled quality and quantity. At the conclusion of the survey
 1054 at the end of the 1990s, the Global Ocean Data Analysis Project (GLODAP) combined and publicly released all of
 1055 the available hydrographic data with high-quality carbon measurements as a single database (Key et al., 2004;
 1056 Sabine et al., 2005). The data were subjected to extensive secondary quality control checks where cruise tracks
 1057 intersected one another, making it the most comprehensive and highest-quality ocean inorganic carbon dataset ever
 1058 generated. A gridded, full-depth global ocean carbon climatology was also created and released as part of the
 1059 project. These data and associated climatology have been extensively used to evaluate carbon distributions as well as
 1060 the accumulation of anthropogenic CO₂ in the ocean. Other regional datasets, like CARINA (CARbon dioxide IN
 1061 the Atlantic Ocean) data synthesis project, an international collaborative effort of the European Union
 1062 CARBOOCEAN program (Key et al., 2010; Tanhua et al., 2010) and PACIFICA (PACIFIC ocean Interior CARbon),
 1063 an international synthesis of Pacific Ocean data organized through the North Pacific Marine Science Organization
 1064 (PICES) (Ishii et al., 2011b; Suzuki et al., 2013), were combined with GLODAP after its initial release. The



1065 GLODAP database is continuing to grow with new data collected as part of the Global Ocean Ship-Based
 1066 Hydrographic Investigations Program (GO-SHIP).

1067 For discrete bottle measurements spanning the entire oceanic water column, GLODAPv2 (No. 3) and the Java
 1068 Ocean Atlas (JOA, No. 4) are the primary data products. Most cruise datasets contributing to these two data products
 1069 overlap, but the key difference lies in their approach to data adjustment. The former applies crossover and inversion
 1070 analysis for bias correction, while the latter presents the data without such adjustments. GLODAPv2 achieves
 1071 consistency by applying adjustments based on deep-ocean offsets, whereas JOA provides the data in its original
 1072 form. While there is substantial overlap between the two, data from a specific expedition might differ slightly due to
 1073 GLODAPv2's secondary quality control adjustments. Both GLODAPv2 and JOA offer global coverage, but several
 1074 independent regional data products are also available, such as SNAPO-CO₂ (No. 6), CODAP-NA (No. 8), AZMP
 1075 Carbon (No. 9), MOCHA (No. 10), and ARIOS (No. 11). Data from these regional products often partially or fully
 1076 overlap with GLODAPv2 and JOA.

1077 **a) GLODAPv2 and CODAP-NA:**

1078 All cruise datasets contributing to CODAP-NA were forwarded to the GLODAPv2 quality control team in 2022.
 1079 Data from select cruises with deep-water sampling (>1500 m), enabling crossover analysis, were subsequently
 1080 incorporated into the GLODAPv2.2022 data product update (Lauvset et al., 2022).

1081 **b) GLODAPv2 and SPOTS:**

1082 Some time-series data are included in both GLODAPv2 and the Synthesis Product for Ocean Time-Series (SPOTS).
 1083 Usually, data present in both products have not been measured on dedicated time-series cruises but rather collected
 1084 as part of a larger cruise passing by a time series location. As the quality control of SPOTS is restricted to assigning
 1085 method flags, adjustments that are applied as a result of the QC of GLODAP are not present in SPOTS. Additional
 1086 crossover analyses between SPOTS and GLODAP have revealed a good consistency (Lange et al., 2024).

1087 **3.2.3 RECCAP2 and GCB:**

1088 RECCAP2 and GCB are not data products themselves, but analyses and syntheses of data-based and model-based
 1089 products. Users should be aware that there is a large degree of overlap between the $f\text{CO}_2$ - products and GOBMs that
 1090 contributed to both RECCAP2 and GCB. However, RECCAP2 and GCB serve different purposes. GCB is updated
 1091 annually to the latest complete calendar year and its main purpose is to present and estimate of the magnitude (and
 1092 uncertainty) of the ocean CO₂ sink for that year, while RECCAP2 is presents a deeper analysis of the magnitude,
 1093 trends, and variability of the ocean CO₂ sink over the period 1985-2018.



3.2.4 Jiang et al. (2019) and Jiang et al. (2023):

Both products contain the projection of surface ocean pH, total hydrogen ion content, and buffer capacity from 1750 to 2100. However, the former is based on one GFDL model ESM2M, while the latter is based on a consortium of 14 Earth system models, and additional observational data. The latter also contains the projection of seven other OA variables, including carbonate ions, aragonite saturation state, calcite saturation state, $f\text{CO}_2$, DIC, and TA.

4 Data availability

Access links for all data products mentioned in this paper are provided in their respective paragraphs. Additionally, their access links are available in Table 7 below.

Table 7. Access links for the compiled ocean carbonate chemistry data products. N/A is short for not applicable.

No.	Name	Data access link	DOI	Reference
1	SOCAT	https://socat.info/	https://doi.org/10.25921/9w-pn-th28	Bakker et al. (2016, 2024)
2	LDEO Surface $p\text{CO}_2$ Database	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0160492.html	https://doi.org/10.3334/cdiac/otg.ndp088(v2015)	Takahashi et al. (2017)
3	GLODAPv2	https://glodap.info/	https://doi.org/10.25921/zyrq-ht66	Lauvset et al. (2023a, 2024)
4	JOA Suite	https://joa.ucsd.edu/	N/A	Swift and Osborne (2022)
5	WOD	https://www.ncei.noaa.gov/products/world-ocean-database	https://doi.org/10.25923/z885-h264	Mishonov et al. (2024)
6	SNAPCO- CO_2	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0285681.html	https://doi.org/10.17882/95414	Metzl et al. (2023, 2024)
7	Arctic Ocean anthropogenic carbon estimates	https://www.seanoe.org/data/00927/103920/	https://doi.org/10.17882/103920	Terhaar et al. (2024)
8	CODAP-NA	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0219960.html	https://doi.org/10.25921/531n-c230	Jiang et al. (2020, 2021)
9	AZMP Carbon	N/A	https://doi.org/10.20383/102.0673	Cyr et al. (2022)
10	MOCHA	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0277984.html	https://doi.org/10.25921/2vve-fh39	Kennedy et al., (2024)
11	ARIOS	https://digital.csic.es/handle/10261/205135	https://doi.org/10.20350/digitalCSIC/12498	Pérez et al. (2020)
12	Marine inorganic carbon chemistry observations in the northern Gulf of Alaska	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0277034.html	https://doi.org/10.25921/x9sg-9b08	Monacci et al. (2023, 2024)



13	Coral Reef Carbonate Chemistry Off the Florida Keys	https://www.ncei.noaa.gov/access/metadata/landing-page/bin/iso?id=gov.noaa.node:NCRMP-CO3-Atlantic	https://doi.org/10.25921/vfz0-dg77	Manzello et al. (2018)
14	Salish cruise data package and multi-stressor data product	https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/SalishCruise_DataPackage.html , https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/SalishCruises_DataProduct.html	https://doi.org/10.25921/zgk5-ep63 , https://doi.org/10.25921/5g29-q841	Alin et al. (2021, 2023, 2024a, 2024b)
15	BATS	https://demo.bco-dmo.org/project/2124	https://doi.org/10.26008/1912/bco-dmo.894099.4 , https://doi.org/10.26008/1912/bco-dmo.893182.4 , https://doi.org/10.26008/1912/bco-dmo.926534.4 , https://doi.org/10.26008/1912/bco-dmo.893521.6 , https://doi.org/10.26008/1912/bco-dmo.917255.5 , https://doi.org/10.26008/1912/bco-dmo.939210.7 , https://doi.org/10.26008/1912/bco-dmo.3782.6 , https://doi.org/10.26008/1912/bco-dmo.3918.8 , https://doi.org/10.26008/1912/bco-dmo.881861.5	Bates et al., 2024a,b,c,d,e; Johnson et al., 2024a,b,c; Steinberg and Cope, 2024
16	HOT	https://www.bco-dmo.org/project/2101 , https://doi.org/10.5281/zenodo.15060930	https://doi.org/10.1575/1912/bco-dmo.3773.1 , https://doi.org/10.5281/zenodo.15060931	Winn et al. (1994, 1998); Dore et al. (2003, 2009, 2014, 2025); Knor et al. (2023)



17	ESTOC	N/A	https://doi.org/10.1594/PANGAEA.959856 , https://doi.pangaea.de/10.1594/PANGAEA.856590 , https://doi.pangaea.de/10.1594/PANGAEA.856615 , https://doi.pangaea.de/10.1594/PANGAEA.856608 , https://doi.pangaea.de/10.1594/PANGAEA.856616 , https://doi.pangaea.de/10.1594/PANGAEA.856593 , https://doi.pangaea.de/10.1594/PANGAEA.856612 , https://doi.pangaea.de/10.1594/PANGAEA.856614 , https://doi.pangaea.de/10.1594/PANGAEA.856607 , https://doi.pangaea.de/10.1594/PANGAEA.956272	González-Dávila and Santana-Casiano (2023)
18	Point B Time-series	N/A	https://doi.org/10.1594/PANGAEA.727120	Gattuso et al. (2021b)
19	Ny-Ålesund Time-series	N/A	https://doi.org/10.1594/PANGAEA.957028	Gattuso et al. (2023)
20	SPOTS	https://www.bco-dmo.org/dataset/896862	https://doi.org/10.26008/1912/bco-dmo.896862.2	Lange et al. (2024a, 2024b)
21	Autonomous $p\text{CO}_2$ and pH time series from 40 surface buoys	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0173932.html	https://doi.org/10.7289/v5db8043	Sutton et al. (2018)
22	Updated Takahashi delta $f\text{CO}_2$ and flux climatology	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0282251.html	https://doi.org/10.25921/295g-sn13	Fay et al. (2023)
23	MPI-ULB-SOM-FFN	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0209633.html	https://doi.org/10.25921/qb25-f418	Landschützer et al. (2020a, 2020b)
24	VLIZ SOM-FFN	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0160558.html	https://doi.org/10.7289/V5Z899N6	Landschützer et al. (2016), Jersild et al. (2017)
25	JMA-MLR	https://www.data.jma.go.jp/kaiyou/english/co2_flux/co2_flux_data_en.html	N/A	Iida et al. (2021)
26(a)	OceanSODA-ETHZv1	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0220059.html	https://doi.org/10.25921/m5wx-ja34	Gregor et al. (2020)
26(b)	OceanSODA-ETHZv2	N/A	https://doi.org/10.5281/zenodo.11206365	Gregor et al. (2024b)
27	LDEO-HPD $f\text{CO}_2$ product	https://zenodo.org/records/4760205	https://doi.org/10.5281/zenodo.4760205	Gloege et al. (2022)



28	LDEO HPD with extended temporal coverage	https://zenodo.org/records/13891722	https://doi.org/10.5281/zenodo.13891722	Bennington et al. (2022a)
29	LDEO fCO ₂ - Residual Method	https://zenodo.org/records/13941548	https://doi.org/10.5281/zenodo.13941548	Bennington et al. (2022b)
30(a)	CMEMS -LSCEv1	https://data.ipsl.fr/catalog/srv/eng/catalog.search#/metadata/a2f0891b-763a-49e9-af1b-78ed78b16982	https://doi.org/10.14768/a2f0891b-763a-49e9-af1b-78ed78b16982	Chau et al. (2022)
30(b)	CMEMS -LSCEv2	https://data.marine.copernicus.eu/product/MULTIOBS_GLO_BIO_CARBO_N_SURFACE_MYNRT_015_008/services	https://doi.org/10.48670/moi-00047	Chau et al. (2024a, 2024b)
31	CarboScope (Jena-MLS)	https://www.bgc-jena.mpg.de/CarboScope/?ID=oc	10.17871/CarboScope-oc_v2024E (or analogously for previous and upcoming releases)	Rödenbeck et al. (2022)
32	UOEx-Watson	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0301544.html	https://doi.org/10.25921/2dp5-xm29	Watson et al. (2025)
33	NIES-ML3	https://db.cger.nies.go.jp/DL/10.17595/20220311.001.html.en	https://doi.org/10.17595/20220311.001	Zeng (2022), Zeng et al. (2022)
34	CSIR-ML6	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0206205.html	https://doi.org/10.25921/z682-mn47	Gregor et al. (2019b)
35	AOML-ET	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0298989.html	https://doi.org/10.25921/0s8y-q287	Wanninkhof et al. (2024, 2025)
36	ULB-SOM-FFN-coastalv2.1	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0279118.html	https://doi.org/10.25921/4sdep068	Roobaert et al. (2023, 2024)
37	RFR-LME	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0287551.html	https://doi.org/10.25921/h8vw-e872	Sharp et al. (2024a, 2024b)
38	ReCAD -NAACOM-pCO ₂	https://zenodo.org/records/14038561	https://doi.org/10.5281/zenodo.1150097	Wu et al. (2025)
39	Gridded surface OA indicators, and air-sea CO ₂ fluxes in the northern Caribbean Sea	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0207749.html	https://doi.org/10.25921/2swk-9w56	Wanninkhof et al. (2019)
40	OA data in the Gulf of Mexico/Gulf of America and wider Caribbean from 2014 to 2020	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0245950.html	https://doi.org/10.25921/tt1c-dx53	van Hooijdonk (2022)
41	Regional pCO ₂ climatology of the Baltic Sea	N/A	https://doi.org/10.1594/PAN/GAEA.961119	Bittig et al. (2023)



42	Global interior ocean mapped climatology from GLODAPv2	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0286118.html	https://doi.org/10.3334/cdiac/otg.ndp093_glodapv2	Lauvset et al. (2016, 2023b)
43	Global aragonite saturation state climatology	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0139360.html	https://doi.org/10.7289/v5q81b4p	Jiang et al. (2015a, 2015b)
44	MOBO-DIC (Version 2020)	https://www.ncei.noaa.gov/access/ocean-carbon-acidification-data-system/oceans/ndp_104/ndp104.html	https://doi.org/10.25921/yvzj-zx46	Keppler et al. (2020)
45	Monthly global interior ocean TA climatology	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0222470.html	http://doi.org/10.20350/DIGITALCSIC/8564	Broullon et al., (2019)
46	Monthly global interior ocean DIC climatology	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0222469.html	http://doi.org/10.20350/digitalCSIC/10551	Broullon et al., (2020)
47	MOBO-DIC (Version 2023)	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0277099.html	https://doi.org/10.25921/z31n-3m26	Keppler et al. (2023)
48	Metrics of acidification in the ocean interior	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0290073.html	https://doi.org/10.25921/rdtr-9t74	Fassbender et al. (2023), Fassbender (2024)
49	Anthropogenic CO ₂ from 1994 to 2007	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0186034.html	https://doi.org/10.25921/wdn2-pt10	Gruber et al. (2019a, 2019b)
50	Decadal trends in anthropogenic CO ₂ From 1994 to 2014	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0279447.html	https://doi.org/10.25921/ppcf-w020	Müller et al. (2023)
51	Progression of Ocean Interior Acidification over the Industrial Era	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0298993.html	https://doi.org/10.25921/tefm-x802	Müller et al. (2024)
52	CODAP-NA climatology	https://www.ncei.noaa.gov/data/oceans/ncei/ocads/metadata/0270962.html	https://doi.org/10.25921/g8pb-zy76	Jiang et al. (2022b), Jiang et al. (2024)
53	SeaFlux	https://zenodo.org/records/8280457	https://doi.org/10.5281/zenodo.5482547	Gregor & Fay. (2021)
54	RECCAP2	https://zenodo.org/records/7990823	https://doi.org/10.5281/zenodo.7990823	Müller (2023)
55	Global Carbon Budget	https://zenodo.org/records/14639761	https://doi.org/10.5281/zenodo.14639761	Hauck et al. (2025)
56	Decadal trends in the ocean carbon sink	N/A	https://doi.org/10.6084/m9.figshare.8091161.v1	DeVries et al. (2019)
57	ECCO-Darwin	https://data.nas.nasa.gov/ecco/	N/A	Carroll et al. (2020)



58	Global surface ocean pH, acidity, and buffer capacity from 1770 to 2100	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0206289.html	https://doi.org/10.25921/kgqr-9h49	Jiang et al. 2019(a), Jiang et al. (2019b)
59	Global surface ocean acidification indicators from 1750 to 2100	https://www.ncei.noaa.gov/data/oceans/nci/ocads/metadata/0259391.html	https://doi.org/10.25921/9ker-bc48	Jiang et al. (2022c), Jiang et al. (2023)
60	Simulated and constrained ocean carbon sink from 1850 to 2100 for the global ocean and the Southern Ocean	https://www.seanoe.org/data/00927/103934/ , https://www.seanoe.org/data/00927/103938/	https://doi.org/10.17882/103934 , https://doi.org/10.17882/103938	Terhaar et al. (2021c), Terhaar et al. (2022b)

1103

1104 5 Summary

1105 The synthesis and gridded data products presented here reflect significant community-based efforts that have
1106 been made to advance understanding of the ocean's role in global carbon cycling. This synthesis provides a
1107 comprehensive overview of key data compilations and gridded data products essential for coastal and global ocean
1108 carbonate chemistry research. It highlights the key features of each product, serving as a resource for researchers
1109 seeking the necessary data for their work.

1110 Author contributions

1111 L-QJ prepared the initial draft. All authors contributed to the writing of the manuscript. The first 20 authors are
1112 listed based on their contributions, while the remaining authors are listed alphabetically by their last names.

1113 Competing interests

1114 One of the (co-)authors, Anton Velo (Instituto de Investigaciones Mariñas, IIM - CSIC, Vigo, Spain), is a
1115 member of the editorial board of the Earth System Science Data.

1116 Acknowledgements

1117 We extend our gratitude to all scientists who collected and measured the original data and those who compiled
1118 and quality controlled these data products. We thank Bronte Tilbrook (CSIRO Oceans and Atmosphere and
1119 Australian Antarctic Program Partnership, University of Tasmania, Australia), and Dwight Gledhill (NOAA/OAR
1120 Ocean Acidification Program, United States) for their contributions to the writing of this paper. We also thank Pierre
1121 Friedlingstein (University of Exeter, Exeter, UK) for his valuable insights and discussions, which helped shape the



1122 vision of this paper. This is PMEL contribution 5728.

1123 **Financial support**

1124 Funding for L-QJ was from NOAA Ocean Acidification Program (<https://ror.org/02bfn4816>) and NOAA grant
 1125 NA24NESX432C0001 (Cooperative Institute for Satellite Earth System Studies - CISESS) at the Earth System
 1126 Science Interdisciplinary Center, University of Maryland. The authors acknowledge funding from the United States
 1127 National Science Foundation (grant no. OCE-2140395) to the Scientific Committee on Oceanic Research (SCOR,
 1128 United States) for the International Ocean Carbon Coordination Project (IOCCP). SKL and NL acknowledge
 1129 funding by the Horizon Europe research and innovation program under grant agreement 101083922 (OceanICU
 1130 Improving Carbon Understanding). Funding for GLODAPv2 was provided by the EU FP7 project CarboChange
 1131 (grant agreement 264879) and the Research Council of Norway project DECApH (grant agreement 214513), and the
 1132 2019-2023 updates were funded by the EU Horizon 2020 innovation action EuroSea (grant agreement 862626).
 1133 AJF, SRA, and RAF were supported by the NOAA Pacific Marine Environmental Laboratory. GAM was funded by
 1134 NOAA NA24OARX431G0151-T1-01. JPG was supported by the Service d'Observation Rade de Villefranche (SO-
 1135 Rade), the Service d'Observation en Milieu Littoral (SOMLIT/CNRS- INSU), the Coastal Observing System for
 1136 Northern and Arctic Seas (COSYNA), the two Helmholtz large-scale infrastructure projects ACROSS and MOSES,
 1137 the French Polar Institute (IPEV), and the European Commission, Horizon 2020 Framework Programme (grant nos.
 1138 871153, 951799, 727890 and 869154). Discrete samples reported by JPG were analyzed for CT and AT by the
 1139 Service National d'Analyse des Paramètres Océaniques du CO₂. AV and FFP was supported by BOCATS2
 1140 (PID2019-104279GB-C21) project funded by MICIU/AEI/10.13039/501100011033, by FICARAM+ (PID2023-
 1141 148924OB-I00) and by EuroGO-SHIP project (Horizon Europe #101094690). NM and CLM acknowledge Institut
 1142 National des Sciences de l'Univers du Centre National de la Recherche Scientifique (INSU/CNRS) and
 1143 Observatoire des Science de l'Univers (OSU ECCE-Terra) for supporting the SNAPO-CO₂ facility housed by the
 1144 LOCEAN laboratory in Paris/France. The AZMP Carbon dataset is a contribution to Fisheries and Oceans Canada
 1145 Atlantic Zone Monitoring Program. Funding for the HOT program and WHOTS mooring maintenance were
 1146 provided by the National Science Foundation (grant no. OCE-9617409). CMEMS-LSCE is supported by the
 1147 European Copernicus Marine Environment Monitoring Service (CMEMS, grant no. 83-CMEMSTAC-MOB). The
 1148 work of LGr was supported by the ESA OceanHealth-OA project (contract number 4000137603/22/I-DT), the
 1149 European Space Agency (OceanSODA project, grant no. 4000112091/14/I-LG), the European Commission
 1150 (COMFORT project, grant no. 820989), and the Horizon 2020 (4C project, grant no. 821003). JH received funding
 1151 from the Initiative and Networking Fund of the Helmholtz Association (Helmholtz Young Investigator Group
 1152 Marine Carbon and Ecosystem Feedbacks in the Earth System [MarESys], Grant VH-NG-1301), and from the ERC-
 1153 2022-STG OceanPeak (Grant 101077209). AK was funded by the NOAA Global Ocean Monitoring and Observing
 1154 program (<https://ror.org/037bamf06>). J.T. was funded by the Swiss National Science Foundation under grant #
 1155 PZ00P2_209044 (ArcticECO). HCB acknowledges funding by the German Federal Ministry of Education and
 1156 Research under grants no. 03F0877D (C-SCOPE project) and 03F0773A (BONUS INTEGRAL project). TD



acknowledges support from the US National Science Foundation through Grant # 1948955. RW, LB, AJS, SRA, and
 RAF acknowledge support from the Office of Oceanic and Atmospheric Research of NOAA, U.S. Department of
 Commerce, including resources from the Global Ocean Monitoring and Observing Program and the Ocean
 Acidification Program (Open Funder Registry numbers 100018302 and 100018228, respectively). SRA and RAF
 acknowledge support from the Washington Ocean Acidification Center.

References

- Alin, S.R., Newton, J. A., Greeley, D., Curry, B., Herndon, J., Kozyr, A., and Feely, R. A.: A Compiled Data
 Product of Profile, Discrete Biogeochemical Measurements from 35 Individual Cruise Data Sets Collected from a
 Variety of Ships in the Southern Salish Sea and Northern California Current System (Washington State Marine
 Waters) from 2008-02-04 to 2018-10-19 (NCEI Accession 0238424), NOAA National Centers for Environmental
 Information, Dataset, <https://doi.org/10.25921/zgk5-ep63>, 2022.
- Alin, S.R., Newton, J. A., Feely, R. A., Greeley, D., Herndon, J., and Kozyr, A.: A Multi-Stressor Data Product for
 Marine Heatwave, Hypoxia, and Ocean Acidification Research, Including Calculated Inorganic Carbon Parameters
 from the Southern Salish Sea and Northern California Current System from 2008-02-04 to 2018-10-19 (NCEI
 Accession 0283266), NOAA National Centers for Environmental Information, Dataset,
<https://doi.org/10.25921/5g29-q841>, 2023.
- Alin, S. R., Newton, J. A., Feely, R. A., Greeley, D., Curry, B., Herndon, J., and Warner, M.: A decade-long cruise
 time series (2008–2018) of physical and biogeochemical conditions in the southern Salish Sea, North America,
 Earth Syst. Sci. Data, 16, 837–865, <https://doi.org/10.5194/essd-16-837-2024>, 2024a.
- Alin, S. R., Newton, J. A., Feely, R. A., Siedlecki, S., and Greeley, D.: Seasonality and response of ocean
 acidification and hypoxia to major environmental anomalies in the southern Salish Sea, North America (2014–
 2018), Biogeosciences, 21, 1639–1673, <https://doi.org/10.5194/bg-21-1639-2024>, 2024b.
- Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven,
 G., Montenegro, A., and Tokos, K.: Atmospheric lifetime of fossil fuel carbon dioxide, Annu. Rev. Earth Planet.
 Sci., 37(1), 117–134, <https://doi.org/10.1146/annurev.earth.031208.100206>, 2009.
- Archer, D.: The Global Carbon Cycle, Princeton University Press, JSTOR, <https://doi.org/10.2307/j.ctvc4m4hx8>,
 2010.
- Bach, L. T., Gill, S. J., Rickaby, R. E., Gore, S., and Renforth, P.: CO₂ removal with enhanced weathering and ocean
 alkalinity enhancement: Potential risks and co-benefits for marine pelagic ecosystems, Front. clim., 1,
<https://doi.org/10.3389/fclim.2019.00007>, 2019.
- Bakker, D. C. E., Pfeil, B., Smith, K., Hankin, S., Olsen, A., Alin, S. R., Cosca, C., Harasawa, S., Kozyr, A., Nojiri,
 Y., O'Brien, K. M., Schuster, U., Telszewski, M., Tilbrook, B., Wada, C., Akl, J., Barbero, L., Bates, N. R., Boutin,



- 1189 J., Bozec, Y., Cai, W.-J., Castle, R. D., Chavez, F. P., Chen, L., Chierici, M., Currie, K., de Baar, H. J. W., Evans,
 1190 W., Feely, R. A., Fransson, A., Gao, Z., Hales, B., Hardman-Mountford, N. J., Hoppema, M., Huang, W.-J., Hunt,
 1191 C. W., Huss, B., Ichikawa, T., Johannessen, T., Jones, E. M., Jones, S. D., Jutterström, S., Kitidis, V., Körtzinger,
 1192 A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Manke, A. B., Mathis, J. T., Merlivat, L., Metzl, N., Murata, A.,
 1193 Newberger, T., Omar, A. M., Ono, T., Park, G.-H., Paterson, K., Pierrot, D., Ríos, A. F., Sabine, C. L., Saito, S.,
 1194 Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Sieger, R., Skjelvan, I., Steinhoff, T., Sullivan, K. F., Sun, H., Sutton,
 1195 A. J., Suzuki, T., Sweeney, C., Takahashi, T., Tjiputra, J., Tsurushima, N., van Heuven, S. M. A. C., Vandemark,
 1196 D., Vlahos, P., Wallace, D. W. R., Wanninkhof, R., and Watson, A. J.: An update to the Surface Ocean CO₂ Atlas
 1197 (SOCAT version 2), *Earth Syst. Sci. Data*, 6, 69–90, <https://doi.org/10.5194/essd-6-69-2014>, 2014.
- 1198 Bakker, D. C. E., Pfeil, B., Landa, C. S., Metzl, N., O'Brien, K. M., Olsen, A., Smith, K., Cosca, C., Harasawa, S.,
 1199 Jones, S. D., Nakaoka, S., Nojiri, Y., Schuster, U., Steinhoff, T., Sweeney, C., Takahashi, T., Tilbrook, B., Wada,
 1200 C., Wanninkhof, R., Alin, S. R., Balestrini, C. F., Barbero, L., Bates, N. R., Bianchi, A. A., Bonou, F., Boutin, J.,
 1201 Bozec, Y., Burger, E. F., Cai, W.-J., Castle, R. D., Chen, L., Chierici, M., Currie, K., Evans, W., Featherstone, C.,
 1202 Feely, R. A., Fransson, A., Goyet, C., Greenwood, N., Gregor, L., Hankin, S., Hardman-Mountford, N. J., Harlay, J.,
 1203 Hauck, J., Hoppema, M., Humphreys, M. P., Hunt, C. W., Huss, B., Ibáñez, J. S. P., Johannessen, T., Keeling, R.,
 1204 Kitidis, V., Körtzinger, A., Kozyr, A., Krasakopoulou, E., Kuwata, A., Landschützer, P., Lauvset, S. K., Lefèvre, N.,
 1205 Lo Monaco, C., Manke, A., Mathis, J. T., Merlivat, L., Millero, F. J., Monteiro, P. M. S., Munro, D. R., Murata, A.,
 1206 Newberger, T., Omar, A. M., Ono, T., Paterson, K., Pearce, D., Pierrot, D., Robbins, L. L., Saito, S., Salisbury, J.,
 1207 Schlitzer, R., Schneider, B., Schweitzer, R., Sieger, R., Skjelvan, I., Sullivan, K. F., Sutherland, S. C., Sutton, A. J.,
 1208 Tadokoro, K., Telszewski, M., Tuma, M., van Heuven, S. M. A. C., Vandemark, D., Ward, B., Watson, A. J., and
 1209 Xu, S. (2016). A multi-decade record of high-quality /CO₂ data in version 3 of the Surface Ocean CO₂ Atlas
 1210 (SOCAT), *Earth Syst. Sci. Data*, 8, 383–413, <https://doi.org/10.5194/essd-8-383-2016>.
- 1211 Bakker, D. C. E., Alin, S. R., Bates, N., Becker, M., Gkritzalis, T., Jones, Steve D., Kozyr, A., Lauvset, S. K., Metzl,
 1212 N., Nakaoka, S.-i., O'Brien, K. M., Olsen, A., Pierrot, D., Steinhoff, T., Sutton, A. J., Takao, S., Tilbrook, B., Wada,
 1213 C., Wanninkhof, R., Akl, J., Arbilla, L. A., Arruda, R., Azetsu-Scott, K., Barbero, L., Beatty, C. M., Berghoff, C. F.,
 1214 Bittig, H. C., Burger, E. F., Campbell, K., Cardin, V., Collins, A., Coppola, L., Cronin, M., Cross, J. N., Currie, K.
 1215 I., Emerson, S. R., Enright, M. P., Enyo, K., Evans, W., Feely, R. A., Flohr, A., Gehrung, M., Glockzin, M.,
 1216 González-Dávila, M., Hamnca, S., Hartman, S., Howden, S. D., Kam, K., Kamb, L., Körtzinger, A., Kosugi, N.,
 1217 Lefèvre, N., Lo Monaco, C., Macovei, V. A., Maenner Jones, S., Manalang, D., Martz, T. R., Mdokwana, B.,
 1218 Monacci, N. M., Monteiro, P. M. S., Mordy, C., Morell, J. M., Murata, A., Neill, C., Noh, J.-H., Nojiri, Y., Ohman,
 1219 M. D., Olivier, O., Ono, T., Petersen, W., Plueddemann, A. J., Prytherch, J., Rehder, G., Rutgersson, A., Santana-
 1220 Casiano, J. M., Schlitzer, R., Send, U., Skjelvan, I., Sullivan, K. F., T'Jampens, M., Tadokoro, K., Telszewski, M.,
 1221 Theetaert, H., Tsanwani, M., Vandemark, D., van Ooijen, E., Vecchia, M. H., Voynova, Y. G., Wang, H., Weller, R.
 1222 A., Woosley, R. J. (2024). Surface Ocean CO₂ Atlas Database Version 2024 (SOCATv2024) (NCEI Accession
 1223 0293257). NOAA National Centers for Environmental Information. Dataset. <https://doi.org/10.25921/9wpn-th28>,



- 1224 2024.
- 1225 Barth, A., Beckers, J.-M., Troupin, C., Alvera-Azcárate, A., and Vandenbulcke, L.: DIVAnd-1.0: n-dimensional
- 1226 variational data analysis for ocean observations, *Geosci. Model Dev.*, 7, 225-241, <https://doi.org/10.5194/gmd-7->
- 1227 225-2014, 2014.
- 1228 Bates, N.R., and Johnson, R.J.: Forty years of ocean acidification observations (1983-2023) in the Sargasso Sea at
- 1229 the Bermuda Atlantic Time-series Study (BATS) site, *Frontiers in Marine Science*, 10,
- 1230 <https://doi.org/10.3389/fmars.2023.1289931>, 2023.
- 1231 Bates, N., Johnson, R. J., Lomas, M.W., Steinberg, D.K., Lopez, P., Hayden, M., May, R., Derbyshire, L., Lethaby,
- 1232 P. J., Smith, D., and Lomas, D.: Determination of carbon, nitrogen, and phosphorus content in sinking particles at
- 1233 the Bermuda Atlantic Time-series Study (BATS) site from December 1988 to December 2023 using a Particle
- 1234 Interceptor Trap System (PITS). Biological and Chemical Oceanography Data Management Office (BCO-DMO).
- 1235 (Version 4) Version Date 2024-11-14 [if applicable, indicate subset used]. doi:10.26008/1912/bco-dmo.894099.4
- 1236 [access date], 2024a.
- 1237 Bates, N., Johnson, R. J., Lethaby, P. J., Smith, D., and Medley, C.: Primary productivity estimates from the
- 1238 incubation of seawater collected at the Bermuda Atlantic Time-series Study (BATS) site from December 1988
- 1239 through December 2023. Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version
- 1240 4) Version Date 2024-11-14 [if applicable, indicate subset used]. doi:10.26008/1912/bco-dmo.893182.4 [access
- 1241 date], 2024b.
- 1242 Bates, N., Johnson, R. J., Lethaby, P. J., Smith, D., and Medley, C.: HPLC and fluorometric derived phytoplankton
- 1243 pigment concentrations from seawater collected on BATS Validation cruises from June 1996 to June 2023.
- 1244 Biological and Chemical Oceanography Data Management Office (BCO-DMO). (Version 4) Version Date 2024-09-
- 1245 18 [if applicable, indicate subset used]. doi:10.26008/1912/bco-dmo.926534.4 [access date], 2024c.
- 1246 Bates, N., Johnson, R. J., Lethaby, P. J., Medley, C., and Smith, D.: HPLC and fluorometric derived phytoplankton
- 1247 pigment concentrations from seawater collected at the Bermuda Atlantic Time-series Study (BATS) site from
- 1248 October 1988 through December 2023. Biological and Chemical Oceanography Data Management Office (BCO-
- 1249 DMO). (Version 6) Version Date 2023-09-13 [if applicable, indicate subset used]. doi:10.26008/1912/bco-
- 1250 dmo.893521.6 [access date], 2024d.
- 1251 Bates, N., Johnson, R. J., Smith, D., Lethaby, P. J., Lomas, M.W., and Lomas, D.: Discrete bottle samples collected
- 1252 during BATS Validation (BVAL) cruises from April 1991 through June 2023. Biological and Chemical
- 1253 Oceanography Data Management Office (BCO-DMO). (Version 5) Version Date 2024-10-01 [if applicable, indicate
- 1254 subset used]. doi:10.26008/1912/bco-dmo.917255.5 [access date], 2024e.
- 1255 Bennington, V., Gloege, L., and McKinley, G. A.: Variability in the global ocean carbon sink from 1959 to 2020 by
- 1256 correcting models with observations, *Geophys. Res. Lett.*, 49(14), <https://doi.org/10.1029/2022gl098632>, 2022a.



- 1257 Bennington, V., Galjanic, T., and McKinley, G. A.: Explicit physical knowledge in machine learning for Ocean
 1258 Carbon Flux Reconstruction: The $p\text{CO}_2$ -residual method. *J. Adv. Model. Earth Syst.*, 14(10),
 1259 <https://doi.org/10.1029/2021ms002960>, 2022b.
- 1260 Bertin, C., Carroll, D., Menemenlis, D., Dutkiewicz, S., Zhang, H., Matsuoka, A., Tank, S., Manizza, M., Miller, C.
 1261 E., Babin, M., Mangin, A., and Le Fouest, V.: Biogeochemical River runoff drives intense coastal Arctic Ocean CO_2
 1262 outgassing, *Geophys. Res. Lett.*, 50(8), <https://doi.org/10.1029/2022gl102377>, 2023.
- 1263 Bittig, H. C., Steinhoff, T., Claustre, H., Fiedler, B., Williams, N. L., Sauzède, R., Körtzinger, A., and Gattuso, J.-P.:
 1264 An alternative to static climatologies: Robust estimation of open ocean CO_2 variables and nutrient concentrations
 1265 from T, S, and O_2 data using Bayesian Neural Networks, *Frontiers in Marine Science*, 5,
 1266 <https://doi.org/10.3389/fmars.2018.00328>, 2018.
- 1267 Bittig, H. C., Jacobs, E., Neumann, T., and Rehder, G.: A regional $p\text{CO}_2$ climatology of the Baltic Sea, PANGAEA
 1268 [data set], <https://doi.org/10.1594/PANGAEA.961119>, 2023.
- 1269 Bittig, H.C., Jacobs, E., Neumann, T., and Rehder, G.: A regional $p\text{CO}_2$ climatology of the Baltic Sea from in situ
 1270 $p\text{CO}_2$ observations and a model-based extrapolation approach, *Earth Syst. Sci. Data*, 16(2), 753–773,
 1271 <https://doi.org/10.5194/essd-16-753-2024>, 2024.
- 1272 Boyer, T. P., Garcia, H. E., Locarnini, R. A., Zweng, M. M., Mishonov, A. V., Reagan, J. R., Weathers, K. A.,
 1273 Baranova, O. K., Seidov, D., Smolyar, I. V.: World Ocean Atlas 2018, NOAA National Centers for Environmental
 1274 Information, <https://www.ncei.noaa.gov/archive/accession/NCEI-WOA18>, 2018.
- 1275 Brett, A., Leape, J., Abbott, M., Sakaguchi, H., Cao, L., Chand, K., Golbuu, Y., Martin, T. J., Mayorga, J., and
 1276 Myksovoll, M. S.: Ocean Data Need a sea change to help navigate the Warming World. *Nature*, 582(7811), 181–183,
 1277 <https://doi.org/10.1038/d41586-020-01668-z>, 2020.
- 1278 Broecker W. S.: Glacial to interglacial changes in ocean chemistry, *Prog. Oceanogr.*, 11(2), 151–97,
 1279 [https://doi.org/10.1016/0079-6611\(82\)90007-6](https://doi.org/10.1016/0079-6611(82)90007-6), 1982.
- 1280 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., González-
 1281 Dávila, M., Jeansson, E., Kozyr, A., and van Heuven, S. M. A. C.: A global monthly climatology of total alkalinity:
 1282 a neural network approach, *Earth Syst. Sci. Data*, 11, 1109–1127, <https://doi.org/10.5194/essd-11-1109-2019>, 2019.
- 1283 Broullón, D., Pérez, F. F., Velo, A., Hoppema, M., Olsen, A., Takahashi, T., Key, R. M., Tanhua, T., Santana-
 1284 Casiano, J. M., and Kozyr, A.: A global monthly climatology of oceanic total dissolved inorganic carbon: A Neural
 1285 Network approach, *Earth Syst. Sci. Data*, 12(3), 1725–1743, <https://doi.org/10.5194/essd-12-1725-2020>, 2020.
- 1286 Cai, W.-J., Hu, X., Huang, W.-J., Murrell, M. C., Lehrter, J. C., Lohrenz, S. E., Chou, W.-C., Zhai, W., Hollibaugh,
 1287 J. T., Wang, Y., Zhao, P., Guo, X., Gundersen, K., Dai, M., and Gong, G.-C.: Acidification of subsurface coastal



- 1288 waters enhanced by eutrophication, *Nature Geoscience*, 4(11), 766–770, <https://doi.org/10.1038/ngeo1297>, 2011.
- 1289 Carroll, D., Menemenlis, D., Adkins, J. F., Bowman, K. W., Brix, H., Dutkiewicz, S., Fenty, I., Gierach, M. M.,
1290 Hill, C., Jahn, O., Landschützer, P., Lauderdale, J. M., Liu, J., Manizza, M., Naviaux, J. D., Rödenbeck, C., Schimel,
1291 D. S., Van der Stocken, T., and Zhang, H.: The ECCO-Darwin data-assimilative global ocean biogeochemistry
1292 model: Estimates of seasonal to multidecadal surface ocean $p\text{CO}_2$ and air-sea CO_2 flux, *J. Adv. Model. Earth Syst.*,
1293 12(10), <https://doi.org/10.1029/2019ms001888>, 2020.
- 1294 Carroll, D., Menemenlis, D., Dutkiewicz, S., Lauderdale, J. M., Adkins, J. F., Bowman, K. W., Brix, H., Fenty, I.,
1295 Gierach, M. M., Hill, C., Jahn, O., Landschützer, P., Manizza, M., Mazloff, M. R., Miller, C. E., Schimel, D. S.,
1296 Verdy, A., Whitt, D. B., and Zhang, H.: Attribution of space-time variability in global-ocean dissolved Inorganic
1297 Carbon, *Global Biogeochemical Cycles*, 36(3), <https://doi.org/10.1029/2021gb007162>, 2022.
- 1298 Carroll, D., Menemenlis, D., Zhang, H., Mazloff, M., McKinley, G., Fay, A., Dutkiewicz, S., Lauderdale, J., and
1299 Fenty, I.: Evaluation of the ECCO-Darwin Ocean Biogeochemistry State Estimate vs. In-situ Observations (ver 1.0),
1300 Zenodo, <https://doi.org/10.5281/zenodo.10627664>, 2024.
- 1301 Carter, B. R., Williams, N. L., Gray, A. R., and Feely, R. A.: Locally interpolated alkalinity regression for global
1302 alkalinity estimation, *Limnology and Oceanography: Methods*, 14(4), 268–277, <https://doi.org/10.1002/lom3.10087>,
1303 2016.
- 1304 Carter, B. R., Feely, R. A., Williams, N. L., Dickson, A. G., Fong, M. B., and Takeshita, Y.: Updated methods for
1305 global locally interpolated estimation of alkalinity, ph, and nitrate, *Limnology and Oceanography: Methods*, 16(2),
1306 119–131, <https://doi.org/10.1002/lom3.10232>, 2017.
- 1307 Carter, B. R., Bittig, H. C., Fassbender, A. J., Sharp, J. D., Takeshita, Y., Xu, Y.-Y., et al.: New and updated global
1308 empirical seawater property estimation routines, *Limnol. and Oceanogr.: Methods*, 19(12), 785–809,
1309 <https://doi.org/10.1002/lom3.10461>, 2021.
- 1310 Chau, T.-T.-T., Gehlen, M., and Chevallier, F.: A seamless ensemble-based reconstruction of surface ocean $p\text{CO}_2$
1311 and air–sea CO_2 fluxes over the global coastal and open oceans. *Biogeosciences*, 19(4), 1087–1109,
1312 <https://doi.org/10.5194/bg-19-1087-2022>, 2022.
- 1313 Chau, T.-T.-T., Gehlen, M., Metzl, N., and Chevallier, F.: CMEMS-LSCE: a global, 0.25° , monthly reconstruction
1314 of the surface ocean carbonate system, *Earth Syst. Sci. Data*, 16, 121–160, [https://doi.org/10.5194/essd-16-121-](https://doi.org/10.5194/essd-16-121-2024)
1315 2024, 2024a.
- 1316 Chau, T.-T.-T., Chevallier, F. and Gehlen, M.: Global analysis of surface ocean CO_2 fugacity and air-sea fluxes with
1317 low latency, *Geophys. Res. Lett.*, <https://doi.org/10.1029/2023GL106670>, 2024b.
- 1318 Chen, C.-T. A., Lui, H.-K., Hsieh, C.-H., Yanagi, T., Kosugi, N., Ishii, M., and Gong, G.-C.: Deep oceans may



- 1319 acidify faster than anticipated due to global warming, *Nat. Clim. Change.*, 7(12), 890–894,
 1320 <https://doi.org/10.1038/s41558-017-0003-y>, 2017.
- 1321 Clement, D., and Gruber, N.: The EMLR(C*) method to determine decadal changes in the global ocean storage of
 1322 Anthropogenic CO₂. *Global Biogeochemical Cycles*, 32(4), 654–679, <https://doi.org/10.1002/2017gb005819>, 2018.
- 1323 Cooley, S. R., and Doney, S. C.: Anticipating ocean acidification’s economic consequences for commercial
 1324 fisheries. *Environmental Research Letters*, 4(2), 024007, <https://doi.org/10.1088/1748-9326/4/2/024007>, 2009.
- 1325 Crisp, D., Dolman, H., Tanhua, T., McKinley, G. A., Hauck, J., Bastos, A., Sitch, S., Eggleston, S., and Aich, V.:
 1326 How well do we understand the land-ocean-atmosphere carbon cycle? *Reviews of Geophysics*, 60(2),
 1327 <https://doi.org/10.1029/2021RG000736>, 2022.
- 1328 Cyr, F., Gibb, O., Azetsu-Scott, K., Chassé, J., Galbraith, P., Maillet, G., Pepin, P., Punshon, S., and Starr, M.:
 1329 Ocean carbonate parameters on the Canadian Atlantic Continental Shelf, Federated Research Data Repository,
 1330 <https://doi.org/10.20383/102.0673>, 2022.
- 1331 Delaigue, L., Sulpis, O., Reichart, G. -J., and Humphreys, M. P.: The changing biological carbon pump of the South
 1332 Atlantic Ocean. *Global Biogeochemical Cycles*, 38(9), <https://doi.org/10.1029/2024gb008202>, 2024.
- 1333 DeVries, T., Holzer, M. and Primeau, F.: Recent increase in oceanic carbon uptake driven by weaker upper-ocean
 1334 overturning, *Nature*, 542, 215–218, <https://doi.org/10.1038/nature21068>, 2017.
- 1335 DeVries, T., Le Quéré, C., Andrews, O., Berthet, S., Hauck, J., Ilyina, T., Landschützer, P., Lenton, A., Lima, I. D.,
 1336 Nowicki, M., Schwinger, J., and Séférian, R.: Decadal trends in the ocean carbon sink, *Proceedings of the National*
 1337 *Academy of Sciences*, 116(24), 11646–11651, <https://doi.org/10.1073/pnas.1900371116>, 2019.
- 1338 DeVries, T.: The Ocean Carbon Cycle, *Annual Review of Environment and Resources*, 47(1), 317–341,
 1339 <https://doi.org/10.1146/annurev-environ-120920-111307>, 2022.
- 1340 DeVries, T., Yamamoto, K., Wanninkhof, R., Gruber, N., Hauck, J., Müller, J. D., Bopp, L., Carroll, D., Carter, B.,
 1341 Chau, T., Doney, S. C., Gehlen, M., Gloege, L., Gregor, L., Henson, S., Kim, J. H., Iida, Y., Ilyina, T.,
 1342 Landschützer, P., Le Quéré, C., Munro, D., Nissen, C., Patara, L., Pérez, F. F., Resplandy, L., Rodgers, K. B.,
 1343 Schwinger, J., Séférian, R., Sicardi, V., Terhaar, J., Triñanes, J., Tsujino, H., Watson, A., Yasunaka, S., Zeng, J.
 1344 (2023). Magnitude, trends, and variability of the Global Ocean Carbon Sink from 1985 to 2018. *Global*
 1345 *Biogeochemical Cycles*, 37(10). <https://doi.org/10.1029/2023gb007780>, 2023.
- 1346 Dickson, A. G.: Standard potential of the reaction: $\text{AgCl(s)} + 1/2 \text{H}_2\text{(g)} = \text{Ag(s)} + \text{HCl(aq)}$, and the standard acidity
 1347 constant of the ion HSO_4^- in synthetic seawater from 273.15 to 318.15K, *Journal of Chemical Thermodynamics*,
 1348 22(2), 113–127, [https://doi.org/10.1016/0021-9614\(90\)90074-Z](https://doi.org/10.1016/0021-9614(90)90074-Z), 1990.



- 1349 Doney, S. C., Busch, D. S., Cooley, S. R., and Kroeker, K. J.: The impacts of ocean acidification on marine
 1350 ecosystems and reliant human communities, *Annual Review of Environment and Resources*, 45(1), 83–112,
 1351 <https://doi.org/10.1146/annurev-environ-012320-083019>, 2020.
- 1352 Dore, J. E., Lukas, R., Sadler, D. W., and Karl, D. M.: Climate-driven changes to the atmospheric CO₂ sink in the
 1353 subtropical North Pacific Ocean, *Nature*, 424, 754–757, <https://doi.org/10.1038/nature01885>, 2003.
- 1354 Dore, J. E., Lukas, R., Sadler, D. W., Church, M. J., and Karl, D. M.: Physical and biogeochemical modulation of
 1355 ocean acidification in the central North Pacific, *Proc. Natl. Acad. Sci.*, 106, 12235–12240,
 1356 <https://doi.org/10.1073/pnas.0906044106>, 2009.
- 1357 Dore, J. E., Church, M. J., Karl, D. M., Sadler, D. W., and Letelier, R. M.: Paired windward and leeward
 1358 biogeochemical time series reveal consistent surface ocean CO₂ trends across the Hawaiian Ridge, *Geophys. Res.*
 1359 *Lett.*, 41, 6459–6467, <https://doi.org/10.1002/2014GL060725>, 2014.
- 1360 Dore, J.E., White, A.E., and Karl, D.M.: Hawaii Ocean Time-series (HOT) Carbon Dioxide (Version 1) [Data set].
 1361 Zenodo. <https://doi.org/10.5281/zenodo.15060931>, 2025.
- 1362 Dunne, John P., John, J. G., Shevliakova, E., Stouffer, R. J., Krasting, J. P., Malyshev, S. L., Milly, P. C., Sentman,
 1363 L. T., Adcroft, A. J., Cooke, W., Dunne, K. A., Griffies, S. M., Hallberg, R. W., Harrison, M. J., Levy, H.,
 1364 Wittenberg, A. T., Phillips, P. J., and Zadeh, N.: GFDL’s ESM2 global coupled climate–carbon earth system
 1365 models. part II: Carbon system formulation and baseline simulation characteristics, *Journal of Climate*, 26(7), 2247–
 1366 2267, <https://doi.org/10.1175/jcli-d-12-00150.1>, 2013.
- 1367 Dunne, J. P., Hewitt, H. T., Arblaster, J., Bonou, F., Boucher, O., Cavazos, T., Durack, P. J., Hassler, B., Jukes, M.,
 1368 Miyakawa, T., Mizielinski, M., Naik, V., Nicholls, Z., O’Rourke, E., Pincus, R., Sanderson, B. M., Simpson, I. R.,
 1369 and Taylor, K. E.: An Evolving Coupled Model Intercomparison Project Phase 7 (CMIP7) and Fast Track in
 1370 Support of Future Climate Assessment, <https://doi.org/10.5194/egusphere-2024-3874>, 2024.
- 1371 Durack, P. J., Taylor, K. E., Gleckler, P. J., Meehl, G. A., Lawrence, B. N., Covey, C., Stouffer, R. J., Levvasseur,
 1372 G., Ben-Nasser, A., Denvil, S., Stockhause, M., Gregory, J. M., Jukes, M., Ames, S. K., Antonio, F., Bader, D. C.,
 1373 Dunne, J. P., Ellis, D., Eyring, V., Fiore, S. L., Joussaume, S., Kershaw, P., Lamarque, J.-F., Lautenschlager, M.,
 1374 Lee, J., Mauzey, C. F., Mizielinski, M., Nassisi, P., Nuzzo, A., O’Rourke, E., Painter, J., Potter, G. L., Rodriguez,
 1375 S., Williams, D. N.: The Coupled Model Intercomparison Project (CMIP): Reviewing Project History, Evolution,
 1376 Infrastructure and Implementation, <https://doi.org/10.5194/egusphere-2024-3729>, 2025.
- 1377 Edmond, J. M.: High precision determination of titration alkalinity and total carbon dioxide content of sea water by
 1378 potentiometric titration, *Deep Sea Research and Oceanographic Abstracts*, 17(4), 737–750,
 1379 [https://doi.org/10.1016/0011-7471\(70\)90038-0](https://doi.org/10.1016/0011-7471(70)90038-0), 1970.



- 1380 Fassbender, A. J., Carter, B. R., Sharp, J. D., Huang, Y., Arroyo, M. C., and Frenzel, H.: Amplified subsurface
 1381 signals of ocean acidification, *Global Biogeochemical Cycles*, 37, e2023GB007843,
 1382 <https://doi.org/10.1029/2023GB007843>, 2023.
- 1383 Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G.,
 1384 Rödenbeck, C., Roobaert, A., and Zeng, J.: SeaFlux: harmonization of air–sea CO₂ fluxes from surface *p*CO₂ data
 1385 products using a standardized approach, *Earth Sys. Sci. Data*, 13(10), 4693–4710, [https://doi.org/10.5194/essd-13-](https://doi.org/10.5194/essd-13-4693-2021)
 1386 4693-2021, 2021.
- 1387 Fay, A. R., and McKinley, G. A.: Observed regional fluxes to constrain modelled estimates of the Ocean Carbon
 1388 Sink, *Geophysical Res. Lett.*, 48(20), <https://doi.org/10.1029/2021gl095325>, 2021.
- 1389 Fay, A. R., Munro, D. R., McKinley, G. A., Pierrot, D., Sutherland, S. C., Sweeney, C., and Wanninkhof, R.:
 1390 Updated climatological mean Δ /CO₂ and net sea–air CO₂ flux over the global open ocean regions, *Earth Syst. Sci.*
 1391 *Data*, 16, 2123–2139, <https://doi.org/10.5194/essd-16-2123-2024>, 2024.
- 1392 Feely, R. A., Jiang, L.-Q., Wanninkhof, R., Carter, B. R., Alin, S. A., Bednaršek, N., and Cosca, C. E.: Acidification
 1393 of the global surface ocean: What we have learned from observations, *Oceanography*, 36, 120–129,
 1394 <https://doi.org/10.5670/oceanog.2023.222>, 2023.
- 1395 Fennel, K., Mattern, J. P., Doney, S. C., Bopp, L., Moore, A. M., Wang, B., and Yu, L.: Ocean biogeochemical
 1396 modelling, *Nat. Rev. Methods Primers*, 2, 76, <https://doi.org/10.1038/s43586-022-00154-2>, 2022.
- 1397 Freeland, H.: A short history of ocean station papa and line P. *Progress in Oceanography*, 75(2), 120–125,
 1398 <https://doi.org/10.1016/j.pocean.2007.08.005>, 2007.
- 1399 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Bakker, D. C. E., Hauck, J., Landschützer, P., Le
 1400 Quéré, C., Luijkx, I. T., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais,
 1401 P., Jackson, R. B., Alin, S. R., Anthoni, P., Barbero, L., Bates, N. R., Becker, M., Bellouin, N., Decharme, B., Bopp,
 1402 L., Brasika, I. B. M., Cadule, P., Chamberlain, M. A., Chandra, N., Chau, T.-T.-T., Chevallier, F., Chini, L. P.,
 1403 Cronin, M., Dou, X., Enyo, K., Evans, W., Falk, S., Feely, R. A., Feng, L., Ford, D. J., Gasser, T., Ghattas, J.,
 1404 Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris, I., Hefner, M., Heinke, J., Houghton, R. A.,
 1405 Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A., Jarníková, T., Jersild, A., Jiang, F., Jin, Z., Joos, F., Kato,
 1406 E., Keeling, R. F., Kennedy, D., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Körtzinger, A., Lan, X.,
 1407 Lefèvre, N., Li, H., Liu, J., Liu, Z., Ma, L., Marland, G., Mayot, N., McGuire, P. C., McKinley, G. A., Meyer, G.,
 1408 Morgan, E. J., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K. M., Olsen, A., Omar, A. M., Ono, T., Paulsen,
 1409 M., Pierrot, D., Pockock, K., Poulter, B., Powis, C. M., Rehder, G., Resplandy, L., Robertson, E., Rödenbeck, C.,
 1410 Rosan, T. M., Schwinger, J., Séférian, R., Smallman, T. L., Smith, S. M., Sospedra-Alfonso, R., Sun, Q., Sutton, A.
 1411 J., Sweeney, C., Takao, S., Tans, P. P., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., van der Werf, G. R., van
 1412 Ooijen, E., Wanninkhof, R., Watanabe, M., Wimart-Rousseau, C., Yang, D., Yang, X., Yuan, W., Yue, X., Zaehle,



- 1413 S., Zeng, J., and Zheng, B.: Global Carbon Budget 2023, *Earth Syst. Sci. Data*, 15, 5301–5369,
- 1414 <https://doi.org/10.5194/essd-15-5301-2023>, 2023.
- 1415 Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Landschützer, P., Le Quéré, C., Li, H.,
- 1416 Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais,
- 1417 P., Jackson, R. B., Alin, S. R., Armeth, A., Arora, V., Bates, N. R., Becker, M., Bellouin, N., Berghoff, C. F., Bittig,
- 1418 H. C., Bopp, L., Cadule, P., Campbell, K., Chamberlain, M. A., Chandra, N., Chevallier, F., Chini, L. P., Colligan,
- 1419 T., Decayeux, J., Djeutchouang, L. M., Dou, X., Duran Rojas, C., Enyo, K., Evans, W., Fay, A. R., Feely, R. A.,
- 1420 Ford, D. J., Foster, A., Gasser, T., Gehlen, M., Gkritzalis, T., Grassi, G., Gregor, L., Gruber, N., Gürses, Ö., Harris,
- 1421 I., Hefner, M., Heinke, J., Hurtt, G. C., Iida, Y., Ilyina, T., Jacobson, A. R., Jain, A. K., Jarníková, T., Jersild, A.,
- 1422 Jiang, F., Jin, Z., Kato, E., Keeling, R. F., Klein Goldewijk, K., Knauer, J., Korsbakken, J. I., Lan, X., Lauvset, S.
- 1423 K., Lefèvre, N., Liu, Z., Liu, J., Ma, L., Maksyutov, S., Marland, G., Mayot, N., McGuire, P. C., Metzl, N.,
- 1424 Monacci, N. M., Morgan, E. J., Nakaoka, S.-I., Neill, C., Niwa, Y., Nützel, T., Olivier, L., Ono, T., Palmer, P. I.,
- 1425 Pierrot, D., Qin, Z., Resplandy, L., Roobaert, A., Rosan, T. M., Rödenbeck, C., Schwinger, J., Smallman, T. L.,
- 1426 Smith, S. M., Sospedra-Alfonso, R., Steinhoff, T., Sun, Q., Sutton, A. J., Séférian, R., Takao, S., Tatebe, H., Tian,
- 1427 H., Tilbrook, B., Torres, O., Tourigny, E., Tsujino, H., Tubiello, F., van der Werf, G., Wanninkhof, R., Wang, X.,
- 1428 Yang, D., Yang, X., Yu, Z., Yuan, W., Yue, X., Zaehle, S., Zeng, N., and Zeng, J.: Global Carbon Budget 2024,
- 1429 *Earth Syst. Sci. Data*, 17, 965–1039, <https://doi.org/10.5194/essd-17-965-2025>, 2025.
- 1430 Garcia, H. E., Locarnini, R. A., Boyer, T. P., Antonov, J. I., Baranova, O. K., Zweng, M. M., Reagan, J. R., and
- 1431 Johnson, D. R.: World Ocean Atlas 2013, Volume 3: Dissolved Oxygen, Apparent Oxygen Utilization, and Oxygen
- 1432 Saturation, edited by: Levitus, S. and Mishonov, A., Technical Ed., NOAA Atlas NES- DIS 75, 27 pp., 2014.
- 1433 Garcia, H. E., Weathers, K., Paver, C. R., Smolyar, I., Boyer, T. P., Locarnini, R. A., Zweng, M. M., Mishonov, A.
- 1434 V., Baranova, O. K., Seidov, D., and Reagan, J. R.: World Ocean Atlas 2018, Volume 4: Dissolved Inorganic
- 1435 Nutrients (phosphate, nitrate and nitrate+nitrite, silicate), A. Mishonov Technical Ed., NOAA Atlas NESDIS 84,
- 1436 35pp, 2018a.
- 1437 Garcia H. E., Weathers, K. W., Paver, C. R., Smolyar, I. V., Boyer, T. P., Locarnini, R. A., Zweng, M. M.,
- 1438 Mishonov, A. V., Baranova, O. K., and Reagan, J. R.: World Ocean Atlas 2018, Volume 3: Dissolved Oxygen,
- 1439 Apparent Oxygen Utilization, and Oxygen Saturation, Mishonov, A., Technical Editor, NOAA Atlas NESDIS 83,
- 1440 2018b.
- 1441 Garcia H. E., Boyer, T. P., Locarnini, R. A., Reagan, J. R., Mishonov, A. V., Baranova, O. K., Paver, C. R., Wang,
- 1442 Z., Bouchard, C. N., Cross, S. L., Seidov, D., and Dukhovskoy, D.: World Ocean Database 2023: User's Manual.
- 1443 A. V. Mishonov, Technical Ed., NOAA Atlas NESDIS 98, pp 129. <https://doi.org/10.25923/j8gq-eee82>, 2024.
- 1444 Gattuso, J.-P., and Hansson, L.: Ocean acidification, Oxford University Press, pp. 326, 2011.
- 1445 Gattuso, J.-P., Epitalon, J.-M., Lavigne, H., and Orr, J.: seacarb: Seawater Carbonate Chemistry, R Package



- 1446 (Version 3.2.16), <https://CRAN.R-project.org/package=seacarb>, 2021a.
- 1447 Gattuso, J.-P., Alliouane, S., Mousseau, L.: Seawater carbonate chemistry in the Bay of Villefranche, Point B
 1448 (France), January 2007 - June 2023 [dataset], PANGAEA, <https://doi.org/10.1594/PANGAEA.727120>, 2021b.
- 1449 Gattuso, J.-P., Alliouane, S., and Fischer, P.: High-frequency, year-round time series of the carbonate chemistry in a
 1450 high-Arctic fjord (Svalbard), *Earth Syst. Sci. Data*, 15, 2809–2825, <https://doi.org/10.5194/essd-15-2809-2023>,
 1451 2023a.
- 1452 Gattuso, J.-P., Alliouane, S., and Fischer, P.: High-frequency, year-round time series of the carbonate chemistry in a
 1453 high-Arctic fjord (Svalbard) [dataset], PANGAEA, <https://doi.org/10.1594/PANGAEA.957028>, 2023b.
- 1454 Gibb, O., Cyr, F., Azetsu-Scott, K., Chassé, J., Childs, D., Gabriel, C.-E., Galbraith, P. S., Maillet, G., Pepin, P.,
 1455 Punshon, S., and Starr, M.: Spatiotemporal variability in pH and carbonate parameters on the Canadian Atlantic
 1456 continental shelf between 2014 and 2022, *Earth Syst. Sci. Data*, 15, 4127–4162, [https://doi.org/10.5194/essd-15-](https://doi.org/10.5194/essd-15-4127-2023)
 1457 4127-2023, 2023.
- 1458 Gloege, L., Yan, M., Zheng, T., and McKinley, G. A.: Improved quantification of ocean carbon uptake by using
 1459 machine learning to merge global models and $p\text{CO}_2$ data, *Journal of Advances in Modeling Earth Systems*, 14(2),
 1460 <https://doi.org/10.1029/2021ms002620>, 2022.
- 1461 González-Dávila, M. and Santana-Casiano J. M.: Long-term trends of pH and inorganic carbon in the Eastern North
 1462 Atlantic: the ESTOC site, *Front. Mar. Sci.*, 10, 1236214, <https://doi.org/10.3389/fmars.2023.1236214>, 2023.
- 1463 Gregor, L., Lebehot, A. D., Kok, S., and Monteiro, P. M. S.: A comparative assessment of the uncertainties of
 1464 Global Surface Ocean CO_2 Estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) – have we hit
 1465 the wall? *Geoscientific Model Development*, 12(12), 5113–5136, <https://doi.org/10.5194/gmd-12-5113-2019>, 2019a.
- 1466 Gregor, L., Lebehot, A. D., Kok, S., Monteiro, P. M. S.: Global surface-ocean partial pressure of carbon dioxide
 1467 ($p\text{CO}_2$) estimates from a machine learning ensemble: CSIR-ML6 v2019a (NCEI Accession 0206205), NOAA
 1468 National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/z682-mn47>, 2019b.
- 1469 Gregor, L., and Gruber, N.: OceanSODA-ETHZ: A global gridded dataset of the surface ocean carbonate system for
 1470 seasonal to decadal studies of ocean acidification (v2023) (NCEI Accession 0220059), NOAA National Centers for
 1471 Environmental Information, Dataset, <https://doi.org/10.25921/m5wx-ja34>, 2020.
- 1472 Gregor, L.: luke-gregor/OceanSODA-ETHZ: code, Zenodo, <https://doi.org/10.5281/zenodo.4455354>, 2021.
- 1473 Gregor, L. and Fay, A.: SeaFlux: harmonised sea-air CO_2 fluxes from surface $p\text{CO}_2$ data products using a
 1474 standardised approach [Data set], Zenodo, <https://doi.org/10.5281/zenodo.5482547>, 2021.
- 1475 Gregor, L., and Gruber, N.: Oceansoda-ETHZ: A global gridded data set of the surface ocean carbonate system for



- 1476 seasonal to decadal studies of ocean acidification, *Earth System Science Data*, 13(2), 777–808,
 1477 <https://doi.org/10.5194/essd-13-777-2021>, 2021.
- 1478 Gregor, L., and Gruber, N.: OceanSODA-ETHZ: A global gridded dataset of the surface ocean carbonate system for
 1479 seasonal to decadal studies of ocean acidification (v2023) (NCEI accession 0220059) [Dataset], NOAA National
 1480 Centers for Environmental Information, <https://doi.org/10.25921/m5wx-ja34>, 2023.
- 1481 Gregor, L., Shutler, J., and Gruber, N.: High-resolution variability of the Ocean Carbon Sink, *Global*
 1482 *Biogeochemical Cycles*, 38(8), <https://doi.org/10.1029/2024gb008127>, 2024a.
- 1483 Gregor, L., Shutler, J., and Gruber, N.: OceanSODA-ETHZ-v2: Surface ocean sea-air CO₂ fluxes from 1982 to 2022
 1484 (8-day by 0.25° x 0.25°) (v2.2024r01), [Data set], Zenodo, <https://zenodo.org/records/11206366>, 2024b.
- 1485 Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr,
 1486 A., Lauvset, S. K., Lo Monaco, C., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T., and
 1487 Wanninkhof, R.: The oceanic sink for anthropogenic CO₂ from 1994 to 2007, *Science*, 363(6432), 1193–1199,
 1488 <https://doi.org/10.1126/science.aau5153>, 2019a.
- 1489 Gruber, N., Clement, D., Carter, B. R., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Key, R. M., Kozyr,
 1490 A., Lauvset, S. K., Lo Monaco, C., Mathis, J. T., Murata, A., Olsen, A., Perez, F. F., Sabine, C. L., Tanhua, T., and
 1491 Wanninkhof, R.: The oceanic sink for anthropogenic CO₂ from 1994 to 2007 - the data (NCEI Accession 0186034),
 1492 NOAA National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/wdn2-pt10>, 2019b.
- 1493 Gruber, N., Bakker, D. C. E., DeVries, T., Gregor, L., Hauck, J., Landschützer, P., McKinley, G. A., and Müller, J.
 1494 D.: Trends and variability in the ocean carbon sink, *Nature Reviews Earth and Environment*, 4(2), 119–134,
 1495 <https://doi.org/10.1038/s43017-022-00381-x>, 2023.
- 1496 Hauck, J., Mayot, N., Landschützer, P., & Jersild, A. (2025). Global Carbon Budget 2024, surface ocean fugacity of
 1497 CO₂ (*f*CO₂) and air-sea CO₂ flux of individual global ocean biogeochemical models and surface ocean *f*CO₂-based
 1498 data-products [Data set]. Zenodo. <https://doi.org/10.5281/zenodo.14639761>
- 1499 Holzer, M., and DeVries, T.: Source-labeled anthropogenic carbon reveals a large shift of preindustrial carbon from
 1500 the ocean to the atmosphere, *Global Biogeochemical Cycles*, 36(10), <https://doi.org/10.1029/2022gb007405>, 2022.
- 1501 Humphreys, M. P., Lewis, E. R., Sharp, J. D., and Pierrot, D.: PyCO₂SYs v1.8: Marine carbonate system
 1502 calculations in Python, *Geoscientific Model Development*, 15(1), 15–43, <https://doi.org/10.5194/gmd-15-15-2022>,
 1503 2022.
- 1504 Iida, Y., Takatani, Y., Kojima, A., and Ishii, M.: Global trends of ocean CO₂ sink and ocean acidification: An
 1505 observation-based reconstruction of surface ocean inorganic carbon variables, *Journal of Oceanography*, 77(2), 323–
 1506 358, <https://doi.org/10.1007/s10872-020-00571-5>, 2021.



- 1507 IPCC: Climate Change 2023: Synthesis Report, Contribution of Working Groups I, II and III to the Sixth
- 1508 Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero
- 1509 (eds.)], IPCC, Geneva, Switzerland, pp. 35-115, <https://doi.org/10.59327/IPCC/AR6-9789291691647>, 2023.
- 1510 Ishii, M., Kosugi, N., Sasano, D., Saito, S., Midorikawa, T., & Inoue, H. Y.: Ocean acidification off the south coast
- 1511 of Japan: A result from time series observations of CO₂ Parameters from 1994 to 2008. *Journal of Geophysical*
- 1512 *Research*, 116(C6), <https://doi.org/10.1029/2010jc006831>, 2011a.
- 1513 Ishii, M., Suzuki, T., and Key, R.: Pacific Ocean Interior Carbon Data Synthesis, PACIFICA, in Progress, PICES
- 1514 Press, 19(1), 20-23, https://www.pices.int/publications/pices_press/volume19/v19_n1/pp_20-23_PACIFICA_f.pdf,
- 1515 2011b.
- 1516 Jersild, A., Landschützer, P., Gruber, N., Bakker, D. C. E.: An observation-based global monthly gridded sea surface
- 1517 *p*CO₂ and air-sea CO₂ flux product from 1982 onward and its monthly climatology (NCEI Accession 0160558),
- 1518 NOAA National Centers for Environmental Information, Dataset, <https://doi.org/10.7289/v5z899n6>, 2017.
- 1519 Jiang, L.-Q., Feely, R. A., Carter, B. R., Greeley, D. J., Gledhill, D. K., and Arzayus, K. M.: Climatological
- 1520 distribution of aragonite saturation state in the global oceans, *Global Biogeochemical Cycles*, 29(10), 1656–1673,
- 1521 <https://doi.org/10.1002/2015GB005198>, 2015a.
- 1522 Jiang, L.-Q., Feely, R. A.: Aragonite saturation state gridded to 1x1 degree latitude and longitude at depth levels of 0,
- 1523 50, 100, 200, 500, 1000, 2000, 3000, and 4000 meters in the global oceans (NCEI Accession 0139360), NOAA
- 1524 National Centers for Environmental Information, Dataset, <https://doi.org/10.7289/v5q81b4p>, 2015b.
- 1525 Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S., and Olsen, A.: Surface ocean pH and buffer capacity: Past,
- 1526 present and future, *Scientific Reports*, 9(1), 18624, <https://doi.org/10.1038/s41598-019-55039-4>, 2019a.
- 1527 Jiang, L.-Q., Carter, B. R., Feely, R. A., Lauvset, S., and Olsen, A.: Global surface ocean pH, acidity, and Revelle
- 1528 Factor on a 1x1 degree global grid from 1770 to 2100 (NCEI Accession 0206289), NOAA National Centers for
- 1529 Environmental Information, Dataset, <https://doi.org/10.25921/kgqr-9h49>, 2019b.
- 1530 Jiang, L.-Q., Feely, R. A., Wanninkhof, R., Greeley, D., Barbero, L., Alin, S., Carter, B. R., Pierrot, D.,
- 1531 Featherstone, C., Hooper, J., Melrose, C., Monacci, N., Sharp, J. D., Shellito, S., Xu, Y.-Y., Kozyr, A., Byrne, R. H.,
- 1532 Cai, W.-J., Cross, J., Johnson, G. C., Hales, B., Langdon, C., Mathis, J., Salisbury, J., and Townsend, D. W.: Coastal
- 1533 Ocean Data Analysis Product in North America (CODAP-NA, Version 2021) (NCEI Accession 0219960), NOAA
- 1534 National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/531n-c230>, 2020.
- 1535 Jiang, L.-Q., Feely, R. A., Wanninkhof, R., Greeley, D., Barbero, L., Alin, S., Carter, B. R., Pierrot, D.,
- 1536 Featherstone, C., Hooper, J., Melrose, C., Monacci, N., Sharp, J. D., Shellito, S., Xu, Y.-Y., Kozyr, A., Byrne, R. H.,
- 1537 Cai, W.-J., Cross, J., Johnson, G. C., Hales, B., Langdon, C., Mathis, J., Salisbury, J., and Townsend, D. W.: Coastal



- 1538 Ocean Data Analysis Product in North America (CODAP-NA) – an internally consistent data product for discrete
- 1539 inorganic carbon, oxygen, and nutrients on the North American ocean margins, *Earth Syst. Sci. Data*, 13, 2777–
- 1540 2799, <https://doi.org/10.5194/essd-13-2777-2021>, 2021.
- 1541 Jiang, L.-Q., Pierrot, D., Wanninkhof, R., Feely, R. A., Tilbrook, B., Alin, S., Barbero, L., Byrne, R. H., Carter, B.
- 1542 R., Dickson, A. G., Gattuso, J.-P., Greeley, D., Hoppema, M., Humphreys, M. P., Karstensen, J., Lange, N.,
- 1543 Lauvset, S. K., Lewis, E. R., Olsen, A., Perez, F. F., Sabine, C., Sharp, J. D., Tanhua, T., Trull, T. W., Velo, A.,
- 1544 Allegra, A. J., Barker, P., Burger, E., Cai, W.-J., Chen, C.-T. A., Cross, J., Garcia, H., Hernandez-Ayon, J. M., Hu,
- 1545 X., Kozyr, A., Langdon, C., Lee, K., Salisbury, J., Wang, Z. A., and Xue, L.: Best practice data standards for
- 1546 discrete chemical oceanographic observations. *Frontiers in Marine Science*, 8,
- 1547 <https://doi.org/10.3389/fmars.2021.705638>, 2022a.
- 1548 Jiang, L.-Q., Boyer, T. P., Paver, C. R., Yoo, H., Reagan, J. R., Alin, S. R., Barbero, L., Carter, B. R., Feely, R. A.,
- 1549 and Wanninkhof, R.: Climatological distribution of ocean acidification indicators from surface to 500 meters water
- 1550 depth on the North American ocean margins from 2003-12-06 to 2018-11-22 (NCEI Accession 0270962), NOAA
- 1551 National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/g8pb-zy76>, 2022b.
- 1552 Jiang, L., Dunne, J., Carter, B. R., Tjiputra, J. F., Terhaar, J., Sharp, J. D., Olsen, A., Alin, S., Bakker, D. C., Feely,
- 1553 R. A., Gattuso, J., Hogan, P., Ilyina, T., Lange, N., Lauvset, S. K., Lewis, E. R., Lovato, T., Palmieri, J., Santana-
- 1554 Falcón, Y., Schwinger, J., Séférian, R., Strand, G., Swart, N., Tanhua, T., Tsujino, H., Wanninkhof, R., Watanabe,
- 1555 M., Yamamoto, A., and Ziehn, T.: Global surface ocean acidification indicators from 1750 to 2100 (NCEI
- 1556 Accession 0259391), NOAA National Centers for Environmental Information, Dataset,
- 1557 <https://doi.org/10.25921/9ker-bc48>, 2022c.
- 1558 Jiang, L., Dunne, J., Carter, B. R., Tjiputra, J. F., Terhaar, J., Sharp, J. D., Olsen, A., Alin, S., Bakker, D. C., Feely,
- 1559 R. A., Gattuso, J., Hogan, P., Ilyina, T., Lange, N., Lauvset, S. K., Lewis, E. R., Lovato, T., Palmieri, J., Santana-
- 1560 Falcón, Y., Schwinger, J., Séférian, R., Strand, G., Swart, N., Tanhua, T., Tsujino, H., Wanninkhof, R., Watanabe,
- 1561 M., Yamamoto, A., and Ziehn, T.: Global surface ocean acidification indicators from 1750 to 2100, *Journal of*
- 1562 *Advances in Modeling Earth Systems*, 15(3), <https://doi.org/10.1029/2022ms003563>, 2023.
- 1563 Jiang, L.-Q., Boyer, T. P., Paver, C. R., Yoo, H., Reagan, J. R., Alin, S. R., Barbero, L., Carter, B. R., Feely, R. A.,
- 1564 and Wanninkhof, R.: Climatological distribution of ocean acidification variables along the North American ocean
- 1565 margins, *Earth System Science Data*, 16(7), 3383–3390, <https://doi.org/10.5194/essd-16-3383-2024>, 2024.
- 1566 Johnson, G. C., Robbins, P. E., and Hufford, G. E. (2001). Systematic adjustments of hydrographic sections for
- 1567 internal consistency. *Journal of Atmospheric and Oceanic Technology*, 18(7), 1234–1244, 2001.
- 1568 Kapsenberg, L., Alliouane, S., Gazeau, F., Mousseau, L., and Gattuso, J.-P.: Coastal ocean acidification and
- 1569 increasing total alkalinity in the northwestern Mediterranean Sea, *Ocean Sci.*, 13, 411–426,
- 1570 <https://doi.org/10.5194/os-13-411-2017>, 2017.



- 1571 Karl, D., Dore, J., Lukas, R., Michaels, A., Bates, N., and Knap, A.: Building the long-term picture: The U.S.
- 1572 JGOFS Time-series programs. *Oceanography*, 14(4), 6–17, <https://doi.org/10.5670/oceanog.2001.02>, 2001.
- 1573 Karl, D. M., and Lukas, R.: The Hawaii Ocean Time-series (HOT) program: Background, rationale and field
- 1574 implementation, *Deep Sea Research Part II*, 43(2–3), 129–156, [https://doi.org/10.1016/0967-0645\(96\)00005-7](https://doi.org/10.1016/0967-0645(96)00005-7),
- 1575 1996.
- 1576 Kennedy, E. G., Zulian, M., Hamilton, S. L., Hill, T. M., Delgado, M., Fish, C. R., Gaylord, B., Kroeker, K. J.,
- 1577 Palmer, H. M., Ricart, A. M., Sanford, E., Spalding, A. K., Ward, M., Carrasco, G., Elliott, M., Grisby, G. V.,
- 1578 Harris, E., Jahncke, J., Rocheleau, C. N., Westerink, S., and Wilmot, M. I.: Multistressor Observations of Coastal
- 1579 Hypoxia and Acidification (MOCHA) Synthesis (NCEI Accession 0277984), NOAA National Centers for
- 1580 Environmental Information, Dataset, <https://doi.org/10.25921/2vve-fh39>, 2023.
- 1581 Kennedy, E. G., Zulian, M., Hamilton, S. L., Hill, T. M., Delgado, M., Fish, C. R., Gaylord, B., Kroeker, K. J.,
- 1582 Palmer, H. M., Ricart, A. M., Sanford, E., Spalding, A. K., Ward, M., Carrasco, G., Elliott, M., Grisby, G. V.,
- 1583 Harris, E., Jahncke, J., Rocheleau, C. N., Westerink, S., and Wilmot, M. I.: A high-resolution synthesis dataset for
- 1584 multistressor analyses along the US West Coast, *Earth Syst. Sci. Data*, 16, 219–243, <https://doi.org/10.5194/essd->
- 1585 16-219-2024, 2024.
- 1586 Keppler, L., Landschützer, P., Gruber, N., Lauvset, S. K., and Stemmler, I.: Seasonal carbon dynamics in the near-
- 1587 global ocean, *Global Biogeochemical Cycles*, 34(12), e2020GB006571, <https://doi.org/10.1029/2020GB006571>,
- 1588 2020.
- 1589 Keppler, L., Landschützer, P., Lauvset, S. K., and Gruber, N.: Recent trends and variability in the oceanic storage of
- 1590 dissolved inorganic carbon, *Global Biogeochemical Cycles*, 37(5), <https://doi.org/10.1029/2022gb007677>, 2023.
- 1591 Key, R. M., Kozyr, A., Sabine, C. L., Lee, K., Wanninkhof, R., Bullister, J. L., Feely, R. A., Millero, F. J., Mordy,
- 1592 C., and Peng, T. -H.: A global ocean carbon climatology: Results from Global Data Analysis Project (GLODAP),
- 1593 *Global Biogeochemical Cycles*, 18(4), <https://doi.org/10.1029/2004gb002247>, 2004.
- 1594 Key, R. M., Tanhua, T., Olsen, A., Hoppema, M., Jutterström, S., Schirnack, C., van Heuven, S., Kozyr, A., Lin, X.,
- 1595 Velo, A., Wallace, D. W. R., and Mintrop, L.: The CARINA data synthesis project: introduction and overview,
- 1596 *Earth Syst. Sci. Data*, 2, 105–121, <https://doi.org/10.5194/essd-2-105-2010>, 2010.
- 1597 Key, R. M., Olsen, A., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T.,
- 1598 Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishi, M., Perez, F. F., and Suzuki, T.: Global Ocean Data
- 1599 Analysis Project, Version 2 (GLODAPv2), ORNL/CDIAC-162, ND-P093, Carbon Dioxide Information Analysis
- 1600 Center, Oak Ridge National Laboratory, U. S. Department of Energy, Oak Ridge, Tennessee,
- 1601 https://doi.org/10.3334/CDIAC/OTG.NDP093_GLODAPv2, 2015.



- 1602 Khesghi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, *Energy*, 20(9), 915–922,
 1603 [https://doi.org/10.1016/0360-5442\(95\)00035-f](https://doi.org/10.1016/0360-5442(95)00035-f), 1995.
- 1604 Knor, L. A. C. M., Sabine, C. L., Sutton, A. J., White, A. E., Potemra, J., and Weller, R. A.: Quantifying net
 1605 community production and calcification at Station ALOHA near Hawai‘i: Insights and limitations from a dual tracer
 1606 carbon budget approach, *Global Biogeochemical Cycles*, 37, e2022GB007672,
 1607 <https://doi.org/10.1029/2022GB007672>, 2023.
- 1608 Kwiatkowski, L., Torres, O., Bopp, L., Aumont, O., Chamberlain, M., Christian, J. R., Dunne, J. P., Gehlen, M.,
 1609 Ilyina, T., John, J. G., Lenton, A., Li, H., Lovenduski, N. S., Orr, J. C., Palmieri, J., Santana-Falcón, Y., Schwinger,
 1610 J., Séférian, R., Stock, C. A., Tagliabue, A., Takano, Y., Tjiputra, J., Toyama, K., Tsujino, H., Watanabe, M.,
 1611 Yamamoto, A., Yool, A., and Ziehn, T.: Twenty-first century ocean warming, acidification, deoxygenation, and
 1612 upper-ocean nutrient and primary production decline from CMIP6 model projections, *Biogeosciences*, 17, 3439–
 1613 3470, <https://doi.org/10.5194/bg-17-3439-2020>, 2020.
- 1614 Landschützer, P., Gruber, N., Bakker, D. C. E., Schuster, U., Nakaoka, S., Payne, M. R., Sasse, T., and Zeng, J.: A
 1615 neural network-based estimate of the seasonal to inter-annual variability of the Atlantic Ocean carbon sink,
 1616 *Biogeosciences*, 10, 7793–7815, <https://doi.org/10.5194/bg-10-7793-2013>, 2013.
- 1617 Landschützer, P., Gruber, N., Bakker, D. C. E., and Schuster, U.: Recent variability of the Global Ocean Carbon
 1618 Sink, *Global Biogeochemical Cycles*, 28(9), 927–949, <https://doi.org/10.1002/2014gb004853>, 2014.
- 1619 Landschützer, P., Gruber, N., and Bakker, D. C. E.: Decadal variations and trends of the Global Ocean Carbon Sink,
 1620 *Global Biogeochemical Cycles*, 30(10), 1396–1417. <https://doi.org/10.1002/2015gb005359>, 2016.
- 1621 Landschützer, P., Laruelle, G. G., Roobaert, A., and Regnier, P.: A uniform $p\text{CO}_2$ climatology combining open and
 1622 coastal oceans, *Earth Syst. Sci. Data*, 12, 2537–2553, <https://doi.org/10.5194/essd-12-2537-2020>, 2020a.
- 1623 Landschützer, P., Laruelle, G. G., Roobaert, A., and Regnier, P.: A combined global ocean $p\text{CO}_2$ climatology
 1624 combining open ocean and coastal areas (NCEI Accession 0209633), NOAA National Centers for Environmental
 1625 Information, Dataset, <https://doi.org/10.25921/qb25-f418>, 2020b.
- 1626 Lange, N., Tanhua, T., Pfeil, B., Bange, H. W., Lauvset, S. K., Grégoire, M., Bakker, D. C., Jones, S. D., Fiedler, B.,
 1627 O’Brien, K. M., and Körtzinger, A.: A status assessment of selected data synthesis products for ocean
 1628 biogeochemistry, *Frontiers in Marine Science*, 10, <https://doi.org/10.3389/fmars.2023.1078908>, 2023.
- 1629 Lange, N., Fiedler, B., Álvarez, M., Benoit-Cattin, A., Benway, H., Buttigieg, P. L., Coppola, L., Currie, K., Flecha,
 1630 S., Gerlach, D. S., Honda, M., Huertas, I. E., Lauvset, S. K., Muller-Karger, F., Körtzinger, A., O’Brien, K. M.,
 1631 Ólafsdóttir, S. R., Pacheco, F. C., Rueda-Roa, D., Skjelvan, I., Wakita, M., White, A., and Tanhua, T.: Synthesis
 1632 product for ocean time series (spots) – a ship-based biogeochemical pilot, *Earth System Science Data*, 16(4), 1901–



- 1633 1931, <https://doi.org/10.5194/essd-16-1901-2024>, 2024a.
- 1634 Lange, N., Fiedler, B., Álvarez, M., Benoit-Cattin, A., Benway, H., Buttigieg, P. L., Coppola, L., Currie, K. I.,
 1635 Flecha, S., Gerlach, D. S., Honda, M. C., Huertas, E. I., Kinkade, D., Muller-Karger, F., Lauvset, S. K., Körtzinger,
 1636 A., O'Brien, K. M., Ólafsdóttir, S., Pacheco, F. C., Rueda-Roa, D., Skjelvan, I., Wakita, M., White, A. E., Tanhua,
 1637 T.: Synthesis Product for Ocean Time Series (SPOTS), Biological and Chemical Oceanography Data Management
 1638 Office (BCO-DMO), (Version 2) Version Date 2024-02-22, <https://doi.org/10.26008/1912/bco-dmo.896862.2>,
 1639 2024b.
- 1640 Laruelle, G. G., Landschützer, P., Gruber, N., Tison, J.-L., Delille, B., and Regnier, P.: Global high-resolution
 1641 monthly $p\text{CO}_2$ climatology for the coastal ocean derived from neural network interpolation, *Biogeosciences*, 14,
 1642 4545–4561, <https://doi.org/10.5194/bg-14-4545-2017>, 2017.
- 1643 Lauvset, S. K., and Tanhua, T.: A toolbox for secondary quality control on ocean chemistry and Hydrographic Data,
 1644 *Limnology and Oceanography: Methods*, 13(11), 601–608, <https://doi.org/10.1002/lom3.10050>, 2015.
- 1645 Lauvset, S. K., Key, R. M., Olsen, A., van Heuven, S., Velo, A., Lin, X., Schirnack, C., Kozyr, A., Tanhua, T.,
 1646 Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Perez, F. F., Suzuki, T., and Watelet, S.: A new
 1647 global interior ocean mapped climatology: The $1^\circ \times 1^\circ$ GLODAP version 2, *Earth System Science Data*, 8(2), 325–
 1648 340, <https://doi.org/10.5194/essd-8-325-2016>, 2016.
- 1649 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Becker, S., Brown, P. J.,
 1650 Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jutterström,
 1651 S., Jones, S. D., Karlsen, M. K., Lo Monaco, C., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B., Schirnack, C.,
 1652 Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., Woosley, R. J., and Key, R. M.: An updated
 1653 version of the global interior ocean biogeochemical data product, GLODAPv2.2021, *Earth Syst. Sci. Data*, 13,
 1654 5565–5589, <https://doi.org/10.5194/essd-13-5565-2021>, 2021.
- 1655 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Alin, S., Álvarez, M., Azetsu-Scott, K.,
 1656 Barbero, L., Becker, S., Brown, P. J., Carter, B. R., da Cunha, L. C., Feely, R. A., Hoppema, M., Humphreys, M. P.,
 1657 Ishii, M., Jeansson, E., Jiang, L.-Q., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F., Pfeil, B.,
 1658 Schirnack, C., Steinfeldt, R., Suzuki, T., Tilbrook, B., Ulfssbo, A., Velo, A., Woosley, R. J., and Key, R. M.:
 1659 GLODAPv2.2022: the latest version of the global interior ocean biogeochemical data product, *Earth Syst. Sci. Data*,
 1660 14, 5543–5572, <https://doi.org/10.5194/essd-14-5543-2022>, 2022.
- 1661 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Becker, S.,
 1662 Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson,
 1663 E., Jones, S. D., Lo Monaco, C., Murata, A., Müller, J. D., Pérez, F. F., Schirnack, C., Steinfeldt, R., Suzuki, T.,
 1664 Tilbrook, B., Ulfssbo, A., Velo, A., Woosley, R. J., Key, R. M.: Global Ocean Data Analysis Project version 2.2023
 1665 (GLODAPv2.2023) (NCEI Accession 0283442), NOAA National Centers for Environmental Information, Dataset,



- 1666 <https://doi.org/10.25921/zyrq-ht66>, 2023a.
- 1667 Lauvset, S. K., Key, R. M., Olsen, A., van Heuven, S. M. A. C., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua,
1668 T., Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., Suzuki, T., Watelet, S.: A new
1669 global interior ocean mapped climatology: the $1^\circ \times 1^\circ$ GLODAP version 2 from 1972-01-01 to 2013-12-31 (NCEI
1670 Accession 0286118), NOAA National Centers for Environmental Information, Dataset,
1671 https://doi.org/10.3334/cdiac/otg.ndp093_glodapv2, 2023b.
- 1672 Lauvset, S. K., Lange, N., Tanhua, T., Bittig, H. C., Olsen, A., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Brown, P.
1673 J., Carter, B. R., Cotrim da Cunha, L., Hoppema, M., Humphreys, M. P., Ishii, M., Jeansson, E., Murata, A., Müller,
1674 J. D., Pérez, F. F., Schirnick, C., Steinfeldt, R., Suzuki, T., Ulfso, A., Velo, A., Woosley, R. J., and Key, R. M.:
1675 The annual update GLODAPv2.2023: the global interior ocean biogeochemical data product, *Earth Syst. Sci. Data*,
1676 16, 2047–2072, <https://doi.org/10.5194/essd-16-2047-2024>, 2024.
- 1677 Lee, K., Tong, L. T., Millero, F. J., Sabine, C. L., Dickson, A. G., Goyet, C., Park, G., Wanninkhof, R., Feely, R. A.,
1678 and Key, R. M.: Global relationships of total alkalinity with salinity and temperature in surface waters of the world's
1679 oceans, *Geophysical Research Letters*, 33(19), <https://doi.org/10.1029/2006gl027207>, 2006.
- 1680 Lee, K., Kim, T.-W., Byrne, R. H., Millero, F. J., Feely, R. A., and Liu, Y.-M.: The universal ratio of boron to
1681 chlorinity for the North Pacific and North Atlantic oceans, *Geochimica et Cosmochimica Acta*, 74(6), 1801–1811,
1682 <https://doi.org/10.1016/j.gca.2009.12.027>, 2010.
- 1683 Lewis, E., and Wallace, D. W. R.: Program Developed for CO₂ System Calculations, ORNL/CDIAC-105, Carbon
1684 Dioxide Information Analysis Center, Oak Ridge National Laboratory, Oak Ridge, TN, 1998.
- 1685 Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., Zweng, M. M.,
1686 Paver, C. R., Reagan, J. R., Johnson, D. R., Hamilton, M., and Seidov, D.: World Ocean Atlas 2013, Volume 1:
1687 Temperature, edited by: Levitus, S. and Mishonov, A., NOAA Atlas NESDIS 73, 40 pp., 2013.
- 1688 Locarnini, R. A., Mishonov, A. V., Baranova, O. K., Boyer, T. P., Zweng, M. M., Garcia, H. E., Reagan, J. R.,
1689 Seidov, D., Weathers, K. W., Paver, C. R., Smolyar, I. V.: World Ocean Atlas 2018, Volume 1: Temperature,
1690 Mishonov, A. [Technical Editor], NOAA Atlas NESDIS 81, 2019.
- 1691 Lueker, T. J., Dickson, A. G., and Keeling, C. D.: Ocean $p\text{CO}_2$ calculated from dissolved inorganic carbon,
1692 alkalinity, and equations for K₁ and K₂: Validation based on laboratory measurements of CO₂ in gas and seawater at
1693 equilibrium, *Marine Chemistry*, 70(1–3), 105–119, [https://doi.org/10.1016/s0304-4203\(00\)00022-0](https://doi.org/10.1016/s0304-4203(00)00022-0), 2000.
- 1694 Ma, D., Gregor, L., and Gruber, N. (2023). Four decades of trends and drivers of global surface ocean acidification,
1695 *Global Biogeochemical Cycles*, 37(7), <https://doi.org/10.1029/2023gb007765>, 2023.
- 1696 Manzello, D. P., Enochs, I. C., Hendee, J. C.: National Coral Reef Monitoring Program: Carbonate chemistry data



- 1697 collected in the Atlantic Ocean, NOAA National Centers for Environmental Information, Dataset,
 1698 <https://doi.org/10.25921/vfz0-dg77>, 2018.
- 1699 Mercier, H., Desbruyères, D., Lherminier, P., Velo, A., Carracedo, L., Fontela, M., & Pérez, F. F.: New insights into
 1700 the eastern subpolar North Atlantic Meridional overturning circulation from OVIDE. *Ocean Science*, 20(3), 779–
 1701 797, <https://doi.org/10.5194/os-20-779-2024>, 2024.
- 1702 Metzl, N., Tilbrook, B., Bakker, D., le Quéré, C., Doney, S., Feely, R., Hood, M., and Dargaville, R.: Surface Ocean
 1703 CO₂ variability and Vulnerability Workshop, Paris, France, 11–14 April 2007. *Eos, Transactions American*
 1704 *Geophysical Union*, 88(28), 287–287, <https://doi.org/10.1029/2007eo280005>, 2007.
- 1705 Metzl, N., Fin, J., Lo Monaco, C., Mignon, C., Alliouane, S., Antoine, D., Bourdin, G., Boutin, J., Bozec, Y.,
 1706 Conan, P., Coppola, L., Diaz, F., Douville, E., Durrieu de Madron, X., Gattuso, J.-P., Gazeau, F., Golbol, M.,
 1707 Lansard, B., Lefèvre, D., Lefèvre, N., Lombard, F., Louanchi, F., Merlivat, L., Olivier, L., Petrenko, A., Petton, S.,
 1708 Pujo-Pay, M., Rabouille, C., Reverdin, G., Ridame, C., Tribollet, A., Vellucci, V., Wagener, T., and Wimart-
 1709 Rousseau, C.: A synthesis of ocean total alkalinity and dissolved inorganic carbon measurements from 1993 to
 1710 2022: the SNAPO-CO₂-v1 dataset, *Earth Syst. Sci. Data*, 16, 89–120, <https://doi.org/10.5194/essd-16-89-2024>,
 1711 2024.
- 1712 Metzl, Nicolas; Fin, Jonathan; Lo Monaco, Claire; Mignon, Claude; Alliouane, Samir; Antoine, David; Bourdin,
 1713 Guillaume; Boutin, Jacqueline; Bozec, Yann; Conan, Pascal; Coppola, Laurent; Diaz, Frédéric; Douville, Eric;
 1714 Durrieu de Madron, Xavier; Gattuso, Jean-Pierre; Gazeau, Frédéric; Golbol, Melek; Lansard, Bruno; Lefèvre,
 1715 Dominique; Lefèvre, Nathalie; Lombard, Fabien; Louanchi, Férial; Merlivat, Liliane; Olivier, Léa; Petrenko, Anne;
 1716 Petton, Sébastien; Pujo-Pay, Mireille; Rabouille, Christophe; Reverdin, Gilles; Ridame, Céline; Tribollet, Aline;
 1717 Vellucci, Vincenzo; Wagener, Thibaut; Wimart-Rousseau, Cathy (2023). A synthesis of ocean total alkalinity and
 1718 dissolved inorganic carbon measurements in the Global Ocean and Mediterranean Sea from 1993 to 2022: the
 1719 SNAPO-CO₂-v1 dataset (NCEI Accession 0285681). [indicate subset used]. NOAA National Centers for
 1720 Environmental Information. Unpublished Dataset. <https://www.ncei.noaa.gov/archive/accession/0285681>, 2023.
- 1721 Mishonov, A. V., Boyer, T. P., Baranova, O. K., Bouchard, C. N., Cross, S. L., Garcia H. E., Locarnini, R. A., Paver,
 1722 C. R., Wang, Z., Seidov, D., Grodsky, A. I., Beauchamp, J. G.: *World Ocean Database 2023*, C. Bouchard, Technical
 1723 Ed., NOAA Atlas NESDIS 97, 206 pp., <https://doi.org/10.25923/z885-h264>, 2024.
- 1724 Monacci, N. M., Cross, J. N., Danielson, S. L., Evans, W., Hopcroft, R. R., Mathis, J. T., Mordy, C. W., Naber, D.
 1725 D., Shake, K. L., Trahanovsky, K., Wang, H., Weingartner, T. J., Whittedge, T. E.: Marine carbonate system
 1726 discrete profile data from the Gulf of Alaska (GAK) Seward Line cruises between 2008 and 2017 (NCEI Accession
 1727 0277034). NOAA National Centers for Environmental Information. Dataset. <https://doi.org/10.25921/x9sg-9b08>,
 1728 2023.



- 1729 Monacci, N. M., Cross, J. N., Evans, W., Mathis, J. T., and Wang, H.: A decade of marine inorganic carbon
 1730 chemistry observations in the northern Gulf of Alaska – insights into an environment in transition, *Earth Syst. Sci.*
 1731 *Data*, 16, 647–665, <https://doi.org/10.5194/essd-16-647-2024>, 2024.
- 1732 Müller, J. D., Gruber, N., Carter, B., Feely, R., Ishii, M., Lange, N., Lauvset, S. K., Murata, A., Olsen, A., Pérez, F.
 1733 F., Sabine, C., Tanhua, T., Wanninkhof, R., and Zhu, D.: Decadal trends in the oceanic storage of anthropogenic
 1734 carbon from 1994 to 2014, *AGU Advances*, 4(4), <https://doi.org/10.1029/2023av000875>, 2023.
- 1735 Müller, J. D.: RECCAP2-ocean data collection, <https://doi.org/10.5281/zenodo.7990823>, 2023.
- 1736 Müller, J. D., and Gruber, N.: Progression of ocean interior acidification over the industrial era, *Science Advances*,
 1737 10(48), <https://doi.org/10.1126/sciadv.ado3103>, 2024.
- 1738 Olsen, A., Key, R. M., van Heuven, S., Lauvset, S. K., Velo, A., Lin, X., Schirnick, C., Kozyr, A., Tanhua, T.,
 1739 Hoppema, M., Jutterström, S., Steinfeldt, R., Jeansson, E., Ishii, M., Pérez, F. F., and Suzuki, T.: The Global Ocean
 1740 Data Analysis Project version 2 (GLODAPv2) – an internally consistent data product for the world ocean, *Earth*
 1741 *Syst. Sci. Data*, 8, 297–323, <https://doi.org/10.5194/essd-8-297-2016>, 2016.
- 1742 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Álvarez, M., Becker, S., Bittig, H. C., Carter, B. R., Cotrim da Cunha,
 1743 L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson, E., Jones, S. D., Jutterström, S., Karlsen, M. K.,
 1744 Kozyr, A., Lauvset, S. K., Lo Monaco, C., Murata, A., Pérez, F. F., Pfeil, B., Schirnick, C., Steinfeldt, R., Suzuki,
 1745 T., Telszewski, M., Tilbrook, B., Velo, A., and Wanninkhof, R.: GLODAPv2.2019 – an update of GLODAPv2,
 1746 *Earth Syst. Sci. Data*, 11, 1437–1461, <https://doi.org/10.5194/essd-11-1437-2019>, 2019.
- 1747 Olsen, A., Lange, N., Key, R. M., Tanhua, T., Bittig, H. C., Kozyr, A., Álvarez, M., Azetsu-Scott, K., Becker, S.,
 1748 Brown, P. J., Carter, B. R., Cotrim da Cunha, L., Feely, R. A., van Heuven, S., Hoppema, M., Ishii, M., Jeansson,
 1749 E., Jutterström, S., Landa, C. S., Lauvset, S. K., Michaelis, P., Murata, A., Pérez, F. F., Pfeil, B., Schirnick, C.,
 1750 Steinfeldt, R., Suzuki, T., Tilbrook, B., Velo, A., Wanninkhof, R., and Woosley, R. J.: An updated version of the
 1751 global interior ocean biogeochemical data product, GLODAPv2.2020, *Earth Syst. Sci. Data*, 12, 3653–3678,
 1752 <https://doi.org/10.5194/essd-12-3653-2020>, 2020.
- 1753 Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A.,
 1754 Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner,
 1755 G.-K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M.-F.,
 1756 Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on
 1757 calcifying organisms, *Nature*, 437(7059), 681–686, <https://doi.org/10.1038/nature04095>, 2005.
- 1758 Orr, J. C., Epitalon, J.-M., and Gattuso, J.-P.: Comparison of ten packages that compute ocean carbonate chemistry,
 1759 *Biogeosciences*, 12, 1483–1510, <https://doi.org/10.5194/bg-12-1483-2015>, 2015.
- 1760 Orr, J. C., Epitalon, J.-M., Dickson, A. G., and Gattuso, J.-P.: Routine uncertainty propagation for the Marine



- 1761 Carbon Dioxide System, *Marine Chemistry*, 207, 84–107. <https://doi.org/10.1016/j.marchem.2018.10.006>, 2018.
- 1762 Oschlies, A., Bach, L. T., Fennel, K., Gattuso, J.-P., and Mengis, N.: Perspectives and challenges of marine carbon
1763 dioxide removal, *Frontiers in Climate*, 6, <https://doi.org/10.3389/fclim.2024.1506181>, 2025.
- 1764 Padin, X. A., Velo, A., & Pérez, F. F.: Arios: A database for ocean acidification assessment in the Iberian Upwelling
1765 System (1976–2018), *Earth System Science Data*, 12(4), 2647–2663, <https://doi.org/10.5194/essd-12-2647-2020>,
1766 2020.
- 1767 Palacio-Castro, A. M., Enochs, I. C., Besemer, N., Boyd, A., Jankulak, M., Kolodziej, G., Hirsh, H. K., Webb, A. E.,
1768 Towle, E. K., Kelble, C., Smith, I., and Manzello, D. P.: Coral Reef carbonate chemistry reveals interannual,
1769 seasonal, and spatial impacts on ocean acidification off Florida, *Global Biogeochemical Cycles*, 37(12),
1770 <https://doi.org/10.1029/2023gb007789>, 2023.
- 1771 Perez, F. F., and Fraga, F.: Association constant of fluoride and hydrogen ions in seawater. *Marine Chemistry*,
1772 21(2), 161–168, [https://doi.org/10.1016/0304-4203\(87\)90036-3](https://doi.org/10.1016/0304-4203(87)90036-3), 1987.
- 1773 Pérez, F. F., Velo, A., Padin, X. A., Doval, M. D., and Prego, R.: ARIOS Database: An Acidification Ocean
1774 Database for the Galician Upwelling Ecosystem, Instituto de Investigaciones Marinas, Consejo Superior de
1775 Investigaciones Científicas (CSIC), <https://doi.org/10.20350/digitalCSIC/12498>, 2020.
- 1776 Pfeil, B., Olsen, A., Bakker, D. C. E., Hankin, S., Koyuk, H., Kozyr, A., Malczyk, J., Manke, A., Metzl, N., Sabine,
1777 C. L., Akl, J., Alin, S. R., Bates, N., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J., Chavez, F.
1778 P., Chen, A., Cosca, C., Fassbender, A. J., Feely, R. A., González-Dávila, M., Goyet, C., Hales, B., Hardman-
1779 Mountford, N., Heinze, C., Hood, M., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Jones, S.
1780 D., Key, R. M., Körtzinger, A., Landschützer, P., Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat,
1781 L., Midorikawa, T., Mintrop, L., Miyazaki, C., Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y.,
1782 Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K., Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Santana-
1783 Casiano, J. M., Salisbury, J., Sarma, V. V. S. S., Schlitzer, R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I.,
1784 Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K., Telszewski, M., Thomas, H., Tilbrook, B., Tjiputra, J.,
1785 Vandemark, D., Veness, T., Wanninkhof, R., Watson, A. J., Weiss, R., Wong, C. S., and Yoshikawa-Inoue, H.: A
1786 uniform, quality controlled Surface Ocean CO₂ Atlas (SOCAT), *Earth Syst. Sci. Data*, 5, 125–143,
1787 <https://doi.org/10.5194/essd-5-125-2013>, 2013.
- 1788 Qi, D., Ouyang, Z., Chen, L., Wu, Y., Lei, R., Chen, B., Feely, R. A., Anderson, L. G., Zhong, W., Lin, H.,
1789 Polukhin, A., Zhang, Y., Zhang, Y., Bi, H., Lin, X., Luo, Y., Zhuang, Y., He, J., Chen, J. and Cai, W.-J.: Climate
1790 change drives rapid decadal acidification in the Arctic Ocean from 1994 to 2020, *Science*, 377, 1544–1550,
1791 [doi:10.1126/science.abo0383](https://doi.org/10.1126/science.abo0383), 2022.
- 1792 Resplandy, L., Hogikyan, A., Müller, J. D., Najjar, R. G., Bange, H. W., Bianchi, D., Weber, T., Cai, W. -J., Doney,



- 1793 S. C., Fennel, K., Gehlen, M., Hauck, J., Lacroix, F., Landschützer, P., Le Quéré, C., Roobaert, A., Schwinger, J.,
 1794 Berthet, S., Bopp, L., Chau, T. T. T., Dai, M., Gruber, N., Ilyina, T., Kock, A., Manizza, M., Lachkar, Z., Laruelle,
 1795 G. G., Liao, E., Lima, I. D., Nissen, C., Rödenbeck, C., Séférian, R., Toyama, K., Tsujino, H., and Regnier, P.: A
 1796 synthesis of global coastal Ocean Greenhouse Gas Fluxes, *Global Biogeochemical Cycles*, 38(1),
 1797 <https://doi.org/10.1029/2023gb007803>, 2024.
- 1798 Revelle, R., and Suess, H. E.: Carbon dioxide exchange between atmosphere and ocean and the question of an
 1799 increase of atmospheric CO₂ during the past decades. *Tellus*, 9(1), 18–27, <https://doi.org/10.1111/j.2153->
 1800 3490.1957.tb01849.x, 1957.
- 1801 Rödenbeck, C., Keeling, R. F., Bakker, D. C., Metzl, N., Olsen, A., Sabine, C., and Heimann, M.: Global surface-
 1802 ocean *p*CO₂ and sea-air CO₂ flux variability from an observation-driven ocean mixed-layer scheme, *Ocean Science*,
 1803 9(2), 193–216, <https://doi.org/10.5194/os-9-193-2013>, 2013.
- 1804 Rödenbeck, Christian, DeVries, T., Hauck, J., Le Quéré, C., & Keeling, R. F.: Data-based estimates of interannual
 1805 sea-air CO₂ flux variations 1957–2020 and their relation to environmental drivers, *Biogeosciences*, 19(10), 2627–
 1806 2652, <https://doi.org/10.5194/bg-19-2627-2022>, 2022.
- 1807 Roobaert, A.: Regnier, P., Landschützer, P., Laruelle, G. G.: A novel sea surface partial pressure of carbon dioxide
 1808 (*p*CO₂) data product for the global coastal ocean resolving trends over the 1982–2020 period (NCEI Accession
 1809 0279118), NOAA National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/4sde-p068>,
 1810 2023.
- 1811 Roobaert, A., Regnier, P., Landschützer, P., and Laruelle, G. G.: A novel sea surface *p*CO₂-product for the global
 1812 coastal ocean resolving trends over 1982–2020, *Earth System Science Data*, 16(1), 421–441,
 1813 <https://doi.org/10.5194/essd-16-421-2024>, 2024.
- 1814 Sabine, C. L., Feely, R. A., Gruber, N., Key, R. M., Lee, K., Bullister, J. L., Wanninkhof, R., Wong, C. S., Wallace,
 1815 D. W., Tilbrook, B., Millero, F. J., Peng, T.-H., Kozyr, A., Ono, T., and Rios, A. F.: The oceanic sink for
 1816 anthropogenic CO₂, *Science*, 305(5682), 367–371, <https://doi.org/10.1126/science.1097403>, 2004.
- 1817 Sabine, C. L., Key, R. M., Kozyr, A., Feely, R. A., Wanninkhof, R., Millero, F. J., Peng, T.-H., Bullister, J. L., and
 1818 Lee, K.: Global Ocean Data Analysis Project (GLODAP): Results and data. ORNL/CDIAC-145, NDP-083, Carbon
 1819 Dioxide Information Analysis Center, Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, TN,
 1820 110 pp. plus 6 Appendices, 2005.
- 1821 Sabine, C. L., Feely, R. A., Millero, F. J., Dickson, A. G., Langdon, C., Mecking, S., and Greeley, D.: Decadal
 1822 changes in Pacific Carbon, *Journal of Geophysical Research: Oceans*, 113(C7),
 1823 <https://doi.org/10.1029/2007jc004577>, 2008.



- 1824 Sabine, C. L., Hankin, S., Koyuk, H., Bakker, D. C. E., Pfeil, B., Olsen, A., Metzl, N., Kozyr, A., Fassbender, A.,
- 1825 Manke, A., Malczyk, J., Akl, J., Alin, S. R., Bellerby, R. G. J., Borges, A., Boutin, J., Brown, P. J., Cai, W.-J.,
- 1826 Chavez, F. P., Chen, A., Cosca, C., Feely, R. A., González-Dávila, M., Goyet, C., Hardman-Mountford, N., Heinze,
- 1827 C., Hoppema, M., Hunt, C. W., Hydes, D., Ishii, M., Johannessen, T., Key, R. M., Körtzinger, A., Landschützer, P.,
- 1828 Lauvset, S. K., Lefèvre, N., Lenton, A., Lourantou, A., Merlivat, L., Midorikawa, T., Mintrop, L., Miyazaki, C.,
- 1829 Murata, A., Nakadate, A., Nakano, Y., Nakaoka, S., Nojiri, Y., Omar, A. M., Padin, X. A., Park, G.-H., Paterson, K.,
- 1830 Perez, F. F., Pierrot, D., Poisson, A., Ríos, A. F., Salisbury, J., Santana-Casiano, J. M., Sarma, V. V. S. S., Schlitzer,
- 1831 R., Schneider, B., Schuster, U., Sieger, R., Skjelvan, I., Steinhoff, T., Suzuki, T., Takahashi, T., Tedesco, K.,
- 1832 Telszewski, M., Thomas, H., Tilbrook, B., Vandemark, D., Veness, T., Watson, A. J., Weiss, R., Wong, C. S., and
- 1833 Yoshikawa-Inoue, H.: Surface Ocean CO₂ Atlas (SOCAT) gridded data products, *Earth Syst. Sci. Data*, 5, 145–153,
- 1834 <https://doi.org/10.5194/essd-5-145-2013>, 2013.
- 1835 Sarma, V. V. S. S., Krishna, M. S., Paul, Y. S., and Murty, V. S. N.: Observed changes in ocean acidity and carbon
- 1836 dioxide exchange in the coastal bay of Bengal—A link to air pollution, *Tellus B: Chemical and Physical*
- 1837 *Meteorology*, 67(1), 24638, <https://doi.org/10.3402/tellusb.v67.24638>, 2015.
- 1838 Schimel, D. S., and Carroll, D.: Carbon cycle–climate feedbacks in the post-paris world, *Annual Review of Earth*
- 1839 *and Planetary Sciences*, 52(1), <https://doi.org/10.1146/annurev-earth-031621-081700>, 2024.
- 1840 Schoderer, M., Bittig, H., Klein, B., Hägele, R., Steinhoff, T., Castro-Morales, K., Cotrim da Cunha, L., Hornidge,
- 1841 A., and Körtzinger, A.: From individual observations to global assessments: Tracing the Marine Carbon Knowledge
- 1842 Value Chain, *Ocean and Society*, 2, <https://doi.org/10.17645/oas.8891>, 2024.
- 1843 Sharp, J. D., Pierrot, D., Humphreys, M. P., Epitalon, J.-M., Orr, J. C., Lewis, E. R., Wallace, D. W. R.: CO2SYSv3
- 1844 for MATLAB (Version v3.2.1), Zenodo, <http://doi.org/10.5281/zenodo.3950562>, 2023.
- 1845 Sharp, J. D., Jiang, L.-Q., Carter, B. R., Lavin, P. D., Yoo, H., Cross, S. L. RFR-LME Ocean Acidification Indicators
- 1846 from 1998 to 2023 (NCEI Accession 0287551). NOAA National Centers for Environmental Information. Dataset.
- 1847 <https://doi.org/10.25921/h8vw-e872>, 2024a..
- 1848 Sharp, J. D., Jiang, L.-Q., Carter, B. R., Lavin, P. D., Yoo, H., and Cross, S. L.: A mapped dataset of surface ocean
- 1849 acidification indicators in large marine ecosystems of the United States, *Sci. Data*, 11, 715,
- 1850 <https://doi.org/10.1038/s41597-024-03530-7>, 2024b.
- 1851 Sridevi, B., and Sarma, V. V. S. S.: Role of river discharge and warming on ocean acidification and pco₂ levels in
- 1852 the Bay of Bengal, *Tellus B: Chemical and Physical Meteorology*, 73 (1), 1–20,
- 1853 <https://doi.org/10.1080/16000889.2021.1971924>, 2021.
- 1854 Sutton, A. J., Feely, R. A., Maenner-Jones, S., Musielwicz, S., Osborne, J., Dietrich, C., Monacci, N., Cross, J.,
- 1855 Bott, R., Kozyr, A.: Autonomous seawater partial pressure of carbon dioxide (pCO₂) and pH time series from 40



- 1856 surface buoys between 2004 and 2017 (NCEI Accession 0173932), NOAA National Centers for Environmental
- 1857 Information, Dataset, <https://doi.org/10.7289/v5db8043>, 2018.
- 1858 Sutton, A. J., Feely, R. A., Maenner-Jones, S., Musielwicz, S., Osborne, J., Dietrich, C., Monacci, N., Cross, J.,
- 1859 Bott, R., Kozyr, A., Andersson, A. J., Bates, N. R., Cai, W.-J., Cronin, M. F., De Carlo, E. H., Hales, B., Howden, S.
- 1860 D., Lee, C. M., Manzello, D. P., McPhaden, M. J., Meléndez, M., Mickett, J. B., Newton, J. A., Noakes, S. E., Noh,
- 1861 J. H., Olafsdottir, S. R., Salisbury, J. E., Send, U., Trull, T. W., Vandemark, D. C., and Weller, R. A.: Autonomous
- 1862 seawater $p\text{CO}_2$ and pH time series from 40 surface buoys and the emergence of anthropogenic trends, *Earth Syst.*
- 1863 *Sci. Data*, 11, 421–439, <https://doi.org/10.5194/essd-11-421-2019>, 2019.
- 1864 Suzuki, T., M. Ishii, M. Aoyama, J. R. Christian, K. Enyo, T. Kawano, R. M. Key, N. Kosugi, A. Kozyr, L. A.
- 1865 Miller, A. Murata, T. Nakano, T. Ono, T. Saino, K. Sasaki, D. Sasano, Y. Takatani, M. Wakita and C. Sabine:
- 1866 PACIFICA Data Synthesis Project. ORNL/CDIAC-159, NDP-092. Carbon Dioxide Information Analysis Center,
- 1867 Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee. doi:
- 1868 10.3334/CDIAC/OTG.PACIFICA_NDP092, 2013.
- 1869 Swift, J. (2022), Java OceanAtlas Data, https://joa.ucsd.edu/Data_homepage (Last accessed: 2025-02-08).
- 1870 Takahashi, T., Feely, R. A., Weiss, R. F., Wanninkhof, R. H., Chipman, D. W., Sutherland, S. C., and Takahashi, T.
- 1871 T.: Global Air-Sea Flux of CO_2 : An estimate based on measurements of sea–air $p\text{CO}_2$ difference, *Proceedings of the*
- 1872 *National Academy of Sciences*, 94(16), 8292–8299, <https://doi.org/10.1073/pnas.94.16.8292>, 1997.
- 1873 Takahashi, T., Sutherland, S. C., Sweeney, C., Poisson, A., Metzl, N., Tilbrook, B., Bates, N., Wanninkhof, R.,
- 1874 Feely, R. A., Sabine, C., Olafsson, J., and Nojiri, Y.: Global sea–air CO_2 flux based on climatological surface ocean
- 1875 $p\text{CO}_2$, and seasonal biological and temperature effects, *Deep Sea Research Part II: Topical Studies in*
- 1876 *Oceanography*, 49(9–10), 1601–1622, [https://doi.org/10.1016/s0967-0645\(02\)00003-6](https://doi.org/10.1016/s0967-0645(02)00003-6), 2002.
- 1877 Takahashi, T., Sutherland, S. C., Wanninkhof, R., Sweeney, C., Feely, R. A., Chipman, D. W., Hales, B., Friederich,
- 1878 G., Chavez, F., Sabine, C., Watson, A., Bakker, D. C. E., Schuster, U., Metzl, N., Yoshikawa-Inoue, H., Ishii, M.,
- 1879 Midorikawa, T., Nojiri, Y., Körtzinger, A., Steinhoff, T., and de Baar, H. J. W.: Climatological mean and decadal
- 1880 change in Surface Ocean $p\text{CO}_2$, and net sea–air CO_2 flux over the Global Oceans, *Deep Sea Research Part II:*
- 1881 *Topical Studies in Oceanography*, 56(8–10), 554–577, <https://doi.org/10.1016/j.dsr2.2008.12.009>, 2009.
- 1882 Takahashi, T., Sutherland, S. C., Kozyr, A.: LDEO Database (Version 2019): Global Ocean Surface Water Partial
- 1883 Pressure of CO_2 Database: Measurements Performed During 1957–2019 (NCEI Accession 0160492), NOAA
- 1884 National Centers for Environmental Information, Dataset, [https://doi.org/10.3334/cdiac/otg.ndp088\(v2015\)](https://doi.org/10.3334/cdiac/otg.ndp088(v2015)), 2017.
- 1885 Takahashi, T., Sutherland, S. C., and Kozyr, A.: Global Ocean Surface Water Partial Pressure of CO_2 Database:
- 1886 Measurements Performed During 1957–2019 (LDEO Database Version 2019) (NCEI Accession 0160492), Version
- 1887 9.9, NOAA National Centers for Environmental Information [data set],



- 1888 [https://doi.org/10.3334/CDIAC/OTG.NDP088\(V2015\)](https://doi.org/10.3334/CDIAC/OTG.NDP088(V2015)), 2021.
- 1889 Tanhua, T., Jones, E. P., Jeansson, E., Jutterström, S., Smethie, W. M., Wallace, D. W., and Anderson, L. G.:
1890 Ventilation of the Arctic Ocean: Mean ages and inventories of Anthropogenic CO₂ and CFC-11, *Journal of*
1891 *Geophysical Research: Oceans*, 114(C1), <https://doi.org/10.1029/2008jc004868>, 2009.
- 1892 Tanhua, T., van Heuven, S., Key, R. M., Velo, A., Olsen, A., and Schirnack, C.: Quality control procedures and
1893 methods of the CARINA database, *Earth Syst. Sci. Data*, 2, 35–49, <https://doi.org/10.5194/essd-2-35-2010>, 2010.
- 1894 Tans, P., and Keeling, R.: Trends in atmospheric carbon dioxide, NOAA Global Monitoring Laboratory (Boulder,
1895 Colorado, USA) and Scripps Institution of Oceanography (LaJolla, California, USA),
1896 <https://gml.noaa.gov/ccgg/trends/> and <https://scrippsco2.ucsd.edu/>, 2025.
- 1897 Terhaar, J., Orr, J. C., Ethé, C., Regnier, P., and Bopp, L.: Simulated Arctic Ocean response to doubling of riverine
1898 carbon and nutrient delivery, *Global Biogeochemical Cycles*, 33, 1048–1070,
1899 <https://doi.org/10.1029/2019GB006200>, 2019.
- 1900 Terhaar, J., Tanhua, T., Stöven, T., Orr, J. C., and Bopp, L.: Evaluation of data-based estimates of anthropogenic
1901 carbon in the Arctic Ocean, *Journal of Geophysical Research: Oceans*, 125(6),
1902 <https://doi.org/10.1029/2020jc016124>, 2020.
- 1903 Terhaar, J., Torres, O., Bourgeois, T., and Kwiatkowski, L.: Arctic Ocean acidification over the 21st century co-
1904 driven by anthropogenic carbon increases and freshening in the CMIP6 model ensemble, *Biogeosciences*, 18, 2221–
1905 2240, <https://doi.org/10.5194/bg-18-2221-2021>, 2021a.
- 1906 Terhaar, Jens, Frölicher, T. L., & Joos, F.: Southern Ocean anthropogenic carbon sink constrained by sea surface
1907 salinity, *Science Advances*, 7(18), <https://doi.org/10.1126/sciadv.abd5964>, 2021b.
- 1908 Terhaar Jens, Frölicher Thomas L., Joos Fortunat (2021c). Simulated and constrained Southern Ocean carbon sink
1909 from 1850 to 2100 based on CMIP5 and CMIP6 models. SEANOE. <https://doi.org/10.17882/103938>
- 1910 Terhaar, J., Frölicher, T. L., and Joos, F.: Observation-constrained estimates of the Global Ocean Carbon Sink from
1911 earth system models, *Biogeosciences*, 19(18), 4431–4457, <https://doi.org/10.5194/bg-19-4431-2022>, 2022a.
- 1912 Terhaar Jens, Frölicher Thomas Lukas, Joos Fortunat (2022b). Simulated and constrained ocean carbon sink from
1913 1850 to 2100 based on CMIP6 models. SEANOE. <https://doi.org/10.17882/103934>
- 1914 Terhaar, J., Frölicher, T. L., and Joos, F.: Ocean acidification in emission-driven temperature stabilization scenarios:
1915 The role of TCRE and non-CO₂ greenhouse gases, *Environmental Research Letters*, 18(2), 02403,
1916 <https://doi.org/10.1088/1748-9326/acaf91>, 2023.
- 1917 Terhaar Jens, Tanhua Toste, Stöven Tim, Orr James C, Bopp Laurent (2024). Observation-based anthropogenic



- 1918 carbon estimates in the Arctic Ocean. SEANO. <https://doi.org/10.17882/103920>
- 1919 van Heuven, S., Pierrot, D., Rae, J. W. B., Lewis, E., Wallace, D. W. R.: MATLAB Program Developed for CO₂
- 1920 System Calculations, ORNL/CDIAC-105b, Carbon Dioxide Information Analysis Center, Oak Ridge National
- 1921 Laboratory, Oak Ridge, TN, USA, 2011.
- 1922 van Hooijdonk, R.: Modeled ocean acidification data in the Gulf of Mexico and wider Caribbean using satellites and
- 1923 climate model data for the Ocean Acidification Products for the Gulf of Mexico and East Coast project from 2014-
- 1924 01-01 to 2020-12-31 (NCEI Accession 0245950), NOAA National Centers for Environmental Information, Dataset,
- 1925 <https://doi.org/10.25921/tt1c-dx53>, 2022.
- 1926 Velo, A., Pérez, F. F., Tanhua, T., Gilcoto, M., Ríos, A. F., and Key, R. M.: Total alkalinity estimation using MLR
- 1927 and neural network techniques, *J. Mar. Syst.*, 111–112, 11–18, <https://doi.org/10.1016/j.jmarsys.2012.09.002>, 2013.
- 1928 Wanninkhof, R., Pierrot, D., Sullivan, K., Barbero, L., and Triñanes, J.: Gridded surface water fugacity of CO₂
- 1929 observations, and calculated pH, aragonite saturation state and air–sea CO₂ fluxes in the northern Caribbean Sea
- 1930 from 2002 through 2019 (NCEI Accession 0207749), NOAA National Centers for Environmental Information,
- 1931 Dataset, <https://doi.org/10.25921/2swk-9w56>, 2019.
- 1932 Wanninkhof, R., Pierrot, D., Sullivan, K., Barbero, L., and Triñanes, J.: A 17-year dataset of surface water fugacity
- 1933 of CO₂ along with calculated pH, aragonite saturation state and air–sea CO₂ fluxes in the northern Caribbean Sea,
- 1934 *Earth Syst. Sci. Data*, 12, 1489–1509, <https://doi.org/10.5194/essd-12-1489-2020>, 2020.
- 1935 Wanninkhof, R., Triñanes, J., Pierrot, D., Munro, D. R., Sweeney, C., and Fay, A. R.: AOML_ET: Partial pressure
- 1936 of CO₂ (*p*CO₂) and sea-air CO₂ fluxes for the global ocean, along with the predictor variables from 1998-01-01 to
- 1937 2023-12-30, using an Extra Trees (extremely randomized trees) machine learning (NCEI Accession 0298989),
- 1938 NOAA National Centers for Environmental Information, Dataset, <https://doi.org/10.25921/0s8y-q287>, 2024.
- 1939 Wanninkhof, R., Triñanes, J., Pierrot, D., Munro, D. R., Sweeney, C., and Fay, A. R.: Trends in sea-air CO₂ fluxes
- 1940 and sensitivities to atmospheric forcing using an extremely randomized trees machine learning approach, *Global*
- 1941 *Biogeochemical Cycles*, 39, <https://doi.org/10.1029/2024GB008315>, 2025.
- 1942 Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G., Landschützer, P., Woolf, D., and Goddijn-
- 1943 Murphy, L.: Revised estimates of ocean-atmosphere CO₂ flux are consistent with ocean carbon inventory, *Nature*
- 1944 *Communications*, 11(1), 1–6, <https://doi.org/10.1038/s41467-020-18203-3>, 2020.
- 1945 Watson, A. J., Schuster, U., Shutler, J. D., Holding, T., Ashton, I. G., Landschützer, P., Woolf, D., and Goddijn-
- 1946 Murphy, L.: Revised estimates of ocean-atmosphere CO₂ flux accounting for near-surface temperature and salinity
- 1947 deviations from 1985-01-01 to 2019-12-31 (NCEI Accession 0301544), NOAA National Centers for Environmental
- 1948 Information, Dataset, <https://doi.org/10.25921/2dp5-xm29>, 2025.



- 1949 Wilkinson, M. D., Dumontier, M., Aalbersberg, I. J., Appleton, G., Axton, M., Baak, A., Blomberg, N., Boiten, J.-
- 1950 W., da Silva Santos, L. B., Bourne, P. E., Bouwman, J., Brookes, A. J., Clark, T., Crosas, M., Dillo, I., Dumon, O.,
- 1951 Edmunds, S., Evelo, C. T., Finkers, R., Gonzalez-Beltran, A., Gray, A. J. G., Groth, P., Goble, C., Grethe, J. S.,
- 1952 Heringa, J., Hoen, P., Hooft, R., Kuhn, T., Kok, R., Kok, J., Scott J. Lusher, S. J., Martone, M. E., Mons, A., Packer,
- 1953 A. L., Persson, B., Rocca-Serra, P., Roos, M., van Schaik, R., Sansone, S.-A., Schultes, E., Sengstag, T., Slater, T.,
- 1954 Strawn, G., Swertz, M. A., Thompson, M., van der Lei, J., van Mulligen, E., Velterop, J., Waagmeester, A.,
- 1955 Wittenburg, P., Wolstencroft, K., Zhao, J., Mons, B.: The Fair Guiding Principles for Scientific Data Management
- 1956 and Stewardship, *Scientific Data*, 3(1), <https://doi.org/10.1038/sdata.2016.18>, 2016.
- 1957 Winn, C. D., MacKenzie, F. T., Carrillo, C. J., Sabine, C. L., and Karl, D. M.: Air-sea carbon dioxide exchange in
- 1958 the North Pacific Subtropical Gyre: Implications for the global carbon budget, *Global Biogeochem Cycles*, 8, 157-
- 1959 163, <https://doi.org/10.1029/94GB00387>, 1994.
- 1960 Winn, C. D., Li, Y.-H., Mackenzie, F. T., and Karl, D. M.: Rising surface ocean dissolved inorganic carbon at the
- 1961 Hawaii Ocean Time-series site, *Mar. Chem.*, 60, 33-47, [https://doi.org/10.1016/S0304-4203\(97\)00085-6](https://doi.org/10.1016/S0304-4203(97)00085-6), 1998.
- 1962 Wu, Z., Lu, W., Roobaert, A., Song, L., Yan, X.-H., and Cai, W.-J.: A machine-learning reconstruction of sea
- 1963 surface $p\text{CO}_2$ in the North American Atlantic Coastal Ocean Margin from 1993 to 2021, *Earth Syst. Sci. Data*.
- 1964 <https://doi.org/10.5194/essd-17-43-2025>, 2025.
- 1965 Zeng, J., Iida, Y., Matsunaga, T., Shirai, T.: Surface ocean CO_2 concentration and air-sea flux estimate by machine
- 1966 learning with modelled variable trends, *Front. Mar. Sci.*, 9, 989233, <https://doi.org/10.3389/fmars.2022.989233>,
- 1967 2022.
- 1968 Zeng, J.: NIES-ML3 ensemble product of surface ocean CO_2 concentrations and air-sea CO_2 fluxes reconstructed by
- 1969 using three machine learning models with new CO_2 trends, National Institute for Environmental Studies (NIES),
- 1970 Japan, <https://doi.org/10.17595/20220311.001>, 2022.
- 1971 Zhang, H., Menemenlis, D., and Fenty, I. G.: ECCO LLC270 ocean-ice state estimate (Tech. Rep.), Jet Propulsion
- 1972 Laboratory, California Institute of Technology, <http://hdl.handle.net/1721.1/119821>, 2018.
- 1973 Zweng, M. M., Reagan, J. R., Antonov, J. I., Locarnini, R. A., Mishonov, A. V., Boyer, T. P., Garcia, H. E.,
- 1974 Baranova, O. K., Johnson, D. R., Seidov, D., and Biddle, M. M.: World Ocean Atlas 2013, Volume 2: Salinity,
- 1975 edited by: Levitus, S. and Mishonov, A., NOAA Atlas NESDIS 74, 39 pp., 2013.
- 1976 Zweng, M. M., Reagan, J. R., Seidov, D., Boyer, T. P., Locarnini, R. A., Garcia, H. E., Mishonov, A. V., Baranova,
- 1977 O. K., Weathers, K. W., Paver, C. R., Smolyar, I. V.: World Ocean Atlas 2018, Volume 2: Salinity, edited by
- 1978 Mishonov, A., NOAA Atlas NESDIS 82, 2019.