

Left knee MRI - 25 September 2026

Report transcription; original PDF unavailable.

Left knee MRI - 25 September 2026

TRANSCRIPTION of the report supplied by the patient; original PDF unavailable.

Clinical history

Assess cause of FFD and progress of semimembranosis oedema on previous scan

Technique

Multiplanar, multisequence MRI without gadolinium.

Comparison

No prior study for comparison.

Findings

Cruciate ligaments are intact.

No meniscal tear. Collateral ligaments are intact. Medial and lateral compartment articular cartilage surfaces are well-maintained with no established osteoarthritic change.

Patellofemoral joint is normally aligned with preserved articular cartilage surfaces. Extensor mechanism tendons and iliotibial band insertion are intact.

No joint effusion or intra-articular loose body. No Baker's cyst. No concerning bone lesion or soft tissue mass. The major neurovascular structures are structurally normal.

There is a focal area of low-grade muscle oedema within the distal muscle belly of the semimembranosus at the distal myotendinous junction, suggesting a subacute low-grade strain. No discrete haematoma or tear defect. The distal free tendon and its distal insertion are intact and of normal signal intensity.

The remainder of the muscles and tendons around the knee joints are normal.

Conclusion

Mild oedema within the distal muscle belly of semimembranosus, at the distal myotendinous junction, suggestive of a subacute low-grade muscle strain. The distal tendon is intact and inserts normally.

No internal joint derangement.