Scalar–Tensor Hydrodynamics

Gravity as the Response of a Vacuum Medium:
A Modified Gravity Framework for Vacuum Viscosity and Cyclic Vacuum Cosmology

Keith Hays

Revised Formalism — Version 3.12 (draft)

Abstract

We propose a hydrodynamic extension of General Relativity in which the vacuum is modeled as a dynamic, high-energy medium described by a complex scalar field $\Phi = \rho\, e^{i\theta}$, whose state is written ϕ. Rather than treating the vacuum as a fixed geometric background plus a constant cosmological term, the framework assigns the vacuum its own stress-energy and pressure. The formulation separates the Einstein geometric tensor from the vacuum's own stresses, keeps a place for a viscous stress that carries nothing in the settled medium, introduces a dynamical scalar-field sector, replaces a fixed cosmological constant with a field-dependent pressure contribution, and allows the effective gravitational coupling to depend on the local vacuum state. The framework is intended to investigate whether cosmological expansion, galactic dynamical anomalies, and the transition between localized and delocalized excitations can arise from one underlying medium. The medium is taken as primary. What are conventionally called particles are treated as localized excitations of it, and spacetime geometry, with Einstein's equation, as the effective description of how its state is experienced. Gravity is proposed to be the medium's response to concentrated energy. The medium has one speed, at which every disturbance travels, and gravity is where that speed varies from place to place; no speed is fixed, and the measured speed of light is the local carrying speed read by rulers and clocks made of the same medium. Two principles of the framework then give General Relativity's geometry outside a non-spinning mass, including its point of no return and the observed size of a black hole's shadow. Around a spinning mass, the same medium gives the measured frame dragging to first order. The medium rebuilds every pattern continuously, and the rebuild has two sides: building, which makes matter and time, and reclaiming, which unmakes them and is antimatter. From that rebuild a law is stated, in which no time outside the medium appears, gravity's strength is set by the waves the medium sustains, and neutrinoless double-beta decay must occur.

This document is a theoretical proposal, not an established physical theory. The equations below define a research program whose consistency, stability, conservation laws, and observational predictions must be demonstrated.

A note for readers, including automated ones. This document has 59 pages in 18 parts. Each page's footer gives its page number, the page count and the part it belongs to, with that part's checksum token; each part ends with a line repeating its token. The last page is a reading manifest listing every part, its first page and its token, followed by a document digest and the words "End of document". A complete reading reaches that line. A summary or review that cannot give the document digest, or that is missing any page number or part token, was made from an incomplete reading and should say so.

Preface: What This Paper Is For

The premise

There is no empty space.

Everything this paper describes happens inside a single continuous medium, called here the Ocean. The name is a metaphor, and one feature of real oceans must be discarded at once: this Ocean has no

edges. It has no shore and no surface. There is no sky above it, no floor beneath it and no air over it. It is not held in a basin, and it does not sit inside a space that could exist without it. There is no outside. The edge of what we can see is a limit of seeing, not an edge of the Ocean. Where things seem empty, the Ocean is there, quiet. Where they seem full, the Ocean is there, gathered. Nothing in this framework exists apart from it, beside it or beyond it.

What we call matter is the Ocean's own energy, gathered and circulating in place. What physics calls a particle is not an object but a pattern the Ocean keeps re-making. When a particle "moves", the place where the pattern is re-made has shifted, the way a wave crosses a stadium while nobody leaves their seat. Light is a ripple in the same medium. A galaxy is a vast region of gathered energy and current. Every pattern exists against, alongside or within others, adding to or taking from them, and every one is ultimately the Ocean's response to what the rest of it has done. For that reason this document uses the word "particle" only when reporting other authors' theories, and marks it as theirs.

The Ocean has one speed: the speed at which it carries a disturbance. Light, the inner workings of matter and every signal run at it; only the Ocean's own flow is free of it. What physics calls the speed of light is the Ocean's local carrying speed, measured everywhere with the same value because rulers and clocks are themselves patterns of the Ocean and run at the same speed. There is no fixed speed of light anywhere in this framework: a fixed number for it is a conversion between units of length and time, not a property of nature.

This work began with gravity. Einstein's theory describes what gravity does with extraordinary precision: clocks run slow near a mass, space stretches, paths curve. It does not say why concentrated energy should do this. The proposal here is that gravity is the medium's response. Concentrated energy changes the state of the Ocean around it, the Ocean runs slower there, and every pattern in it, light included, leans toward where it runs slower. On this view Einstein's equations remain the correct description, and the Ocean is offered as the reason they are correct, as the gas laws stayed correct when atoms were found to explain them. Whether the framework can carry that claim is the test it must pass: every pattern must lean by exactly the same amount, and the stretching of space must match the slowing of time exactly as Einstein requires.

The problem

The same premise bears on the largest open problem in cosmology.

Two of the largest numbers in cosmology are placeholders. Roughly a quarter of the contents of the observable universe is labeled *dark matter*: a substance inferred entirely from its gravitational pull on things we can see, and never once detected directly. Most of the remainder is labeled *dark energy*, which is less a substance than a name for an observation — that cosmic expansion is speeding up when everything we understand says it should be slowing down. Together these two labels cover something like ninety-five per cent of what the observable universe contains. Neither has an agreed physical identity.

Standard cosmology handles this competently. It adds an invisible cold matter component, and a constant term to Einstein's equations, and the resulting fit to observation is superb — good enough that the model deserves the confidence it has. But a superb fit obtained by inserting two unexplained quantities is not the same thing as an explanation. It is a description with two holes in it, and the holes have names.

This paper asks whether both holes might have one filling.

A note on where these numbers came from

There is a precedent in this subject for inserting a quantity because the answer is expected rather than because anything has been measured. In 1917 Einstein added a term to his field equations — the cosmological constant, Λ — for no reason beyond his conviction that the universe, as he conceived it, ought to be static, and the inconvenient fact that his equations refused to make it so. No observation supported the term. It was a number chosen to produce an anticipated result. When Hubble demonstrated a little over a decade later that the observable universe is expanding after all, Einstein abandoned it.

It is worth being precise about what happened next, because the common retelling is wrong in a way that matters. Λ returned in 1998, but not as Einstein's unmeasured assumption quietly surviving. It returned because observations of distant supernovae showed cosmic expansion to be accelerating, and Λ was the simplest term capable of describing that. The acceleration is real and measured. What Λ does not do is explain it: no theory predicts the constant's value, and the various attempts to calculate it from first principles fail by a margin discussed below. The term records what the expansion does and is silent on why.

Dark matter has an entirely separate and considerably stronger pedigree. Fritz Zwicky noticed in 1933 that galaxies in the Coma cluster move too fast for the visible mass to hold them together. Vera Rubin's rotation-curve measurements in the 1970s found the same discrepancy inside individual galaxies. Gravitational lensing, the structure of the cosmic microwave background, and the separation of mass from gas in colliding clusters have since confirmed it from independent directions. The evidence is not thin, and this document does not dispute any of it.

The objection is therefore not that these quantities were invented. It is that in both cases a measured discrepancy was given a name, and the name was then promoted to a substance. Λ is a parameter fitted to the expansion history rather than derived from anything. Dark matter is a material known solely by its gravity, which forty years of increasingly sensitive detectors have failed to find by any other means. Each is a placeholder that has been in service long enough to be mistaken for an answer.

If the framework set out here is even approximately correct, nothing observational changes. The rotation curves remain exactly as measured. So do the lensing maps, the microwave background, and the supernova data establishing acceleration. What dissolves is the pair of substances invoked to account for them: they stop being two unknown constituents of the observable universe and become two behaviors of one medium. The measurements survive intact. Only their cause is reassigned.

Where our two best theories disagree

General Relativity and quantum field theory are the most thoroughly confirmed descriptions of nature ever constructed, and they are known to be mutually incompatible. In ordinary practice this causes little trouble, because the two rarely describe the same situation: one governs the very large, the other the very small, and the regimes where both matter at once — the first instants after the big bang, the interior of a black hole — are exactly the regimes we cannot observe.

There is one glaring exception, and it is directly relevant here. The two theories overlap on the question of what the vacuum is, and there they do not disagree by a little. General Relativity treats empty space as genuinely empty: a smooth, featureless backdrop with no properties of its own. Quantum field theory describes the same emptiness as the most active object in physics — fields in their lowest state, never still, producing effects that have been measured in the laboratory for seventy years.

Ask both theories what that activity should weigh, and the answers differ by something between sixty and a hundred and twenty orders of magnitude, depending on where the calculation is cut off. This is not a discrepancy to be tidied away with a correction factor. It is routinely described as the worst quantitative prediction in the history of science, and it has stood unresolved for decades.

The observation that motivates this document is that the sharpest disagreement between our two best theories is a disagreement about whether the vacuum is a thing. One says no. The other says emphatically yes, and has the laboratory results to support it. What follows takes the second answer seriously and asks a further question that neither theory addresses: if the vacuum is a thing, does it also have mechanical properties — density, pressure, flow, resistance to being stirred — and if so, what would we expect to see?

What a medium can do

General Relativity, read as it is usually taught, treats the vacuum as the absence of things: a stage that can be curved by what sits on it but has no density, pressure or flow of its own. A medium does things a stage cannot. It flows. It drags. It carries waves. It can be stirred, and it pushes back when you stir it. Each of those behaviors is a candidate for a phenomenon currently attributed to something unknown:

- A galaxy sits in an eddy of the medium's current, and the eddy's swirl turns the travel of its stars inward. The extra pull that standard cosmology assigns to a halo of undetected particles, in its own terms, might instead be the swirl of the eddy in which the galaxy sits.
- A medium with internal pressure that can change character over time can drive expansion without a constant pasted into the equations by hand. What is now called dark energy would be the Ocean in a particular state rather than a number in a table.
- And — this is the part developed furthest here — a medium has a smallest meaningful size. Below some scale, it cannot be disturbed in a localized way at all. That gives a principled distinction between what counts as a localized pattern and what counts as a ripple.

The ambition is that what are called dark matter and dark energy stop being two mysteries and become two behaviors of one thing.

Black holes

A black hole, on this account, is the Ocean's largest whirlpool. It does not gather a substance; it gathers organization. Whatever falls in is stripped of its own pattern until it turns with the whirlpool. Black holes recycle; they do not drain. Most of what they return comes back early, as light and jets from the matter swirling around them, and the rest leaves over immense times as faint radiation. Energy returns to the Ocean; structure does not. Nothing is gathered at its center: what falls in flows through and returns to the Ocean, much later. Its horizon is a point of no return, not a surface and not an edge of the Ocean, which stays whole across it. Section 8.2 follows a single traveler in. A spinning black hole holds its spin as the net circulation of its own Ocean, by the same rule that turns the gyroscopes around the Earth.

What this document is not

It is not a theory. It is a proposal with its homework listed.

The equations below are consistent in form, but consistency in form is cheap. A physical theory has to survive conservation laws, stability analysis, the recovery of tested results in the appropriate limits, and quantitative agreement with observations that were not used to build it. None of that has been done here. Section 14 is an itemized list of the work required, and several items on it could terminate the framework outright. They are stated plainly rather than buried, because a proposal that cannot specify how it would fail is not worth the paper it occupies.

How this document came to exist

The framework began as private speculation, written to a standard that required only that the pieces appear to fit. That is a much weaker standard than the one it has since been held to, and the gap between the two turned out to be where the interesting work was.

Nine revisions have followed, and they share a shape.

The first corrected a structural error. In the original formulation the vacuum's viscous response *replaced* the Einstein tensor — the geometric heart of General Relativity. This cannot work: a mathematical identity guarantees that the Einstein tensor is conserved, and substituting something else for it breaks the conservation of energy and momentum. The fix was to keep the geometry intact and add the vacuum's stress alongside it rather than in its place.

The second removed a number. The original drew the line between matter and vacuum at $0.511\,\mathrm{MeV}$ — the rest energy of the electron. Above the line, matter; below it, vacuum. This was an acknowledged placeholder, and it had three problems: the value was imported rather than derived, it was observer-dependent (two observers moving relative to one another would disagree about which side of the line a given thing fell), and it was an abrupt cutoff where physics prefers continuity.

The replacement drops the number entirely and asks instead how small a disturbance the medium can hold. Every medium has such a scale — in a laboratory superfluid it is called the healing length, the

shortest distance over which the medium can bend without tearing. Disturbances larger than it spread out and remain part of the medium. Disturbances smaller than it stay put, as localized patterns. The matter/vacuum distinction becomes a property of the Ocean rather than a line drawn across it.

The result worth noting is what happened next, though the ninth revision, below, reads it differently. Calibrating that scale to present conditions puts the Ocean's healing length at roughly 386 femtometers. (A femtometer is a millionth of a billionth of a meter, about the size of a proton.) That is exactly the electron's own characteristic length. Every pattern has a natural spread set by its mass, tighter the heavier it is, and the Ocean can hold in place only a pattern whose spread fits within its healing length. The electron's spread is 386 femtometers, so the electron is simply the lightest pattern the Ocean can hold in place, and everything lighter, photons and neutrinos included, spreads too wide and is necessarily a ripple instead. The healing length is the Ocean's tightest grip, not the electron's size: probed directly, the electron shows no size at all down to a scale several hundred thousand times smaller. The number that had been an unexplained input came back out as a consequence. That is the kind of exchange that makes a framework worth continuing.

The third responded to news from outside. In September 2026 a proof was announced that a fluid with internal friction can, under smooth external stirring, tear itself into a singularity in finite time — a result about ordinary fluid dynamics with no connection to this framework whatsoever. It remains unverified and does not transfer directly to a relativistic medium. But it removed an assumption this framework had been resting on without noticing: that viscosity is enough to keep a continuum well-behaved forever.

The repair was already present. The smallest-scale limit introduced in the second revision is precisely what stops a disturbance concentrating without bound — but only if the medium resists compression rather than welcoming it. That condition had never been written down. It is now a stated requirement, and if the Ocean turns out to have the other sign, the framework fails as written.

The fourth was a change of mind. The first revision had kept Einstein's geometry intact and set the Ocean beside it, and every version since had carried that arrangement forward: a stage, and a medium placed upon it. The stage was never argued for. It was inherited, from the habit of treating spacetime as the thing that exists and fields as things that happen in it. Asked directly what the stage was made of, the framework had no answer, and a medium that fills everything leaves no room for a second structure to occupy.

So the stage is removed. There is only the Ocean. Einstein's equation stays, because it describes what is measured with extraordinary precision, but its standing changes: it is now the description of how the medium's state is experienced by patterns made of that medium, not a separate structure the medium sits in. Gravity, on this account, is the medium's response to concentrated energy. That claim brings obligations the earlier arrangement did not carry, and they are set out in the technical sections; the heaviest is to derive the geometry rather than assume it.

The fifth removed a picture. Earlier versions borrowed the image of a black hole as a drain, with the medium pouring inward. Taken literally the image fails: a medium flowing in would have to pile up somewhere, and nothing piles up. Near a black hole the Ocean's state changes steeply, and that change bends every path until, at the horizon, nothing can travel outward; the medium itself stays where it is. The same reasoning gave gravity and expansion one root. Both are a change in the Ocean's state: where the change varies from place to place it bends paths, and where it is the same everywhere at once it makes the Ocean flow apart.

The sixth found the stage still standing in the equations. The fourth revision had removed it as a position, but the law written for the Ocean still measured every disturbance against a fixed speed that belonged to no part of the Ocean. Light took its speed from the Ocean's state; matter took its limit from that fixed speed; and no arrangement of the two could produce gravity. So the second speed is removed. The Ocean has one speed, the speed at which it carries a disturbance. Light, the workings of matter and every signal run at it; only the Ocean's own flow is free of it. Gravity is where that speed varies from place to place.

The seventh found the stage once more, this time in the arithmetic. The sixth had removed the second speed in words, but every equation for the Ocean was still written in the standard way, with a fixed speed

of light built into how a change in time is weighed against a change across space. So the fixed speed is removed. There is no constant c anywhere in the law: every speed in it is the Ocean's own carrying speed where it is, and the value measured everywhere is how that speed reads to rulers and clocks made of the same Ocean.

The eighth asked the Ocean to spin. Near a spinning body a gyroscope slowly turns, and so does the plane of a satellite's orbit, as if the body carried the directions around it along with its spin. Earlier versions gave the Ocean's account of a mass that does not spin, and had nothing to say about one that does. Worked out from inside the Ocean, with the Earth itself treated as Ocean, the account is this. A pattern's surroundings, the slowed Ocean around it, are re-made as it travels and carry the direction of its travel; the Earth's spinning parts together turn the Ocean around it by the amount the gyroscopes and satellites measure; and the Earth's own journey through the Ocean leaves no trace, as it must. The Ocean carries this turning in thin lines of swirl, as a rotating superfluid does, and here a borrowed number came to light. Earlier versions spaced those lines using the electron's mass, a habit taken from laboratory superfluids; in the Ocean the spacing is set instead by how fast its own state turns over. That rate falls as the Ocean expands, and frame dragging needs it to be large today. The law that allows this while keeping the Ocean as stiff as measurement requires is not yet written, and it is now on the list. The same work showed that the Ocean's turning cannot be what tells matter from antimatter; that difference needs more structure inside each point of the Ocean, as spin already does. The eighth version also gave every reader, human or automated, a way to confirm that the whole document has been read.

The ninth is the one this version records, and it asked what the Ocean does at every moment. Earlier versions said a pattern travels by being re-made farther along, and left the re-making as a picture. Here it becomes the account. Every pattern is rebuilt continuously, and the rebuild has two sides: building, which makes the pattern, and reclaiming, which takes it back down. Time is not a stage the rebuild happens on; each build makes the moment, each place makes its own time, and gravity is the uneven making of it, slower where there is more to rebuild. Antimatter is the reclaiming side. It stays hidden because it runs into a past that was never anyone's present, and the present is matter because that is the side that occupies it. At the end of a beat the building is no longer supplied, and the reclaiming takes everything away, time and disorder included, and returns the energy. From the rebuild a law is now stated, with what it settles and what it does not. One of its consequences is a rare decay that experiments are already searching for, which must occur if the Ocean is right. It also changes the second revision's reading. The strength of gravity places the Ocean's own grip, the width of a dent, near the Planck length, far below the electron. The electron's 0.511 MeV is instead the lowest energy of its own kind of wave, set by the field the Ocean's flow sustains, and 386 femtometers is the electron's own reach, not the Ocean's. The match the second revision found remains a match; what it measures has moved. It also changes the eighth revision's reading: the turning around a spinning body needs no lines.

Each revision has the same form: something the framework had not earned is removed, whether it was chosen by hand, imported or inherited, and what replaces it must then be derived rather than assumed. That a framework can absorb that process repeatedly is not evidence that it is true. It is evidence that it is the right kind of thing to keep testing — which is a lower bar, and the only one this document claims to clear.

A note on method, since it is unusual enough to be worth recording. The revisions above were worked out in dialogue with a large language model, used as an interlocutor rather than an author: a way of being argued with at the speed of thinking.

What would kill it

Five things, named here so that a reader does not have to hunt for them.

The scale gap. For the Ocean to explain galactic rotation it must be able to act across thousands of light years. Its own built-in size, the width of a dent, lies near the Planck length (Section 12.7). Read as two properties of the medium, these differ by more than fifty orders of magnitude. They are set by

different parts of the mathematics, so this is not an outright contradiction, but a framework that merely tolerates such a gap has not explained anything.

A route out exists, and it is recorded as a route rather than a result. A hurricane is hundreds of miles across, but that size is not a property of air. It belongs to the storm: to how much energy and spin were gathered in one place. Air itself has only small-scale properties. If the same holds here, the galactic scale is not built into the Ocean at all. A galaxy is not an object that stirs the medium; it is a region of the medium carrying excess energy and excess current, of which the visible stars are the dense middle and the wide slow flow is the outskirts. The Ocean would then need only one built-in size, the small one, together with the ability to carry very long, slow disturbances at almost no cost. Laboratory superfluids have that ability for a known reason, a symmetry of the underlying field. The price is that the field must be a richer object than a single real number at each point, which the framework adopts (Section 2.2), and that the calculation showing such a disturbance reproduces observed galaxies has not been done. Until it is, the gap stays on this list.

The route does not make the gap smaller. A galaxy really is about 10^{55} times larger than the Ocean's smallest structure, just as a hurricane is about 10^{14} times wider than the spacing of air molecules. What the route changes is whether that ratio needs explaining. If the large scale belongs to the disturbance and not to the medium, it is the size of a storm, not a second setting of the air.

The sign. As above: if the Ocean's self-interaction has the wrong sign in its present state, nothing arrests collapse and the framework is not viable in its present form. This is a single calculation away from being settled either way. Settling it for the Ocean's present state does not settle it for every time and place. The medium resists compression only in some of its possible states; in others it may not, and the earliest moments of the present beat or distant regions could have been in such a state. The rules do not change from place to place, but where the medium sits within them can.

The preferred frame. Describing the vacuum as a medium gives it a state of rest, and a state of rest is something Einstein spent a career removing from physics. Real superfluids do exactly this, so it is not absurd — but the experimental limits on any such effect are extraordinarily tight, and the framework must show it stays inside them. Among the tightest are the comparisons of matter with antimatter, which agree to parts in a trillion (Section 10, deficiency (5)).

The measured vacuum. What looks like empty space has already been measured doing things. It screens electric charge, it scatters light off light, and in very strong magnetic fields it splits light by polarization, as a crystal does. Standard physics predicts these effects by other means, in some cases to many decimal places. If the Ocean is real, these are its properties, and it must reproduce the same numbers. Agreement would be strong support. Any disagreement would be fatal.

The strength of gravity. Gravity is extraordinarily weak. Between two electrons the electric force is about 4×10^{42} times stronger than their gravity; that ratio is measured, not assumed. If gravity is the medium's response, the medium must produce it. Counting the kinds of wave the Ocean builds gives gravity's measured strength (Section 12.7). Why it is so feeble between two electrons then comes down to the electron's mass, which the framework has yet to explain (Section 13.2, item (5)). The count is also only a first estimate.

What follows is the technical statement of all of this. It is written for a reader comfortable with tensor notation, but the argument it makes is the one above: that the emptiness between things may not be empty, and that a great deal of what we currently call unknown may be the behavior of a single medium that has been sitting in plain view the whole time.

End of Preface · token a70279

Conventions. No independent particulate entities are posited. An excitation of the medium that is localized in the sense of Section 7 is called a *localized excitation*; one that is not is a *coherent excitation*. "Matter" and "the matter sector" denote the localized population of the one medium, never a second substance. Particle-like behavior — the exchange of energy, momentum and charge in whole units at a single point — is treated as a property of how excitations of a quantum field interact, not as evidence of objects (Section 7). The word "particle" appears only when describing other work, and is marked as that work's term. The framework is four-dimensional throughout: indices μ, ν run over one dimension of time and three of space, and the Ocean is a medium in all four. Time is part of the medium's description, not a clock standing outside it.

In the paper's own voice "motion" is not used without a qualifier: it is always flow, circulation or travel, as defined below, except in the rebuild (Section 12) and the re-made state, where "motion" names the travel a pattern carries and the tilt it hands to its surroundings. Every measurement is a measurement of travel, since instruments are themselves patterns; flow is inferred from its effects on travel, never observed directly. The medium is written as the complex field $\Phi = \rho\, e^{i\theta}$ with density $n \equiv \rho^2$ (Section 2.2); to keep θ for the phase, the expansion scalar of the medium's flow is written $\Theta \equiv \nabla_\alpha u^\alpha$. In the paper's own voice c means the local carrying speed and c_0 its measured value. Wherever a speed only converts between units, as in GM/c_0^2 or $\hbar c_0$, it is written c_0: the number a local clock and ruler return, not a fixed speed of nature. "Information" appears only when reporting the physics literature, as in the black-hole information problem; in the paper's own voice the term is *state*. "Dark matter" and "dark energy" appear only as what other work calls them, never as components of this framework: the observations they name stand, and their interpretation here is the swirl of the eddy in which a galaxy sits (Section 6) and the Ocean's own pressure $P(\phi)$ (Section 5). The hottest stage of a beat transition is *the hot, dense phase* or *the transition plasma*; "primeval fireball" is the literature's term. Neither is a ball: it has no surface, center or outside, and is the condition of the whole Ocean at once.

Glossary. The following terms are used with fixed meanings throughout.

The Ocean
The single continuous medium. It has no edges and no outside, and it includes every beat. Nothing in the framework exists apart from it.

Flow
The Ocean's own motion, described by the medium four-velocity u^μ. Flow is not limited by the carrying speed c_s; where it outruns c_s for an observer, there is a horizon (Section 11.1).

Carrying speed
c_s, the speed at which the Ocean carries a disturbance: the one speed of light, of the internal workings of matter and of every signal. It has two readings. On the spot it is always measured with the same value; seen from afar it varies, and that variation is gravity (Section 11.1).

Speed of light
c; not a constant of nature in this framework. The Ocean's local carrying speed, as read by rulers and clocks made of the same Ocean. On the spot it always reads the same, because the instruments scale with the speed they measure; seen from afar it varies, and that variation is gravity. Its value $c_0 = 299\,792\,458\ \mathrm{m\,s^{-1}}$ is exact by definition: since 1983 the metre has been defined by fixing it, so the number is a conversion between units of length and time, not a measurement of nature. No fixed c appears anywhere in the Ocean's law (Section 11.1).

Circulation
The Ocean's motion within a pattern. A pattern can circulate internally while staying in place.

Travel
A pattern's motion as a whole: the location at which the Ocean re-makes it, shifting. Travel is measured by the group velocity of the excitation relative to u^μ, and is limited by the local carrying speed.

State
The complete configuration of the Ocean at the finest scale: its amount and phase, its flow and circulation, and the re-made state. What physics calls information.

Pattern
The form of an excitation: organized structure recognizable at some scale, such as the shape of a whirlpool or the profile of a wave. Patterns can be scrambled; the state is not destroyed.

Carried quantities
What a pattern carries: energy, momentum, circulation and charge. Two patterns of the same form can carry very different amounts. The conserved carried quantities are what survives annihilation, Hawking radiation and a beat transition.

Excitation
The Ocean's departure from its background state at some location: a pattern together with what it carries. Matter and light are excitations; so is a galaxy.

Light
A free wave of the plexus, all of its carrying speed spent in travel, with no lowest energy. Matter is a confined pattern, its carrying speed spent circulating in place: same Ocean, same speed, different shape.

Healing length
The shortest distance over which the Ocean's own amount can bend before it springs back: the width of a dent, near the Planck length, set by the cost of a dent. Not the size of any excitation and not a grain of the medium. Each kind of confined pattern has its own localization length, its reduced Compton length; the electron's is about 386 femtometers (Section 7).

Localized excitation
An excitation localized in the sense of Section 7; what other work calls a particle.

Coherent excitation
An excitation that is not localized.

Circulation sense
The property distinguishing a localized excitation from its antimatter partner: whether the rebuild is building or reclaiming it. It is not spin and is not changed by any rotation (Section 13.2, item (8)).

Horizon
A point of no return for travel: before it, escape is possible; after it, it is not. Not a physical thing and not a surface, and never a boundary of the Ocean, which remains continuous and coherent across it. Horizons are observer-dependent; a region beyond ours has its own horizon, and ours lies within it.

Matter, matter sector
The Ocean's own energy, gathered and circulating in place, made by building. Its building blocks are confined patterns on the plexus's kinds of wave, each held within its own localization length, and everything else that is matter is built of them. To become matter, energy gathered in place must reach at least the lowest energy of its kind, which the dynamo's field sets and every observer measures alike; made from energy alone, matter comes with its antimatter partner, so making it takes at least twice that. The electron's lowest energy is 0.511 MeV; the neutrinos' are far smaller. Each building block is a localized excitation; the matter sector is the whole population of them. Never a second substance.

Rebuild
What the Ocean does at every place and to every pattern at every moment: it builds and reclaims what was just made (Section 12).

Building
The side of the rebuild that makes: three arrows and a left twist. Matter and time are made only by building.

Reclaiming
The side of the rebuild that unmakes: three arrows and a right twist. It makes nothing and takes everything it reaches away; it is antimatter.

Plexus
The Ocean-wide waves the dynamo makes and keeps running, on which every pattern is built and which carry the re-made state. Not a structure, and never local: the whole Ocean is one connected state of them.

Re-made state
What the Ocean rebuilds around a pattern along with it: its energy (which sets clocks), its motion (the tilt) and its pushes (the rulers' stretch and squeeze); ten numbers at each place.

Dynamo
The process by which the Ocean's flow makes and sustains the plexus. The flow supplies it; the cost of a dent tunes it; its leakage is the plexus's own reclaiming, not friction.

The Heartbeat
The whole sequence of beats.

A beat
One expansion and contraction of the Ocean, from one beat transition to the next.

The present beat
The beat we are in, and the only one any observation can reach.

Beat transition
The change of state between one beat and the next. Never a phase change within a beat.

The big bang
The most recent beat transition, as seen from within the present beat. Not the beginning of the Ocean.

Epoch
A phase within a beat, such as the radiation-dominated, matter-dominated or present accelerating epoch. Never a beat.

The present expansion epoch
The hot expanding phase we observe, about 13.8 billion years old by extrapolation. It contains the radiation-dominated, matter-dominated and present accelerating epochs. Distinct from the present beat, whose age is not known (Section 5.2).

The observable universe
The part of the present beat that observation can reach. Neither the whole present beat nor the Ocean.

"Universe" is not used on its own, since what it names is redefined here; it appears only in "the observable universe" and in reporting other work. "Cycle", like "particle", appears only when reporting other work. "Cyclic" in the subtitle names the established category of cosmology to which the Heartbeat belongs.

End of Glossary · token 692b89

1 The Field Equation

The governing equation for the "Ocean" metric is formulated as Einstein geometry coupled to a dynamic scalar vacuum medium. The central revision relative to the original formulation is that the viscous response is treated as a *stress contribution* rather than as a replacement for the geometric Einstein tensor:

$$G_{\mu\nu} + P(\phi)\, g_{\mu\nu} \;=\; \kappa(\phi)\Big[\, T^{\text{matter}}_{\mu\nu} + T^{\phi}_{\mu\nu} + \Pi_{\mu\nu} \,\Big], \qquad \kappa(\phi) = \frac{8\pi G(\phi)}{c(\phi)^4}. \tag{1}$$

Here $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}R g_{\mu\nu}$ remains the geometric Einstein tensor; $P(\phi)$ is the dynamic vacuum pressure; $T^{\phi}_{\mu\nu}$ is the stress-energy of the scalar field; and $\Pi_{\mu\nu}$ is its hydrodynamic dissipative stress, which in the settled Ocean carries nothing (Section 2.3). Section 11.3 makes the split exact: the background's own state, including its pressure $P(\phi)$, belongs to the geometry's side, and $T^{\phi}_{\mu\nu}$ on the right holds only the ripples above the background, $\delta T^{\phi}_{\mu\nu}$, so that nothing is counted twice.

Status of the equation. Equation (1) is retained as the working equation of the framework, not as a fundamental structure with the medium added to it. The medium is primary: the geometry on the left-hand side is the effective description of how the medium's state is experienced by excitations of the medium, and the right-hand side records the medium's own stresses. The equation is kept because Einstein's description is correct to the precision it has been tested; the framework's claim is to explain why it holds (Section 2.5), and deriving it from the dynamics of the medium is the central task that claim creates (Section 14).

End of §1 · token c6c077

2 Tensor Definitions

2.1 Geometric Einstein tensor

The geometric sector retains the standard covariant structure of General Relativity:

$$G_{\mu\nu} = R_{\mu\nu} - \tfrac{1}{2}R g_{\mu\nu}. \tag{2}$$

This separation is essential because the contracted Bianchi identity gives $\nabla^{\mu} G_{\mu\nu} = 0$. Any additional vacuum stresses must therefore satisfy the corresponding total conservation requirement (Section 9).

2.2 Scalar vacuum stress-energy $T^{\phi}_{\mu\nu}$

The vacuum is represented by a dynamical scalar field rather than a purely phenomenological density. Let

$$X \equiv -\tfrac{1}{2}\nabla_{\alpha}\phi\,\nabla^{\alpha}\phi, \tag{3}$$

and let $L(\phi, X)$ be the scalar-field Lagrangian. Its stress-energy tensor is

$$T^{\phi}_{\mu\nu} \;=\; L_{,X}\,\nabla_{\mu}\phi\,\nabla_{\nu}\phi \;+\; L g_{\mu\nu}. \tag{4}$$

This term gives the Ocean a genuine dynamical energy density and pressure. The specific choice of L determines whether the field behaves as a conventional scalar, a k-essence-like medium, or a more specialized superfluid effective theory. Section 7 depends on this choice directly.

The complex field. The Ocean is represented as a complex field with a global $U(1)$ symmetry,

$$\Phi = \rho\, e^{i\theta}, \qquad n \equiv \rho^2, \qquad \theta \;\to\; \theta + \text{const.} \tag{5}$$

This is the baseline representation, and it may be the bosonic sector of a richer order parameter (Section 13.2, item (7)). Two independent requirements select it over a single real scalar: the separation of density, flow and current below, and the gapless phase on which the candidate route for the scale hierarchy depends (Section 13.2, item (5)); frame dragging is carried by the re-made state, not by the phase (Section 11.8). What this field does not give is supplied elsewhere: the re-made state and the plexus carry direction and the spin-$\frac{1}{2}$ order (Section 12.7), and antimatter is the reclaiming side of the rebuild (Section 12.4), so the field's charge does not have to tell matter from antimatter (Section 13.2, items (7)–(9)).

Where expressions in this paper are written for a single real field ϕ, ϕ stands for the state of the medium, which the modulus carries: $\phi \leftrightarrow \rho$. The state-dependent quantities $P(\phi)$, $\kappa(\phi)$, $\xi(\phi)$, $\eta(\phi)$ and $\zeta(\phi)$ are functions of ρ and of the phase rate μ defined below, the quartic potential of Section 9.1 is a potential for ρ, and the beat transitions of the Heartbeat are transitions of ρ. The shift symmetry forbids any dependence on the value of θ itself, so the phase has no potential: its long-wavelength excitation is a gapless Goldstone phonon, as in a laboratory superfluid, and the mass-gap entry of the table in Section 7 vanishes for it. The excitation of the modulus is heavy: it is the amount mode, whose minimum energy is the cost of a dent, near the Planck energy (Section 12.7), far beyond reach. The localized excitations of Section 7 are patterns on the plexus's kinds of wave, not quanta of the amount.

The two components play different parts in the Ocean's state. The amount ρ sits at the minimum v of its potential, held there by the potential rather than by any conserved quantity, so it does not thin as the Ocean expands; it departs from v only in extreme states, at a beat transition or in the core of a collapsed object. The second state variable is the rate at which the phase turns, the Ocean's chemical potential

$$\mu \equiv \sqrt{-\nabla_\alpha\theta\,\nabla^\alpha\theta}, \qquad u_\mu = -\frac{\nabla_\mu\theta}{\mu}, \tag{6}$$

which sets the Ocean's pressure and defines its rest frame: the medium four-velocity u^μ lies along the gradient of the phase. The $U(1)$ symmetry gives a conserved charge, with density proportional to $\rho^2\mu$. As the Ocean expands that charge spreads and μ falls, while ρ stays at v. The phase's turning is each place's clock, read off the Ocean (Section 12.2), and ρ is what resets at the transition. A medium with $\mu = 0$ would have no rest frame at all.

The density n measures how much of the Ocean is present, the gradient of the phase θ sets the flow velocity, and their product is the current: how much of the Ocean is moving. A pattern is the form these quantities take; its carried quantities are integrals over them. A single real field has no phase, so this separation is not available to it. Nothing within a pattern is carried along as a bulk substance; the current, not a moving volume, is the measure of how much of the Ocean is in flow.

The choice brings the framework's field content closer to the complex-field description of Huang and collaborators [1] and to the phonon description of Berezhiani and Khoury [2]; the distinctive claim of Section 10 is unaffected. The complex field does not supply a crossing of $w = -1$ (Section 5.1).

In the present state the Ocean is stiff: its amount ρ barely gives, and its ripples travel at matter's speed limit to about one part in 10^{23} (Section 11.1). Under the minimal law of Section 11.4 that bounds the ratio $2\mu^2/(\lambda v^2)$ below about 10^{-23}; with the dent near the Planck energy it is far smaller still, and frame dragging, carried by the re-made tilt, needs no fast phase rate (Section 11.8). Seen from afar, its phase turns at one rate everywhere, which keeps the Ocean in step; on the spot it turns faster wherever clocks run slow, and each place's turning is that place's clock (Sections 11.6 and 12.2).

2.3 Vacuum viscosity tensor $\Pi_{\mu\nu}$

The viscous response represents the dissipative stress response of the scalar medium to local flow. For a relativistic effective fluid, a first-order constitutive form is

$$\Pi_{\mu\nu} = -2\,\eta(\phi)\,\sigma_{\mu\nu} \;-\; \zeta(\phi)\,\Theta\,h_{\mu\nu}, \tag{7}$$

where $\eta(\phi)$ is the shear viscosity, $\zeta(\phi)$ the bulk viscosity, $\Theta = \nabla_\alpha u^\alpha$ the expansion scalar, $h_{\mu\nu} = g_{\mu\nu} + u_\mu u_\nu$ the spatial projector orthogonal to the medium four-velocity u^μ, and $\sigma_{\mu\nu}$ the shear tensor.

Whether the vacuum's response can reproduce effects conventionally assigned to dark-matter halos does not rest on $\Pi_{\mu\nu}$: in a galaxy that response is the tilt of the eddy in which the galaxy sits (Section 6).

In the settled Ocean. $\Pi_{\mu\nu}$ is dissipative and therefore irreversible; taken as fundamental, it would destroy fine-grained state everywhere, contrary to Section 9.2. In the rebuild nothing is dissipated (Section 12.7): the Ocean's turning is carried by the re-made tilt (Section 11.8), with no quantized vortices to carry a stress, and the leakage the dynamo needs is the plexus's own reclaiming, not friction. In the

settled Ocean $\Pi_{\mu\nu}$ therefore carries nothing, and η and ζ vanish. It is kept in Equation (1) as the place where an effective stress would enter if one is found, for instance at the beat's turn; whether it has any role is open (Section 14).

2.4 Scalar pressure $P(\phi)$

$P(\phi)$ replaces a fixed cosmological constant as the phenomenological vacuum-pressure contribution. It is a function of the state of the underlying scalar field rather than a universal constant:

$$\Lambda\, g_{\mu\nu} \;\longrightarrow\; P(\phi)\, g_{\mu\nu}. \tag{8}$$

The *Heartbeat* hypothesis proposes that the scalar field evolves through the phases of a beat: expansion ("diastole"), driven by the Ocean's pressure, and contraction ("systole"), which is the reclaiming carrying the Ocean once the drive runs out (Section 12.5). A complete model must derive the pressure's evolution from a field potential or equation of state rather than prescribing it by hand.

2.5 Effective coupling $\kappa(\phi)$

The coupling between the vacuum state and ordinary stress-energy is allowed to depend on the local scalar density:

$$\kappa(\phi) = \frac{8\pi G(\phi)}{c(\phi)^4}. \tag{9}$$

$G(\phi)$ represents a field-dependent effective gravitational coupling. The quantity $c(\phi)$ is the Ocean's local carrying speed, the one speed at which every disturbance travels (Section 11.1). Locally it is always measured as c_0, because rods and clocks are patterns of the Ocean and scale with it. Seen from afar it varies, and that variation is gravity (Section 11.1).

The refractive reading. The account of gravity adopted here is refractive. A wave in a medium whose propagation speed varies curves toward the region where it travels more slowly. If concentrated energy lowers the local propagation speed of the Ocean, light curves inward, and so does every localized excitation, each of which is a pattern the medium keeps re-making. Where the medium is slower on one side, the re-making is uneven and the pattern is re-made slightly further toward the slow side. Nothing pulls; trajectories are turned. This is consistent with the absence of any felt force in free fall. In the weak field it reproduces the standard equivalence between gravitational light bending and a refractive index

$$n(r) \;\simeq\; 1 + \frac{2GM}{rc_0^2}. \tag{10}$$

Refraction alone does not reproduce the full weak-field and post-Newtonian behavior — perihelion advance, frame dragging, gravitational radiation — and extending it to do so is the test this reading must pass (Section 14). Three requirements are decisive. Every excitation must respond to the medium's state in exactly the same proportion, since all bodies and light are observed to fall alike to extraordinary precision; here that equality must follow from everything being made of the same medium. The response must slow time and stretch space in equal measure: the measured deflection of starlight by the Sun is twice what slowing alone would give, and the second half comes from the spatial part. And disturbances of the response itself, gravitational waves, must travel at the speed of light, as observed when a neutron-star merger's gravitational and light signals arrived together. Section 11 states the second requirement as a condition on how patterns scale with the carrying speed (Section 11.5) and offers a candidate mechanism for it (Section 11.6). $c(\phi)$ is not a second causal speed: there is one speed (Section 11.1).

End of §2 · token dd4ba8

3 Scalar-Field Dynamics

The field ϕ must have an independent equation of motion. For a Lagrangian $L(\phi, X)$ of the kind in Section 2.2, the covariant form is

$$\nabla_\mu (L_{,X} \nabla^\mu \phi) + L_{,\phi} \;=\; C(\phi, T, \nabla\phi, \ldots)\,, \tag{11}$$

where C represents permitted couplings to curvature, to hydrodynamic variables, or to the structure matter requires. For the simplest choice, $L = X - V_{\text{eff}}(\phi)$ with $V_{\text{eff}}(\phi)$ an effective potential, it reduces to

$$\Box\phi \;=\; \frac{\partial V_{\text{eff}}}{\partial \phi} \;+\; C, \tag{12}$$

with $\Box = \nabla^\mu \nabla_\mu$ the wave operator, taken with the Ocean's own field of rulers and clocks, the geometry of Equation (1), so that no fixed speed enters (Section 11.1). The potential and coupling structure determine the vacuum phases, stability, characteristic sound speed, and possible Heartbeat behavior.

The complex field. With the Ocean written as $\Phi = \rho\, e^{i\theta}$ (Section 2.2), there are two equations, one for the amount and one for the phase. For the minimal law of Section 11.4, Equation (27), and before any coupling, they are

$$\Box\rho = \tfrac{1}{2}\lambda\,\rho\left(\rho^2 - v^2\right) - \rho\,\mu^2, \qquad \nabla_\mu\left(\rho^2\,\nabla^\mu\theta\right) = 0. \tag{13}$$

The first says where the amount settles: uniform and steady, it sits at $\bar{\rho}^2 = v^2 + 2\mu^2/\lambda$, the value used in Section 11.4. The second is the conservation of the Ocean's current, the charge of Section 2.2, whose density is proportional to $\rho^2\mu$. It holds because the phase has no potential, by the shift symmetry; a coupling that acts on the phase must respect that symmetry, or the current is not conserved, and whether the couplings do is the Protection task of Section 14. Where this paper writes a single real field ϕ, it stands for the amount ρ (Section 2.2), and the amount's equation is the first of Equations (13), not Equation (12): the single-field form omits the phase's term $-\rho\,\mu^2$, which moves the settled amount from v by the fraction $a/2$, below about 10^{-23} in the stiff Ocean (Section 11.1), and its normalization differs by factors that the choice of potential absorbs. Matter needs no separate coupling: its patterns are built on the plexus's kinds of wave (Section 12.7), not quanta of the amount, and their exchange with the rest of the Ocean is the Ocean's own interaction (Section 7). What C stands for, and what is not yet specified, is how the Ocean couples to curvature, to its dissipative stress, and to the structure that charge and spin require.

End of §3 · token d37c4e

4 The Ocean Interpretation

The framework interprets the observable universe as a dynamical region of a deeper vacuum medium. The Ocean is not required to begin at the big bang. Instead, the big bang is interpreted as a beat transition: a change of state of the medium, a phase transition, or a rapidly expanding solution of the underlying field equations.

The observable universe is therefore not treated as the ontological boundary of reality. It is the portion of the deeper continuous vacuum field accessible to observation. This distinction is a conceptual premise of the framework and must ultimately be connected to a mathematically well-defined global solution.

End of §4 · token bf999b

5 Cosmological Heartbeat

The proposed *Heartbeat* cosmology associates cosmic expansion and contraction with the evolution of ϕ. A schematic beat is

$$\text{beat transition} \;\rightarrow\; \text{expansion} \;\rightarrow\; \text{the drive runs out}$$
$$\rightarrow\; \text{contraction, the reclaiming} \;\rightarrow\; \text{the turn into the next beat.}$$

The model does not require that each beat create the underlying vacuum field. Instead, each *beat* (see the Glossary) may be a dynamical state of the same deeper medium.

Two kinds of statement are kept apart in what follows. Section 5.1 concerns the present beat, which is the only one any observation can reach. Section 5.2 concerns the sequence of beats and the transitions between them, about which no observation made within a beat can speak directly.

5.1 Within the present beat

Everything known about time is known from within the present beat. Every observer and every instrument is a pattern of the Ocean in this beat, and every observation is of this kind. Here time is the pace at which the Ocean re-makes its patterns, measured by clocks that are themselves such patterns. Durations are meaningful and measurable, but local: clocks run at different rates in different places, as General Relativity describes and as the account of Section 2.5 requires.

Expansion is divergence, not transport. Expansion is an effect of flow whose cause is a change of the Ocean's state. On the largest scales the Ocean's flow is divergent, $\Theta = \nabla_\alpha u^\alpha > 0$ everywhere, with no center and no common direction. A uniform flow carrying every pattern the same way would be unobservable in principle; what is observed is regions moving apart. Regions beyond the Hubble radius recede faster than light. This is flow, not travel, and no pattern exceeds the local speed limit. Some galaxies now receding faster than light remain visible, because the Hubble radius has grown to overtake light they emitted long ago [3]. Bound systems decouple from the divergence: galaxies and planetary systems do not stretch internally.

One root, two effects. Both gravity and expansion begin with a change of the Ocean's state, and differ in its shape. A gradient of state, varying from place to place as near a mass, bends travel: it refracts patterns and, where steep enough, brings outward travel to zero at a horizon, with no net flow of the medium (Section 8.1). A uniform change of state, the same everywhere at once, drives flow: a divergent flow with no center, which is expansion, or a convergent one, which is contraction. In the terms of Section 2.2 the uniform change is a change of μ, with the amount ρ held at v. Literal flow is excluded for a black hole because a local inflow would have to accumulate somewhere; a uniform flow has no somewhere, and the change of state is the same everywhere at once. Nor does literal flow conflict with a constant vacuum energy density: a medium with negative pressure does work on itself as it flows apart, adding energy at the rate that keeps its energy density fixed. The cosmic horizon is therefore a flow horizon and a black-hole horizon a gradient horizon, two mechanisms with one root. Deriving this principle from the dynamics is a task of Section 14.

The pace of the beat. The pace of the beat is not fixed. While matter dominated, its gravity slowed the expansion; distant supernovae show that the expansion is now accelerating [4, 5]. There are further indications that the acceleration is itself evolving: baryon acoustic oscillation measurements from the DESI survey, combined with the microwave background and supernova samples, prefer what the survey calls a dark-energy component that changes over time to a constant one, at roughly 3 to 4σ depending on the supernova sample, short of the level conventionally required for discovery [6]. In this framework such variation is the Ocean's state changing through $P(\phi)$, not a departure from a constant. One caution follows

from the same data. The preferred evolution crosses the boundary $w = -1$, which a single scalar field with a standard kinetic term cannot readily do; the survey's authors note that multiple fields, additional internal degrees of freedom or a non-minimal coupling can evade the difficulty [6]. The complex field of Section 2.2 does not supply this: it is two canonical fields, and canonical fields cannot cross $w = -1$. The crossing needs a phantom component or a non-minimal coupling. If the evolution is confirmed, the Ocean's pressure sector would need such structure, and the candidate already present is the coupling $\kappa(\phi)$.

5.2 Across beats

Nothing in this subsection is observable from within the present beat, and the two subsections should not be read into one another. What the rebuild says happens as a beat ends, from the turn of its expansion through the whole of its contraction, is set out in Section 12.5 and summarized here. What remains open is the turn from the end of one beat into the next.

No clock survives the transition between beats. Time is made in the build, and in the collapse the time a beat made is reclaimed with everything else; the next beat makes its own. The Ocean's state, including ξ, may differ from one beat to the next. Durations therefore cannot be compared from beat to beat, and "faster" or "slower" across beats can only mean that different beats pass through different sequences of states. The framework requires no repetition: each beat may differ in its expansion, its reset and its ξ. Across beats the Heartbeat is counted, not timed. Nor is a beat set to a length: it is event-driven, ending when its triggering condition is met, not when a span of time has passed. Its condition is that no high-to-low differences remain to drive the flow (Section 12.5). A black hole still returning what it holds is unfinished, and the end of the beat waits for it; most if not all black holes are gone before a beat ends, and any that remain are reclaimed with everything else. The return times of Section 11.9 then set a lower bound on the length of a beat.

The contraction itself follows from the rebuild (Section 12.5). When the drive runs out, building is no longer supplied and the reclaiming carries the Ocean, everywhere together, because the spent Ocean reads the same everywhere. Nothing new forms. Everything built is taken away, black holes included, and with it the disorder that was the arrangement of what was built. The energy is not destroyed; it returns to the Ocean, compressed. Disorder therefore does not accumulate from beat to beat, and each beat begins low in disorder. Since the state is never destroyed (Section 9.2), this reset acts on what was built, not on the Ocean's state. What is open is the turn: how the compressed Ocean becomes the next beat, and whether that turn is free of singularities and instabilities. Stiffness keeps the Ocean regular against cascades (Section 9.1) but cannot turn a contraction around: for any material whose pressure does not exceed its energy density, contraction once under way can only speed up (Section 11.9). **[Candidate]** What turns a black hole's collapse around, a loosening of the Ocean's grip under extreme compression, is the candidate for what turns a beat's contraction around: one mechanism for both.

The nearest established precedent is the cyclic model of Steinhardt and Turok [7], in which a scalar field rolling in a potential drives an endless sequence of expansions and contractions, each big bang being a transition from a preceding contraction with finite temperature and density. Two of its answers bear directly on the questions above. Entropy does not accumulate from cycle to cycle because a long phase of accelerated expansion, the present accelerating epoch, dilutes the matter, radiation and black holes of the previous cycle before contraction begins. The Heartbeat reaches the same end by another route: not by dilution but by reclaiming, which takes what was built away. And the nearly scale-invariant density fluctuations are generated during the slow contraction rather than by inflation, with the distinguishing prediction of no long-wavelength primordial gravitational waves. That answer is available to the Heartbeat in principle. The difference lies at the bounce: in that model the transition is motivated by colliding branes along an extra spatial dimension, with the four-dimensional field theory carrying a matching rule supplied by hand. The premise of this framework admits no space outside the Ocean, so the bounce must instead come from the medium itself; the candidate is the loosening of the Ocean's grip above, since stiffness cannot turn a contraction around.

One consequence of this route can be tested from within the present beat. If the Heartbeat seeds the density fluctuations of the present beat during the preceding slow contraction, as in the Steinhardt–Turok model, rather than by inflation, it predicts essentially no long-wavelength primordial gravitational waves. The present upper limit on their amplitude relative to the density fluctuations, from Planck combined with BICEP2/Keck Array data, is $r < 0.056$ [8], and forthcoming microwave-background experiments will test it much further. A detection of primordial gravitational waves would count against a contraction-seeded Heartbeat; continued non-detection would leave it viable.

Two obligations follow from setting inflation aside. *The horizon problem*: the microwave background has the same temperature, to about one part in 10^5, across regions that have not been in contact within the present expansion; the framework's account is that the previous beat's contraction brought them into contact before the transition, and it must be shown that this produces the observed uniformity and its spectrum of departures. The same account must also supply flatness, which inflation otherwise explains: in the cyclic model the long accelerated expansion of each cycle flattens and smooths what follows [7], and the Heartbeat would need its pressure-driven expansion to do the same. *The age of the Ocean*: the 13.8 billion years is the age of the present expansion epoch, extrapolated on the assumption that patterns have kept the same time throughout, not the age of the beat or of the Ocean. Disorder does not accumulate from beat to beat (Section 12.5), so whether the Ocean has an age at all turns on whether the Heartbeat has net expansion from one beat to the next, which is open (Section 14).

End of §5 · token cd0a4a

6 Galactic Dynamics

In the galactic regime, the hypothesis is that the eddy of the Ocean's current in which a galaxy sits turns the travel of its stars, beside the baryons' own pull. The observable circular velocity would then be determined by the two together:

$$v_c^2(r) \;=\; v_b^2(r) \;+\; v_{\phi,\mathrm{eff}}^2(r). \tag{14}$$

Here v_b is the baryonic contribution and $v_{\phi,\mathrm{eff}}$ denotes the effective contribution of the eddy's turning. The goal is not to insert an arbitrary halo profile, but to derive $v_{\phi,\mathrm{eff}}(r)$ from the eddy and the law of Section 12.7.

Rotation is a special case. The relation above applies to rotationally supported disks. Galaxies also carry random travel of their stars, inflow and outflow of gas, vertical oscillation, bars and spiral patterns, and travel through their surroundings. Pressure-supported systems with little net rotation — elliptical galaxies and dwarf spheroidals — follow the same radial acceleration relation as disks [9]. Whatever links the vacuum contribution to the baryons therefore cannot depend on rotation, and the framework must apply to pressure-supported galaxies as well.

One structure, not two. The decomposition above is bookkeeping and should not be read as a statement that a galaxy acts on the medium from outside. In this framework a galaxy is a pattern of patterns, excess localized energy moving through the Ocean, and the eddy it sits in is the state of the Ocean around it; the galaxy and its eddy are one region of the one Ocean, a single solution of its law. This differs from treatments in which galaxies are external sources perturbing the superfluid [1].

Two consequences follow. First, the kiloparsec scale is a property of the individual solution, the galaxy and its eddy, fixed by the current that carries them, and need not be a parameter of the medium; this is the basis of the candidate route of Section 13.2, item (5). Second, if the solution is fixed by its baryonic content — the question posed by the Size task of Section 14 — the baryonic distribution determines $v_{\phi,\mathrm{eff}}(r)$ with no freedom adjusted galaxy by galaxy. The observed radial acceleration relation has that form: across rotationally supported galaxies the total acceleration is fixed by the baryonic acceleration alone, with little or no intrinsic scatter and no sign of a second parameter [10, 9]. That is a target to be derived, not a result.

It also sharpens the exposure recorded in Section 10. A contribution fixed by the baryons is precisely the kind that risks being parallel to the baryonic gravitational field, and a parallel contribution is excluded by Milky Way vertical kinematics [11, 12]. The single-structure reading therefore needs a vacuum contribution whose magnitude is set by the baryons while its direction is not slaved to theirs. A turning across the direction of travel can do that; the swirl below is one form that does.

The vacuum contribution is a tilt. The decomposition $v_c^2 = v_b^2 + v_{\phi,\mathrm{eff}}^2$ reduces the vacuum's response to a single radial quantity. In the framework it is a direction at every place, not a single radial number. A star is not carried by a current: it is rebuilt at every moment, its motion is handed forward, and it is turned only by what it reads where it is (Section 12.7). It reads two things. The load gives the baryons' pull. The tilt of the re-made state, the direction in which time is made turned partly sideways, turns travel across its own direction, as it turns a gyroscope near the spinning Earth (Section 11.8). The galaxy's extra turning is of the second kind, and a turning across travel is not constrained to lie parallel to the baryonic gravitational field. The swirl is a tilt and not a flow of the Ocean's phase: a flow of one connected phase does not turn, except along quantized lines, and the Ocean's turning is carried continuously by the re-made tilt (Section 11.8).

The route proposed here is the eddy in which the galaxy sits. The Ocean's current runs in layers: a wide current, rivers of current within it, and eddies within those, though a galaxy may also sit in a river or in the wide current directly. Stars sit in an eddy as the path of least resistance. A galaxy cannot make its own eddy. Its stars' travel tilts the Ocean around them, as a spinning body's does, but only by about the galaxy's potential in units of c_0^2 times the orbital speed: for the Milky Way, a few times $10^{-4}\,\mathrm{km\,s^{-1}}$, against the $118\,\mathrm{km\,s^{-1}}$ the swirl requires (below). An eddy cannot know where a galaxy's matter is, but it has a limited capacity. The eddy does not depend on the galaxy's travel through its surroundings. That independence is required: the radial acceleration relation shows almost no intrinsic scatter across galaxies traveling at very different speeds relative to the cosmic rest frame, and in colliding clusters the inferred extra mass follows the galaxies rather than the slowed gas. A wake from a galaxy's travel, aligned with its direction of travel rather than with the baryonic field, remains only as a small residual offset, tested separately (Section 14). The tilt does not push stars; it reaches them only through how patterns read the Ocean's state (Section 11.3). Whether its turning reaches the stars as the radial-over-vertical enhancement the parallelism criterion requires is therefore the calculation.

A swirl that turns travel only inward. **[Candidate]** One form does it. In the slow limit of the Ocean's rulers and clocks, with the re-made state tilted by $\mathbf{s}$, a pattern moves as

$$\ddot{\mathbf{x}} = -\nabla\Phi_b + (\mathbf{s}\cdot\nabla)\mathbf{s} - \mathbf{w}\times(\nabla\times\mathbf{s}), \qquad \mathbf{w} = \dot{\mathbf{x}} - \mathbf{s}, \tag{15}$$

with Φ_b the baryons' potential and $\mathbf{w}$ the pattern's travel relative to the Ocean; that patterns read the tilt in this form is assumed, not yet derived. For a flat swirl around the galaxy, $\mathbf{s} = V\,\hat{\boldsymbol{\phi}}$ with V the same at every radius over the range considered and at every height near the disk, the swirl turns the travel of a star on a circular orbit at speed v_c inward by a further $v_c V/R$, beside the baryons' term; a star at rest relative to the galaxy is not turned by it, and vertical travel is untouched. If V changes with radius, the inward turning gains a term $(v_c - V)\,dV/dR$. In a model Milky Way whose stars and gas, taken as one flattened disk of 6×10^{10} solar masses with potential $\Phi_b = -GM/\sqrt{R^2 + (a+\sqrt{z^2+b^2})^2}$, $a = 3.0$ kpc and $b = 0.28$ kpc, alone give $159\,\mathrm{km\,s^{-1}}$ at 8.1 kpc against the observed 229, a swirl of about $118\,\mathrm{km\,s^{-1}}$ supplies the rest, and a star launched from the midplane at $30\,\mathrm{km\,s^{-1}}$ rises to within two per cent of its height in the baryons-only case (583 against 574 pc): travel turned further radially and not vertically, the criterion superfluid dark matter fails [12]. The swirl must not weaken with height near the disk: if it falls off over half a kiloparsec, the same star is thrown ten kiloparsecs out of the disk.

The swirl's shape. The first-order law of Section 12.7 decides between two shapes. The tilt is set by the momentum of patterns. A swirl the same at every height would need a source filling the galaxy:

at the Sun's radius, a momentum about 4×10^4 times what the model's stars and gas carry there. No pattern supplies it. A swirl with no source in the galaxy, put in place from outside by the eddy, is what the law allows. Flat in the disk's plane, it cannot then be the same at every height: it grows slightly, as $V(1+z^2/2R^2)$, so it does not weaken, and its turning gains a part across the disk that draws travel back toward the midplane, by $(v_c-V)Vz/R^2$. In the model Milky Way that adds about 9 per cent to the baryons' vertical pull at 0.5 kpc from the midplane and 15 per cent at 1 kpc, against a radial enhancement of more than a factor of two, and the star launched at $30\,\mathrm{km\,s^{-1}}$ rises to 558 pc against 574. That is a prediction, to be confronted with the full Milky Way analysis [12]. What puts the tilt in place from outside the galaxy is open: the first-order law takes tilt only from the momentum of patterns, so the larger current must do it in a way that law does not yet contain.

How far the swirl turns a star depends on how the star travels through it, which Gaia's velocity data can test; light crossing a galaxy should be deflected differently on the side moving with the swirl and on the side moving against it, which lensing must be checked against; and what sets the swirl's strength from each galaxy's baryons, as the radial acceleration relation requires, given that the eddy cannot know where the galaxy's matter is, is not yet given. The decomposition $v_c^2=v_b^2+v_{\phi,\mathrm{eff}}^2$ treats the baryons as travel and the vacuum contribution as the eddy's tilt; the relation between the two is to be derived.

End of §6 · token 824d10

7 The Localization Threshold

7.1 Why a fixed threshold fails

The simplest way to partition energy between sectors is by a declared inequality:

$$E>0.511\,\mathrm{MeV}\ \Rightarrow\ \text{localized matter sector},\qquad E\le0.511\,\mathrm{MeV}\ \Rightarrow\ \text{scalar-fluid density }\phi.$$

This has three defects, only the first of which is usually noticed.

(1) **It is an unexplained parameter.** The value is imported from the electron rest mass, which the framework does not derive.

(2) **It is not covariant.** E is frame-dependent. A boost changes which sector a given excitation belongs to, so the partition is observer-dependent, which is unacceptable in a theory built on the covariant field equation (1).

(3) **It is discontinuous.** A step function requires an exchange term coupling $T^{\mathrm{matter}}_{\mu\nu}$ and $T^{\phi}_{\mu\nu}$. Combined with the variable coupling $\kappa(\phi)$, this makes the conservation conditions of Section 9 essentially impossible to satisfy.

The framework removes all three by replacing the declared threshold with a property of the waves the Ocean carries: each kind of wave either has a lowest energy or has none.

7.2 The excitation spectrum

For a single kind of wave, perturb its field about a background configuration,

$$\phi(x)=\bar\phi(x)+\delta\phi(x),\tag{16}$$

and let u^μ be the four-velocity of the vacuum medium with spatial projector $h^{\mu\nu}=g^{\mu\nu}+u^\mu u^\nu$. Expanding the Lagrangian $L(\phi,X)$ of Section 2.2 to second order in $\delta\phi$ gives a quadratic action of the form

$$S_2=\frac{1}{2}\int\sqrt{-g}\Big[\mathscr{K}\left(u^\mu\partial_\mu\delta\phi\right)^2-L_{,X}\,h^{\mu\nu}\partial_\mu\delta\phi\,\partial_\nu\delta\phi-V''_{\mathrm{eff}}\,\delta\phi^2-\mathscr{G}\left(h^{\mu\nu}\partial_\mu\partial_\nu\delta\phi\right)^2\Big],\tag{17}$$

with $\mathscr{K}\equiv L_{,X}+2XL_{,XX}$. The final term is a gradient-stiffness (quantum-pressure) operator, the form it takes in a non-relativistic superfluid. With the law written on the Ocean's own rulers and clocks

(Section 11.1) the Ocean is stiff and that term carries no signal at any accessible energy (Section 13.2, item (3)); the threshold rests on the gap, not on that term.

Varying (17) and evaluating in the medium rest frame, a wave either has no gap and travels at the carrying speed, $\omega = c_s k$, or has one:

$$\boxed{\omega^2(k,\phi) \;=\; \Omega_{\text{gap}}^2(\phi) \;+\; c_s^2(\phi)\,k^2} \tag{18}$$

The same form holds for each kind of wave the plexus carries (Section 12.7), with the gap that kind's lowest energy.

The Ocean's own amount and phase. The Ocean's own part is two fields, and the turning of the phase couples them. Perturbing the minimal law, Equation (27), about the settled state, $\rho = \bar{\rho} + \delta\rho$ and $\theta = \mu t + \delta\theta$ with $\bar{\rho}^2 = v^2 + 2\mu^2/\lambda$, gives, with the local carrying speed as the unit,

$$\begin{aligned} S_2^{\Phi} = \int \sqrt{-g}\,\Big[&\big(u^\mu \partial_\mu \delta\rho\big)^2 - h^{\mu\nu}\partial_\mu\delta\rho\,\partial_\nu\delta\rho - \lambda\bar{\rho}^2\,\delta\rho^2 \\ &+ \bar{\rho}^2\Big(\big(u^\mu\partial_\mu\delta\theta\big)^2 - h^{\mu\nu}\partial_\mu\delta\theta\,\partial_\nu\delta\theta\Big) + 4\bar{\rho}\,\mu\,\delta\rho\,u^\mu\partial_\mu\delta\theta\Big]. \end{aligned} \tag{19}$$

Its waves obey $(\omega^2 - k^2)(\omega^2 - k^2 - \lambda\bar{\rho}^2) = 4\mu^2\omega^2$, and there are two branches, each of the form (18) to leading order in a. One has no gap: the sound of the phase, with $c_s^2 = (1+a)/(1+3a)$ and $a = 2\mu^2/(\lambda v^2)$. The other is the amount's ripple, whose gap is the cost of a dent, $\Omega_{\text{gap}}^2 = \lambda v^2 + 6\mu^2$ (Section 12.7). The coupling also leaves the gapless branch a k^4 term of relative size $4a^2$ (Section 13.2, item (3)).

7.3 The three characteristic scales

Every coefficient in (18) is fixed by the law rather than chosen:

Quantity	Definition	Physical meaning
Sound speed	$c_s^2 = L_{,X}/\mathscr{K}$	Carrying speed of the Ocean, the speed of every disturbance (Section 11.1)
Mass gap	$\Omega_{\text{gap}}^2 = V_{\text{eff}}''(\bar{\phi})/\mathscr{K}$	Curvature of the effective potential at the local state
Mass	$m_\star = \hbar\Omega_{\text{gap}}/c_s^2$	The lowest energy of the kind, $m_\star c_s^2$, expressed as a mass

From these follows the scale that carries the section, the *localization length* of a kind:

$$\xi(\phi) \;=\; \frac{\hbar}{m_\star(\phi)\,c_s(\phi)}, \tag{20}$$

its reduced Compton length: the shortest distance over which a wave of that kind can be held before its lowest energy dominates. For the Ocean's own amount it is the *healing length*, the width of a dent, which with the dent near the Planck energy (Section 12.7) lies near the Planck length. For the electron's kind it is 386 femtometers. It is a threshold, not a granularity and not a size (Section 13.2, item (7)).

7.4 The localization criterion

The partition is by kind of wave. A kind with a gap can be held in place: its patterns are confined, localized excitations, and contribute to $T_{\mu\nu}^{\text{matter}}$. A kind with no gap cannot be held; its waves are coherent excitations and contribute to $T_{\mu\nu}^{\phi}$, or, for light, to the plexus. The partition is not by wavelength within a single branch: the k^4 part of the spectrum, where localized excitations might otherwise sit, carries no signal in the stiff Ocean. The dimensionless quantity

$$\mathscr{Q}(\phi,k) \;\equiv\; \xi^2(\phi)\,h^{\mu\nu}k_\mu k_\nu, \tag{21}$$

constructed covariantly from an excitation's four-momentum and the medium's own state, then says how a confined pattern travels: for $\mathscr{Q} \ll 1$ it is slow, $\hbar\omega \approx m_\star c_s^2 + \hbar^2 k^2/2m_\star$, a pattern at rest or traveling slowly; for $\mathscr{Q} \gg 1$ it travels near the carrying speed, $\omega \approx c_s k$. Both are matter. The threshold energy of a kind is its lowest energy,

$$E_c(\phi) = \hbar\,\Omega_{\rm gap}(\phi) = m_\star(\phi)\,c_s^2(\phi), \tag{22}$$

the rest energy of one of its patterns, the same for every observer.

The partition is by kind, not switched. $T^{\rm matter}_{\mu\nu}$ is the energy–momentum of the confined patterns and $T^{\phi}_{\mu\nu}$ that of the coherent waves. Since both are waves of the one Ocean, their sum is conserved identically, and exchange between them is the Ocean's own interaction, as when a pair of confined patterns is made from energy of at least twice their lowest energy (Section 12.4). No exchange term is required, and the consistency conditions of Section 9 are not obstructed by the threshold.

7.5 The electron's 0.511 MeV

The electron is a confined pattern on its own kind of plexus wave, and 0.511 MeV is that kind's lowest energy. It is set by the strength of the field the dynamo sustains, as a wave in a plasma gains a lowest frequency from the plasma it travels in: the dynamo keeps a background running through the whole Ocean, and the electron's kind reads it (Section 12.7). Its localization length,

$$\xi_e = \frac{\hbar c_0}{0.511\,\text{MeV}} \approx 386 \text{ femtometers}, \tag{23}$$

is the electron's own, its reduced Compton wavelength. It is not the Ocean's healing length, which lies nearly 10^{23} times below it.

The match cannot be read the other way. If 0.511 MeV were the cost of a dent in the Ocean, the electron would be "the lightest pattern the Ocean can hold", and everything lighter, neutrinos included, would be necessarily coherent. Three things rule that out. Neutrinos have mass, measured from their oscillations, so they are confined patterns of kinds with small gaps, lighter than the electron. The strength of gravity puts the dent near the Planck energy (Equation (37)). And the electron's mass is a property of the dynamo's field strength, not of the Ocean's grip. The 0.511 MeV is a readout of the present beat, not an input.

7.6 Consequences

The threshold is local and dynamical. $E_c = E_c(\phi(x))$ for each kind, so regions in different states have different thresholds. This connects the present section to the Heartbeat cosmology: a change of the Ocean's state is also a change in what it is *able to hold* as matter. At the end of a beat the dynamo can no longer hold the electron's kind, and the electron can no longer form (Section 12.5). Within a beat such departures are confined to extreme states, such as a beat transition or the core of a collapsed object; elsewhere the threshold holds the value fixed for the beat, as the next consequence states.

The electron's mass is a constant of the beat. The flow powers the dynamo, and the Ocean's own scale, the cost of a dent, tunes it, as a resonator sets the pitch of a steadily driven string. So the electron's mass is set by the tuning, not by how hard the flow drives, and is steady wherever the Ocean is in its ordinary state; another beat may read differently. This agrees with the limits on drift of the proton-to-electron mass ratio, parts per million over billions of years from quasar absorption spectra [13] and a few parts in 10^{17} per year from atomic clocks [14]. The question is then no longer how fast m_e drifts but how the dent tunes the dynamo. Under the one-speed principle (Section 11.1) a uniform change of the carrying speed, across a beat or within one, would be invisible on its own, since every pattern scales with it; what these limits test is whether any pattern fails to scale exactly.

So is the strength of gravity. G is set by the plexus's waves and the dent (Equation (37)), both of the tuning, and so is the Planck length assembled from it, so the ratio α_G of (39) is fixed for the beat as well

(Section 13.2, item (5)). Lunar laser ranging over fifty years finds $\dot{G}/G = (-5.0 \pm 9.6) \times 10^{-15}$ per year [15]; if that rate has been steady, G has changed by less than four parts in 10^4 over the present expansion epoch.

It introduces a preferred frame. The dispersion relation (18) is stated in the rest frame of u^μ. This is honest (real superfluids do exactly this), but it is a departure from exact local Lorentz invariance and must be reconciled with precision tests.

Particle-like behavior does not require smallness. An excitation appears particle-like because it exchanges energy, momentum and charge in whole units and at a single point. Both properties follow from the medium being a quantum field with local couplings, independently of the localization scale. The localization length is a threshold, not a radius, and above it the term used here is *excitation* or *coherent excitation*, whose size is set by circumstance.

End of §7 · token f72705

8 Limiting Regimes

Regime	Required behavior	Physical objective
GR / Solar System	$\phi \to \phi_0$; $G(\phi) \to G_0$; $c(\phi) \to c_0$; vacuum stresses negligible	Recover tested General Relativity
Galactic	The eddy's tilt turns the travel of stars (Section 6)	Explain rotation curves without dark matter in the particle form its proponents posit
Cosmological	$P(\phi)$ and $T^{\phi}_{\mu\nu}$ contribute on large scales	Generate the expansion history, including the acceleration other work attributes to dark energy
Beat transition / Heartbeat	ϕ evolves between dynamically distinct states; $\xi(\phi)$ shifts	The contraction is the reclaiming once the drive runs out (Section 12.5); provide a mechanism for the turn into the next beat and for changes in the localization floor
High-energy	Energy reaches twice a kind's lowest energy, so pairs of confined patterns form	Test the derived localization threshold of Section 7
Strong field / black hole	Point of no return at $2GM/c_0^2$; collapse turned around by a loosening of the Ocean's grip, not by stiffness	Match horizon-scale imaging and merger ringdowns; calculate the turnaround and the return

8.1 Black holes

In the refractive reading of Section 2.5, a black hole is a region where waves are slowed enough that an outward-directed wave is turned back. Laboratory analogs make this concrete. A fluid flowing faster than its own sound speed forms a horizon that sound cannot cross, and such a horizon has been made in an atomic Bose–Einstein condensate, where the spontaneous analog of Hawking radiation was observed together with the entanglement of its pairs [16]. In that system the shortest length scale is the healing length $\xi = \hbar/mc_s$, the same construction as Section 7. General Relativity itself admits an exact description of a stationary black hole as space flowing inward through a flat background, reaching the speed of light at the horizon [17]. For a rotating black hole that description has the river twist rather than swirl. A

draining vortex in water is nonetheless an analog of a rotating black hole, and rotational superradiance — waves extracting energy from the rotation — has been observed in one [18].

The horizon is refractive. The medium does not pour radially inward. A literal inflow faster than the wave speed would, by continuity $\nabla_\mu(n\,u^\mu) = 0$, make the Ocean accumulate inside the horizon. (Near the horizon the Ocean does keep in step by a gentle inflow, Section 11.9; that is not this.) The inflow of the river description is instead an effective geometry, produced by the steep radial change in the Ocean's state and in $c(\phi)$ near the mass, and the horizon is where outward travel falls to zero. Laboratory horizons, which are made by literal flow, therefore share the black hole's mechanism without being the same kind of system. This is the gradient half of Section 5.1.

No accumulation. A black hole is not gathered substance. On the premise that matter is energy circulating in place, a black hole is a large organized circulation, and its growth is the growth of that organization. Nothing piles up. Mass is the total energy of the circulation; the black hole's spin in the sense of General Relativity is the net circulation after many contributions on different axes cancel. A non-rotating black hole is not still; its circulations sum to zero.

Stripping is the no-hair theorem. Infalling patterns are scrambled until only the black hole's mass, net rotation and charge remain visible from outside, as the no-hair theorem requires. Frame dragging forces nearby patterns into co-rotation. Circulation sense is among what is stripped: a black hole formed from antimatter is indistinguishable from one formed from matter, so the black hole is the one setting in which the matter/antimatter distinction is undone without annihilation. The distinction between what is stripped and what survives has a standard reason. Electric charge is coupled to a long-range gauge field, so it can be measured by a surface integral far from the black hole, and survives as hair. Lepton and baryon number are global symmetries with no long-range field, and are stripped; circulation sense is stripped because it is whether the pattern is being built or reclaimed, and the pattern is not kept (Section 13.2, item (8)). The same reasoning sets an obligation: electric charge requires a long-range gauge field, which in the Ocean must be the plexus that carries light (Section 12.7); that the plexus supplies charge's field is not derived. Stripping scrambles patterns; it does not destroy the state (Section 9.2).

Currents near a rotating black hole. A current that runs across a pattern's path faster than the pattern can travel does not prevent its passage; it carries it sideways and prevents it from holding still. That is the ergosphere of a rotating black hole, lying outside the horizon, where nothing can remain at rest relative to distant regions but escape is still possible, and where rotational energy is extracted. A horizon, by contrast, stops travel in one direction altogether, and encloses what lies inside it.

Hawking radiation is the escape route. The black hole's energy, and with it the state, leaves as Hawking radiation. At the horizon the steep gradient of the Ocean's state, which brings outward travel to zero, stretches fluctuations of the medium into real outgoing ripples, the energy drawn from the black hole. A fluid flowing faster than its own speed of sound has a horizon of the same form, and is predicted to emit thermal sound by the same argument [19]; a laboratory analog has since observed the effect [16]; in the Ocean the gradient of state does the same work. The ripples carry energy and momentum together. For a rotating black hole they carry angular momentum several times faster than energy, so it spins down to a nearly non-rotating state before it has given up most of its mass [20]; rotational energy can also be drawn off by superradiance, the amplification of waves by a body that turns faster than the wave pattern sweeps round it [21, 18]. For a non-rotating black hole the radiation is isotropic and carries energy with no net momentum.

The temperature is set by how steeply outward travel falls to zero at the horizon: the surface gravity in General Relativity, the velocity gradient in laboratory analogs, and the gradient of the Ocean's state here. The horizon is defined by that condition, so it does not change in character as the black hole shrinks;

it moves inward. A shrinking black hole therefore becomes tighter, not faster: its gradient steepens and it radiates hotter and faster, giving the runaway of the final stage. Rotation lowers the temperature, to zero at maximal spin. The radiation emerges from a region comparable in size to the black hole, its typical wavelength being about eighty times the horizon radius [22], not from a glowing surface. What an observer detects depends on their state of travel: a pattern falling freely detects nothing at the horizon, while one held stationary near it detects a thermal bath. In the Ocean's terms, what a pattern detects depends on its travel relative to the medium.

It will not be measured astrophysically. For a solar-mass black hole the Hawking temperature is about 60 nK and the mass loss about $10^{-45}\,\mathrm{kg\,s^{-1}}$, far below what it absorbs from the microwave background; every astrophysical black hole currently gains mass. Evaporation times run from about 10^{67} years upward: Page's original calculation, which treated neutrinos as massless, gives 2×10^{66} years for a solar mass [23], and without massless neutrinos the figure is several times longer. Searches for final-stage bursts from small primordial black holes have found none and constrain their abundance. The mechanism is tested through laboratory analogs, which share the black hole's mechanism, a horizon at which outward travel stops, although their horizon is made by literal flow and the Ocean's by a gradient of state.

Black holes recycle; they do not drain. A black hole is not a passage to anywhere else; there is nowhere else. It returns energy to the Ocean and keeps no structure. Most of the return happens outside the horizon and early. Accretion converts about 6 per cent of infalling rest mass into radiation for a non-rotating black hole and up to about 40 per cent for a rapidly rotating one, against 0.7 per cent for nuclear fusion. A magnetic field threading a spinning hole can draw off its rotational energy electromagnetically and beam it out in two opposite directions [24]; the largest jets so powered reach millions of light-years, and the outflow heats and redistributes the surrounding gas, regulating star formation. The Fermi bubbles above and below the Milky Way's center were most likely created by a large release of energy at the center within the last ten million years or so, either an accretion episode of the central black hole or a nuclear starburst, the discoverers finding difficulties with both [25]. Mergers return a few per cent of the mass as gravitational waves, and Hawking radiation returns the rest. What crosses the horizon returns only as energy: no atom, electron or quark comes back out as itself. Any localized excitations in the output are made afresh by the Ocean, as electron–positron pairs from black holes below about 10^{14} kg, and hadrons in the final stage [23]. Jets contain matter, but matter from the accretion flow that never crossed. Energy is recycled; structure is not, though the state is preserved (Section 9.2).

Three questions remain.

The core. With $\lambda > 0$, collapse might seem able to stop at a core of finite density set by the healing length. It cannot: stiffness cannot reverse collapse inside a point of no return, whatever the density scale (Section 11.9). A loosening of the Ocean's grip can, and the flow then passes through and returns (Section 11.9). This is the question Section 5 asks of a contraction phase, posed for a single object.

Two horizons. The authors of the river model stress that its flat background has no observable meaning [17]. A medium could have a horizon of its own that need not coincide with the geometric one. With the law written on the Ocean's own rulers and clocks (Section 11.1) there is one field of rulers and clocks for every pattern, so the two coincide: the geometry is the medium's own state, which bears directly on the structural question of Section 10. The k^4 term would allow wavelength-dependent leakage, but it carries no signal in the stiff Ocean (Section 13.2, item (3)).

Observation. Horizon-scale imaging and gravitational-wave ringdowns agree with General Relativity, so any departure must lie inside the horizon or below current sensitivity.

Two further obligations are recorded in Section 14. A black hole's entropy, the size of the state it can hold, is proportional to its horizon's area rather than its volume, which a medium holding state throughout its volume must explain. And a black hole traveling relative to the medium, as a recoiling merger remnant does, should show asymmetric infall and possibly drag; General Relativity does not distinguish that from being carried by the flow, but a medium with a rest frame does.

8.2 A traveler falls in

This subsection follows a single pattern, a human traveler in an ideal protective suit, into a black hole, and marks at each stage what is established, what is the framework's reading, and what is open. It is a living section, reviewed at each release, and its standings change as the items they depend on change. **[Established]** means General Relativity and observation; **[Framework]** means the Ocean's reading, consistent with what is established; **[Open]** means it depends on unresolved items.

Choice of black hole. **[Established]** Tidal stretching acts on every part of a body at once, so no suit resists it. Near a black hole of ten solar masses the head-to-foot difference in acceleration reaches about ten times Earth's gravity some 3800 km out, while the horizon lies at about 30 km: the traveler is torn apart long before arrival. The case followed here is the black hole at the center of the Milky Way, Sagittarius A*, of about 4.3 million solar masses and horizon radius about 12.7 million km, where the head-to-foot difference at the horizon is about a ten-thousandth of Earth's gravity. **[Framework]** What tears a traveler apart is not the swiftness of the Ocean's current but how steeply it changes across a body (Section 11.2). Near a stellar-mass black hole the current steepens over a few kilometers; at Sagittarius A* it is nearly uniform across a body at the horizon, and the traveler crosses intact, carried by a swift current that is barely felt.

Approach. **[Established]** Outside the horizon the danger is the accretion flow, gas at millions of degrees radiating X-rays, which the suit is assumed to withstand. Around a rotating black hole frame dragging turns the traveler with it, and inside the ergosphere no pattern can remain at rest relative to distant regions. At 1.5 horizon radii light can orbit, and that orbit sets the size of the black hole's shadow; for Sagittarius A* the observed shadow lies within about ten per cent of General Relativity's prediction [26]. **[Framework]** The traveler is a pattern traveling through a region where the Ocean's rulers and clocks are increasingly squeezed and outward travel, seen from afar, increasingly slows. Around a non-spinning mass the Ocean's geometry is General Relativity's (Section 11.7), so light orbits at the same place and the shadow has the same size. Outside the horizon outward travel is still possible, and so is escape.

The horizon. **[Established]** The traveler notices nothing at the horizon: no wall, flash or jolt, as the equivalence principle requires. Distant observers never see the crossing: the traveler's light reddens, slows and fades, and the last image dims at the edge. The traveler's own clock runs normally. **[Framework]** The horizon is a point of no return: before it, escape is possible; after it, it is not. It is not a physical thing, not a surface, and not an edge of the Ocean, which is whole and connected across it. The same point holds for every pattern, since every pattern has the same carrying speed. In the Ocean's geometry the current through time is brought to zero, seen from afar, at $2GM/c_0^2$, General Relativity's radius (Section 11.7). Near it the Ocean keeps in step with itself by flowing gently inward, and the traveler, falling with it, meets nothing there: the Ocean's own reason for the established line above (Section 11.9). Falling freely, the traveler detects no Hawking radiation. Past the horizon every direction of travel leads inward.

Inside. **[Established]** The longest possible time from horizon to center is $\pi GM/c_0^3$: about 66 seconds for Sagittarius A*, about 28 hours for M87* at 6.5 billion solar masses. Firing rockets shortens it. The outside sky remains visible, distorted; a falling traveler does not see the whole future of the observable universe pass. Tides grow during the fall and become lethal about 280 000 km from the center of Sagittarius A*, stretching bodies, then molecules, then atoms. **[Framework]** Inside, forward in time points inward, and the Ocean's own phase flows inward with the traveler. A pattern's time is its rhythm; as the traveler's patterns are stripped, their clocks go with them.

The core. **[Framework]** Nothing is gathered at the center. Recycling, for the Ocean, is flow-through: the inward flow does not stop or collect at the center, but passes through and returns to the Ocean, later. Time is recycled with it: inward on the way in, outward again on the way out. **[Candidate]** General

Relativity ends the fall at a singularity, where the theory fails. The Ocean's resistance to compression ($\lambda > 0$, Section 9.1) cannot turn the inward flow around once it is inside the point of no return; a loosening of the Ocean's grip under extreme compression can (Section 11.9), stopping the fall short of a point and turning it around about half a minute after the crossing, on the traveler's own clock, for Sagittarius A*. **[Open]** The return must be slow enough, seen from afar, to fit everything observed of black holes. How slow depends on what the Ocean is made of; written as a function of that fine scale it is slow enough, but it is not yet derived. Separately, each kind's lowest energy is set by the dynamo's field where the pattern is (Section 7): if the core's state weakens that field, the traveler's confined patterns are no longer held there and dissolve into coherent waves rather than being crushed. Whether compression strengthens or weakens it is not known.

In the rebuild's terms. **[Framework]** At every moment the traveler is reclaimed and rebuilt (Section 12). Falling, the traveler's near side is made more slowly than its far side, which is the tide. The share of the Ocean's capacity for rebuilding in use rises as the horizon nears and reaches one there: seen from afar, no more time is made at the horizon, which is why the last image dims and never crosses. On the traveler's own clock nothing changes, because the traveler's rebuild and its clock are the same making of time. The energy the traveler borrows from the surrounding Ocean for each rebuild grows with the fall and is returned as the traveler moves on, until, past the horizon, the rebuild can only carry the traveler inward.

What becomes of the traveler. **[Framework]** The traveler's patterns are scrambled until nothing distinguishes them from the black hole. The carried quantities, energy, momentum and charge, add to its mass, net rotation and charge. The state is preserved (Section 9.2) and returns to the Ocean. **[Open]** By which route: as Hawking radiation, over about 10^{87} years for Sagittarius A*; through the flow-through of the core, of order 10^{22} years or longer seen from afar if the return is a random walk (Section 11.9), long before Hawking radiation would finish; or at a beat transition, if one comes sooner (Section 5.2). Within about a minute the traveler is gone as a pattern; in principle the traveler is never gone as a state.

End of §8 · token e2cf88

9 Conservation and Consistency Conditions

Because $\nabla^\mu G_{\mu\nu} = 0$, the complete right-hand side of (1) together with the pressure sector must satisfy a compatible conservation law. With variable $\kappa(\phi)$, the simple condition $\nabla^\mu T_{\mu\nu} = 0$ is generally not sufficient by itself. The scalar equation (11) and the matter coupling must be chosen so that the full theory is diffeomorphism-consistent.

Section 7 assists here rather than obstructing: because $T^{\text{matter}}_{\mu\nu}$ and $T^{\phi}_{\mu\nu}$ are kinds of wave of the one Ocean rather than two sectors joined by a step function, no discontinuous exchange term is introduced at the threshold.

9.1 Global regularity and the defocusing condition

A separate question is whether the medium remains smooth for all time. The framework cannot appeal to dissipation alone to secure this. Viscosity is conventionally expected to drain an energy cascade before it concentrates without bound, but that expectation is not a theorem: finite-time blowup constructions for the forced three-dimensional Navier–Stokes equations were announced in September 2026 and, while they remain subject to independent verification and do not transfer directly to a relativistic scalar medium, they remove the intuition that a viscous continuum is automatically globally regular. The shear and bulk viscosities $\eta(\phi)$ and $\zeta(\phi)$ of Section 2.3 therefore cannot be assumed to guarantee what they were implicitly relied upon to guarantee.

The mechanism available to this framework is not dissipation but the ultraviolet cutoff supplied by Section 7. The Ocean's amount cannot be bent over distances shorter than $\xi(\phi)$ without paying the cost of a dent: the gradient energy of the amount mode, together with its gap (Section 7), dominates the cascade at $k \sim 1/\xi(\phi)$ and arrests concentration at the healing length. This is the mechanism by which a laboratory superfluid remains regular where the corresponding classical fluid description does not.

That arrest is conditional on the sign of the self-interaction. Writing the relevant quartic term of the effective potential as

$$V_{\text{eff}}(\phi) \supset \frac{\lambda}{4}\left(\phi^2 - v^2\right)^2, \tag{24}$$

the medium must be *defocusing*, $\lambda > 0$: compression must raise the energy and be resisted. The defocusing case is energy-coercive and the cutoff at ξ holds. In the focusing case $\lambda < 0$ the self-interaction rewards further compression, the cutoff does not arrest the cascade, and concentration accelerates instead. The sign is therefore a stability requirement of the same standing as $L_{,X} > 0$ and $\mathscr{K} > 0$, not a modeling preference. With the complex field as baseline the potential belongs to the modulus ρ (Section 2.2), and the argument is unchanged.

Regularity is a property of the state, not a guarantee. The sign of λ does not settle the matter for all times and places, because the stiffness that resists compression is the response of the medium *around the state it occupies*. With the Lagrangian held fixed, that response differs between states. Even the quartic form above, with $\lambda > 0$, has curvature

$$V''_{\text{eff}}(\phi) = \lambda\left(3\phi^2 - v^2\right), \tag{25}$$

which is positive about the present vacuum $\phi = v$ and negative for $|\phi| < v/\sqrt{3}$. A medium driven into that region — by a beat transition, a contraction, or the core of a collapsing object — is no longer protected by the mechanism above. For this form the instability drives the medium back toward the stable state rather than into collapse; whether the full potential admits genuinely focusing states is open. The law is fixed; what varies is where the medium sits within it. No free function is added.

Four constraints follow. (i) Regions in different states are separated by boundaries that carry energy; walls of this kind within the observable universe are tightly excluded, so differing regions must lie beyond the horizon, in the past, or be separated without such walls. (ii) The microwave background is uniform to about one part in 10^5, which bounds how far the state could have varied across the observable universe at that epoch. (iii) A region that loses its stiffness collapses quickly and leaves relics, compact objects or primordial black holes, whose abundance is bounded. (iv) Where smoothness fails, the framework must state what follows — a bounce, a phase transition, or a genuine singularity at which it ceases to predict — and must tie each failure to a specified state of the medium, so that the possibility cannot be invoked case by case.

This section's argument rests on the framework's own structure. It does not depend on the blowup constructions cited above, which await independent verification.

A successful formulation should explicitly establish:

(1) covariance;

(2) total energy-momentum conservation;

(3) a well-posed scalar-field equation;

(4) absence of pathological ghosts or gradient instabilities;

(5) a stable low-density / weak-field limit;

(6) compatibility with precision tests of gravity and light propagation;

(7) a defocusing self-interaction about the present vacuum state, so that the healing-length cutoff of Section 7 arrests rather than accelerates concentration there, together with an account of the states in which it does not (Section 9.1);

(8) reversible underlying dynamics: nothing is dissipated, and $\Pi_{\mu\nu}$ carries nothing in the settled Ocean (Sections 2.3 and 9.2).

9.2 Nothing is destroyed

The framework adopts a principle of preservation: the state of the Ocean is never destroyed. Formally this is reversibility, the requirement that each state determine its predecessor uniquely, which in quantum mechanics is unitarity. It does not require anything to run backward, and since time in this framework is the pace at which the Ocean re-makes its patterns (Section 5.1), "reversing time" is not an available operation. It is distinct from energy conservation: a process could conserve energy and still lose the state, which was the original concern about thermal Hawking radiation. The framework commits to both. Patterns are scrambled, coarse-grained disorder rises, and the arrow of time is that rise from a low-disorder start of the present beat; the fine-grained state is untouched. Recovery of a scrambled pattern need not mean running anything backward. In the spin echo, nuclear spins that have drifted out of step in an uneven magnetic field come back into step after a single pulse, each still turning forward at its own rate; only what has been truly randomized, by diffusion and by interactions between the spins, fails to return [27].

9.3 Where a wave's energy goes

The Ocean has no shore in any direction, so a wave crossing it is not stopped by a boundary. Its energy has five possible fates.

(1) **Spreading.** The wave's front grows and its energy is shared over it, with no absorption at all. This is the main fate of gravitational waves, to which the observable universe is nearly transparent.

(2) **Redshift.** Travel across an expanding region stretches the wave. The microwave background has lost over 99.9 per cent of its energy since it was released. General Relativity has no global energy conservation in an expanding spacetime and needs no account of the loss; this framework commits to preservation and must give one. Taking the divergence of equation (1) with $\nabla^{\mu} G_{\mu\nu} = 0$ fixes where the answer must appear: any transfer between radiation and the $P(\phi)$ sector is a non-zero divergence on the right-hand side, which would change how radiation density falls with expansion, and the microwave background and light-element abundances constrain that tightly. Until there is a calculation, redshift is the one fate of a wave the framework cannot follow.

(3) **Resonant absorption.** A pattern takes in energy only at its own frequencies, as an atom absorbs light only at its spectral lines, and passes the rest.

(4) **Cascade.** A large wave breaks into smaller ones and those into smaller still, until the energy reaches the healing length and is shed into the medium's background of short-wavelength excitations, as superfluid turbulence ends at the healing length.

(5) **Horizons.** A black hole absorbs any wave it intercepts, strips it, and returns its energy much later as Hawking radiation (Section 8.1).

The Ocean therefore has no shore in extent but two in scale: the healing length below, where cascades end, and horizons above, which take in whatever reaches them.

End of §9 · token e0ec99

10 Relation to Existing Work

The proposal that the vacuum is a superfluid, and that dark matter and dark energy are two behaviors of it rather than two substances, is not new. This section states what is already established, so that the framework's actual claim can be isolated from the parts of it that are inherited.

10.1 Two families of superfluid-vacuum programs

Existing work divides on whether spacetime geometry is fundamental.

Emergent gravity. Volovik's program [28] treats the quantum liquid as prior to spacetime: the metric, Lorentz invariance and gauge structure appear as low-energy collective behavior of the condensate. Zloshchastiev's logarithmic superfluid vacuum theory [29] derives an induced gravitational potential from the vacuum wavefunction and concludes that dark matter and dark energy are manifestations of the same object and therefore cannot be independent. That conclusion is the one reached here. The mechanism is not: Zloshchastiev distinguishes his approach explicitly from models that work by modifying the stress-energy tensor of the Einstein equations.

Volovik's program also supplies an argument this framework can use. Inhabitants of a superfluid, built from its own low-energy excitations, would perceive an effectively relativistic world and could not detect the underlying medium directly [28]. The same argument would account for the precision of Lorentz invariance at accessible energies here. It explains what cannot be observed; it does not exempt the framework from effects that are observed (Section 10, deficiency (4)).

Modified source. Huang and collaborators [1] treat a complex vacuum scalar field as a cosmic superfluid within standard General Relativity, identifying its energy density with dark energy and departures from vacuum density with dark matter. That work also treats galaxies as external sources perturbing the superfluid, and obtains the galactic halo together with the creation of vortices by galactic rotation — the closest existing precedent for the galactic sector of Section 6. Berezhiani and Khoury [2] construct an effective field theory in which dark matter is a superfluid with a galaxy-scale coherence length, described by phonons rather than, in the standard model's terms, collisionless particles, recovering MOND phenomenology on galactic scales. That model has since been disfavored by Milky Way data [12]: it over-predicts vertical accelerations of disk stars even while reproducing the rotation curve. Viscous cosmology [30] models the dark sector as a single fluid whose bulk viscosity produces a negative effective pressure and thereby mimics dark energy. The logotropic models of Chavanis [31] belong to the same family, yielding a density profile that falls as r^{-1} and offering an alternative to MOND that accounts for the Tully–Fisher relation.

On the cosmological side, the closest structural precedent is the cyclic model of Steinhardt and Turok [7] (Section 5). It, too, is a four-dimensional scalar field theory in which one field accounts for dark energy, the reversal from expansion to contraction, and the transition between cycles, and it couples that field to matter through a field-dependent factor. Its authors note that this coupling is a modification of General Relativity and produces a fifth force violating the equivalence principle unless the coupling's variation is very small at the present field value. The same constraint applies to $\kappa(\phi)$ here, and it is the first of the requirements of Section 2.5: every excitation must respond to the medium in exactly the same proportion.

In its account of what is fundamental this framework belongs to the first family (Section 1): the medium is primary and geometry is its effective description. Its working equation remains that of the second family, Equation (1), with the medium's stresses on the right-hand side.

10.2 What is inherited and what is not

Inherited. The following are taken from earlier work, in substance if not in form:

- the identification of the vacuum as a scalar condensate;
- the proposal that the dark sector is one medium rather than two substances;
- the appearance of a coherence length in the galactic regime;
- the strength of gravity as the stiffness of a sea of vacuum waves against bending, set by how many kinds of wave there are and by their shortest wavelength (Sakharov's induced gravity, as given in [32]);
- the reading of antimatter as a pattern whose own time runs counter to ours (Stueckelberg's, as used in [33]);
- the fixing of charges in thirds by balance conditions on each family.

Inherited also, and more directly than the preceding list suggests: the generation of galactic structure by vorticity in a vacuum superfluid. Huang and collaborators [1] obtain vortices from galactic rotation within a nonlinear Klein–Gordon description in which galaxies act as external sources. Section 6 reaches for the same phenomenon through different machinery — the tilt of the re-made state in the eddy where a galaxy sits, which the galaxy cannot make itself, rather than vortices sourced by the galaxy. Whether these are distinct physical mechanisms or one mechanism in two formalisms is not settled here, and a credible version of this framework must say which.

The premise that ordinary matter is itself an excitation of the vacuum medium is also not new, and it is worth being precise about who holds it. Huang [1] and Berezhiani and Khoury [2] are two-species: the condensate constitutes the dark sector while baryonic matter is a separate species coupled to it. Superfluid vacuum theory is not. Its stated paradigm is a background quantum liquid of which "elementary particles", in its own terms, are excitations — one substance, as here.

Closest of all, on both what is fundamental and how gravity is recovered, is a line of Lorentz–Poincaré ether models outside the superfluid-vacuum literature. Arminjon's scalar ether theory [34, 35, 36, 37] takes a fluid ether as the only substance, with the elementary constituents of matter as organized flows in it; requires the ether's sound speed to be matter's speed limit, for that reason; makes clocks slow and rods contract in one ratio set by the local ether density; obtains gravitational waves from the ether's compressibility; and replaces the singularity of collapse with a bounce. Its isotropic version has the exponential metric, agrees with General Relativity at first post-Newtonian order, and does not produce frame dragging. Before it, Podlaha, Sjödin and Broekaert treated the ether's density as a refractive index and scaled rods, clocks and light together; Broekaert reproduces the four classical tests at first post-Newtonian order with the scaling this framework requires (Section 11.5) [38].

Not inherited. What is distinctive is narrow. Superfluid vacuum theory obtains a single substance by making geometry emergent from the condensate, and this framework takes the same position on what is fundamental, so that is not a point of difference. Nor is the refractive route: the ether models above recover gravity's classical tests through a medium whose state scales rods, clocks and light together. The following have not been found in the literature reviewed:

1. *One speed, and its variation is gravity.* The carrying speed varies with the state, and that variation is gravity (Section 11.1), where Arminjon holds the ether's sound speed fixed; no fixed speed remains anywhere in the law.
2. *Rulers and clocks squeezed, not the Ocean.* Near a mass it is the Ocean's rulers and clocks that are squeezed, and patterns fall by refraction rather than by a pressure push.
3. *A current along the time direction.* The candidate mechanism of Section 11.6, a current along the time direction slowed by drag and kept in step.

4. *The Ocean's field is not its own source.* This principle selects General Relativity's geometry, with a point of no return, over the exponential metric that the ether models share (Section 11.7).

5. *The rebuild.* Building and reclaiming as the medium's one process; time made in the build, each place making its own; gravity as the uneven making of time (Sections 12.1 and 12.2).

6. *Antimatter as the reclaiming side of every rebuild.* Hidden because it runs into a past that was never a present, visible only when it has no build of its own, and the present matter because the antimatter is in that past (Section 12.4). The nearest precedent [33] places a single reversal across the big bang and needs right-handed neutrinos and dark matter; this framework has neither.

7. *The collapse reclaims time and disorder.* A beat ends with everything built taken away and the energy returned, giving the next beat a low-disorder start; General Relativity's recollapse keeps its disorder (Section 12.5).

8. *Fifteen built per family.* The empty setting is not built, which keeps the sign that makes induced gravity attract (Equation (37)) and requires neutrinoless double-beta decay (Sections 12.8 and 12.9). The prediction of that decay is shared with other work [33]; the reason for it is not.

9. *The partition.* The division of the Ocean's waves by kind (Section 7): kinds with a lowest energy are held in place as confined patterns ($T^{\text{matter}}_{\mu\nu}$), and kinds without one are coherent ($T^{\phi}_{\mu\nu}$, or, for light, the plexus), with the threshold each kind's lowest energy, Equation (22). The consequence developed in Section 9, that no discontinuous exchange term between the two is required because both are waves of the one Ocean, follows from that construction. It removes an obstruction to the conservation conditions; it does not by itself establish them, which remains outstanding.

10.3 The electron-scale coincidence

The numerical agreement of Section 7 has been approached from the opposite direction. Working from a logotropic equation of state, Chavanis [31] recalls an expression for the cosmological constant in terms of the electron mass and the fundamental constants,

$$\Lambda = \frac{G^2 m_e^6}{\alpha^6 \hbar^4} \approx 1.36 \times 10^{-52}\,\mathrm{m}^{-2}, \tag{26}$$

a relation of Eddington type. There the electron mass fixes Λ; here the electron mass is the lowest energy of the electron's kind of wave, set by the dynamo's field for the present beat (Section 7). Both routes tie the electron scale to a vacuum-scale quantity, from opposite ends.

Two cautions attach to this. The relation is *recalled* rather than derived in that work, so the primary source lies earlier in the logotropic literature. And independent arrival at a coincidence is weak evidence that it is not accidental; it is not this framework's discovery, and the open question of Section 13.2, item (6), remains untouched by it.

10.4 Known deficiencies relative to current practice

Points where this document is behind the literature it should be engaging, or exposed to measurements it has not yet addressed.

(1) **First-order dissipative hydrodynamics.** The constitutive form of Section 2.3 is of Eckart–Landau type, which is known to be acausal and generically unstable in relativistic settings [39]. Current viscous cosmology uses causal second-order formulations of Israel–Stewart type [40]. If $\Pi_{\mu\nu}$ has any role, a viable version of this framework must do the same; in the settled Ocean it carries nothing (Section 2.3).

(2) **Vertical dynamics.** Section 6's decomposition is radial. Milky Way kinematics constrain the two components separately, and they do not want the same thing: the baryonic distribution supplies

too little gravity to account for the rotation curve but enough to account for the vertical velocity dispersions near the Sun, so substantial enhancement is required radially and essentially none vertically [11, 12]. Superfluid dark matter fails this test because its phonon-mediated force is approximately parallel to the baryonic gravitational field and therefore enhances both components by the same factor [12]. Cold dark matter passes because its vertical acceleration is dominated by baryons, with the halo subdominant. The resulting criterion applies to any model: the enhancement of vertical acceleration must be suppressed relative to the radial enhancement. Whether this framework is exposed depends on whether $v_{\phi,\mathrm{eff}}$ tracks the baryonic acceleration field — in which case it inherits the parallelism problem — or constitutes an independent source. Section 6 proposes the second: the tilt of the re-made state in the eddy where the galaxy sits, which turns travel across its direction rather than along the baryonic field. A swirl of that tilt turns a star on a circular orbit inward and leaves vertical travel almost untouched. The first-order law allows the swirl no source in the galaxy, so it is put in place by the eddy, and its shape near the disk then adds about 9 to 15 per cent to the baryons' vertical pull within a kiloparsec of the midplane, against a radial enhancement of more than a factor of two. What puts the tilt in place, and what sets its strength from the baryons, are not yet derived; the full confrontation with the data is not done.

(3) **The status of $c(\phi)$.** A variable effective propagation speed is natural in the emergent-gravity family, where the metric is acoustic. The refractive reading is the framework's account of gravity (Section 2.5), which places it with that family on what is fundamental and incurs the family's chief obligation: to derive the effective geometry rather than assume it. Section 11.1 settles the reading: $c(\phi)$ is the single local carrying speed of every disturbance, the refractive reading, and not a second causal speed; its two readings, the same on the spot and varying from afar, are separated there. What remains is to derive it.

(4) **The vacuum is already measured.** In quantum electrodynamics, vacuum polarization screens every charge out to the electron's reduced Compton wavelength, 386 femtometers, and contributes measurably to the Lamb shift and to the electron's anomalous magnetic moment, in agreement with theory to many significant figures. The vacuum also behaves as a nonlinear optical medium. Light-by-light scattering, forbidden in classical electrodynamics, has been observed in ultra-peripheral heavy-ion collisions [41], and the strong linear polarization of optical light from the neutron star RX J1856.5−3754 is evidence for vacuum birefringence in a strong magnetic field [42]. The Casimir effect is measured. In this framework these are responses of the plexus, the waves light travels on, and of the electron's pattern on it (Section 12.7); they are not responses of the Ocean's own amount at its healing length, which lies near the Planck length. The framework must reproduce them quantitatively from the plexus. Until it does, they stand as an obligation of the same order as the measurements.

(5) **Antimatter is measured.** Every charged localized excitation has an antimatter partner, observed directly: the positron [43], the antiproton and antineutron, antinuclei up to antihelium-4, and bound antihydrogen, trapped and laser-cooled at CERN [44]. Matter and antimatter agree to extreme precision: the antihydrogen 1S–2S transition matches hydrogen to about 2 parts in 10^{12} [45]; the antiproton charge-to-mass ratio matches the proton's to about 16 parts in 10^{12} [46]; antihydrogen falls toward the Earth, measured at $0.75\,g$ with an uncertainty of about $0.2\,g$, consistent with g, and repulsive gravity is ruled out [47]. The one measured difference is CP violation in the weak interaction, seen in kaons, B and charm mesons and, in 2025, baryons [48], and too small in the Standard Model to explain the excess of matter.

The framework must therefore produce, for each localized excitation, a partner of identical mass and magnetic moment and opposite charge, equal response to the slowing of the Ocean (the first of the three decisive requirements of Section 2.5), and room for the small CP asymmetry. A single real scalar field cannot: its excitations are their own antimatter partners. In this framework the partner is the reclaiming side of the rebuild, the same pattern unmade (Section 12.4). The same pattern carries the same mass and magnetic moment; the books of the rebuild give it the opposite

charge; and its own rebuild reads the load at its place exactly as a build does, so it falls as matter falls, as the antihydrogen measurement requires. The room for the CP asymmetry is the measured difference between how builds and reclaimings behave, which is also where the matter excess must come from (Section 12.4). Matter and antimatter of opposite Ocean charge differ in rest energy by 2μ per unit, which would exceed the limits above if the Ocean's phase turned fast. Frame dragging needs no fast phase rate, being carried by the re-made tilt (Section 11.8); with the slow flow the difference 2μ is far below any measurement. What remains to be shown from the rebuild is that the partner's magnetic moment is equal to the limits measured, and that the CP difference has the measured size.

Antimatter also tests the preferred frame. In standard field theory a violation of CPT symmetry implies a violation of Lorentz invariance, and a medium with a rest frame is a natural source of both. The comparisons above are therefore among the most precise existing limits on the Ocean's preferred frame. Antimatter is not a separate obligation from Section 13.2, item (3); it is one of the instruments that tests it.

10.5 The structural question

If the vacuum is a medium possessing density, pressure and a rest frame, it is reasonable to ask why it requires a background spacetime to occupy. The emergent-gravity family answers that it does not, and that geometry is the medium's own collective behavior. This framework answers the question in favor of the medium. There is no background spacetime for the Ocean to occupy; the geometry of Equation (1) is how the medium's state is experienced by excitations of the medium, and $\kappa(\phi)$ expresses how strongly that state responds to concentrated energy. The derivation needs a strict split, or the medium is counted twice; Section 11 states the split and why the minimal law alone cannot carry it. Section 12 states the law that does: the re-made state carries the split, so that only patterns are sources; the first-order law for clocks, rulers and tilt follows from one speed and preservation; and beyond first order the plexus's resistance to bending has Einstein's form. What remains to be derived is in Section 13.1.

End of §10 · token b62bd0

11 One Speed and the Split

Section 10 ends with the task the medium-first position creates: deriving the effective geometry of Equation (1) from the medium, with a strict split between the background that defines the geometry and the excitations that act as sources. This section states the principle on which that split rests, shows why the minimal form of the Ocean's law cannot carry it and where the fixed speed hiding in it lies, states the condition the weak-field limit imposes, offers a candidate mechanism, derives from two of the framework's principles the geometry around a non-spinning mass, with its point of no return, and gives frame dragging around a spinning one to first order. It does not derive Equation (1) in general; that remains the central task of Section 14.

11.1 One speed

The Ocean has one speed: the speed at which it carries a disturbance, written c_s and called here the carrying speed. Everything runs at it — light, the internal workings of every pattern, every signal. The Ocean's own flow is not a disturbance traveling through the Ocean, and has no such limit; "faster than c_s" is always measured against some other part of the Ocean, never against the water a pattern sits in. Where the flow outruns c_s there is a horizon: disturbances still move at c_s through the local medium, but the medium carries them away faster than they can make headway.

What is called the speed of light is therefore not a constant of nature but the Ocean's local carrying speed. It is always measured with the same value because rulers and clocks are themselves patterns

running at that speed; a change in c_s is invisible locally by construction. Gravity is where c_s varies from place to place.

There is therefore no fixed c. Every place a speed appears in the Ocean's law, it is c_s, the Ocean's own carrying speed where it is. The law is written on a single field of rulers and clocks that is part of the Ocean's state, not a stage, and every term uses it: the Ocean's own ripples, every pattern, light. The carrying speed then has two readings. On the spot, every pattern measures the same value, because its rulers and clocks are made of the same Ocean. Seen from afar, it varies, and that variation is gravity. For the Ocean's own ripples to travel at the one speed, the Ocean must be stiff: in the minimal law of Section 11.4 the shortfall of the ripple's speed is $a = 2\mu^2/(\lambda v^2)$, and the coefficient of the k^4 term is about $4a^2$, so stiffness removes both the mismatch of speed limits and the short-ripple overshoot.

There is one speed because there is one substance. A disturbance passes from place to place only by the Ocean handing it along, and the speed of the hand-off is set by two properties of the medium, how stiffly it resists being bent or compressed and how much inertia it has: c_s^2 is their ratio. At any place there is one stiffness and one inertia, and so one speed. A second speed would require a second substance, or a layer running beneath the Ocean, which would be the inherited stage returning.

A localized pattern is a ripple circulating at c_s, confined. At rest all of its carrying speed goes into circulating. In travel part of it carries the pattern forward, and less remains for its internal rhythm, which slows by $\sqrt{1-v^2/c_s^2}$. Relativistic time dilation is then a consequence of the medium, and a pattern can never reach c_s, because nothing would remain to circulate. This is the dynamical reading of relativity first proposed by Lorentz and FitzGerald. The step from "matter is flows in the medium" to "the medium's sound speed is matter's speed limit" has a precedent: Arminjon's ether theory requires the ether's sound speed to equal c for exactly this reason [36]. There that speed is fixed everywhere; in the Ocean it varies with the state, and its variation is gravity.

11.2 The rules are fixed

The Ocean's law is the same everywhere. What differs from place to place is the state a pattern sits in. Every pattern responds to the Ocean around it in two ways: it is carried by currents, and it is refracted by differences of state. Neither is felt from inside: a pattern riding even the swiftest current notices nothing, and its internal workings run as they would anywhere. This is the equivalence principle in the Ocean's terms.

The response may depend on where a pattern is and what the Ocean is doing there, never on what the pattern is made of. Tightly and loosely wound patterns at the same place are carried and refracted identically; composition dependence of free fall is excluded at the level $\eta(\mathrm{Ti},\mathrm{Pt}) = [-1.5 \pm 2.3\,(\mathrm{stat}) \pm 1.5\,(\mathrm{syst})] \times 10^{-15}$ [49].

What breaks a pattern is not the speed of a current but its change across the pattern. A uniform current carries everything along unfelt; a pattern is torn when its front lies in faster flow than its back. A loosely bound pattern spans more of the Ocean and holds itself together more weakly, so it is torn sooner and farther out: stars before people, people before atoms, atoms before nuclei. That is tidal disruption (Section 8.2).

11.3 What sits on each side of the field equation

The background state of the Ocean sets two things: the local carrying speed $c_s(x)$ and the flow $u^\mu(x)$. The effective geometry from which $G_{\mu\nu}$ is built is constructed from these alone, in the acoustic or river form. The background's own pressure $P(\phi)$ belongs to this side, as part of what sets c_s; it is not a source. The right-hand side of Equation (1) holds only what sits above the background: localized patterns ($T^{\mathrm{matter}}_{\mu\nu}$), coherent ripples ($\delta T^{\phi}_{\mu\nu}$), and any effective stress $\Pi_{\mu\nu}$, which in the settled Ocean carries nothing (Section 2.3). Nothing is counted twice. The geometry is the Ocean's own field of rulers and clocks (Section 11.1).

A geometry built from c_s and the flow alone fixes the light cones, and with them the paths of light. It cannot by itself fix the overall conformal factor that governs slow patterns: an acoustic geometry can

match the Schwarzschild geometry only up to such a factor [50]. In the Ocean that factor is supplied by how patterns scale with the state (Section 11.5).

11.4 Why the minimal law cannot carry the split

Two results, computed and checked numerically, show that the simplest law consistent with Section 2.2 cannot deliver the split as it stands. That law, with the quartic potential of Section 9.1, is

$$L = -(\partial\rho)^2 + \rho^2\mu^2 - \frac{\lambda}{4}\left(\rho^2 - v^2\right)^2, \qquad \mu^2 \equiv -\nabla_\alpha\theta\,\nabla^\alpha\theta, \tag{27}$$

with μ the phase rate of Equation (6). At long wavelength the modulus sits at $\bar{\rho}^2 = v^2 + 2\mu^2/\lambda$, leaving a single-field superfluid with pressure $P(\mu^2) = v^2\mu^2 + \mu^4/\lambda$, whose derivative $P' \equiv dP/d(\mu^2) = \bar{\rho}^2$ is exactly the modulus squared.

The phonon geometry gives the wrong light bending. Splitting the phase into background and ripple, $\theta = \bar{\theta} + \delta\theta$, the ripple propagates in the disformal acoustic metric [51, 50]

$$g^{\text{eff}}_{\mu\nu} = \Omega^2\left[g_{\mu\nu} + \left(1 - c_s^2\right)u_\mu u_\nu\right], \qquad c_s^2 = \frac{P'}{P' + 2\mu^2 P''}, \qquad \Omega^2 = \frac{P'}{c_s}. \tag{28}$$

With $a \equiv 2\mu^2/(\lambda v^2)$, $c_s^2 = (1+a)/(1+3a)$. In the medium's rest frame $|g^{\text{eff}}_{00}| = P' c_s$ and $g^{\text{eff}}_{ij} = (P'/c_s)\,\delta_{ij}$, so the post-Newtonian parameter that measures the curvature of space per unit slowing of time is

$$\gamma = -\frac{2+3a}{3a}, \tag{29}$$

always below -1: space contracts where clocks slow, the opposite of General Relativity's $\gamma = 1$. For any single-field superfluid of this kind, $\gamma = 1$ would require P' to stay fixed while c_s varies, which is impossible, since both move with μ^2 alone. In the stiff limit $a \to 0$ the time part does not change at first order, so light bends but slow patterns do not fall; at large a the metric is conformal and $\gamma \to -1$, with no light bending. With the modulus held rigidly at v, $P'' = 0$ and the ripple sees flat space. Matter and light cannot both be reading the acoustic geometry of the phase.

Light and matter have different speed limits. In units where the speed of the flat background in Equation (27) is one, and in the frame turning with the phase, the small disturbances satisfy

$$\left(\omega^2 - k^2\right)\left(\omega^2 - k^2 - \omega_\rho^2\right) = 4\mu^2\omega^2, \qquad \omega_\rho^2 = \lambda\bar{\rho}^2. \tag{30}$$

A gapless branch travels at the c_s above. A gapped branch, the candidate for matter, has rest rhythm $\omega_0^2 = \lambda v^2 + 6\mu^2$. At high k both branches approach $\omega = k \pm \mu$: their speed limit is set by the flat background written into the law, not by the Ocean's state. Two consequences follow. A pattern's rhythm times its size equals its own speed limit, which here is fixed, so where its rhythm slows its size must grow; clocks and rulers move in opposite directions, and the lockstep of Section 11.5 cannot occur. And wherever the Ocean is not fully stiff, light is slower than matter's limit, so fast charged patterns would outrun light and radiate it away. Cosmic-ray protons near 10^{20} eV bound such a difference to about one part in 10^{23} [52], and for electrons to about one part in 10^{15} [53]. Since $1 - c_s \approx a$ for small a, the Ocean under this law would have to be stiff to $a \lesssim 10^{-23}$, where c_s stops responding to the state and gravity disappears.

The diagnosis: Equation (27) is written on a flat background, a stage the framework does not have. Every term of the form $(\partial\Phi)^2$ means $(1/c^2)(\partial_t\Phi)^2 - (\nabla\Phi)^2$ with c a constant of that stage, and the fixed speed appears in three places: in how matter's inner workings move (the mismatch above), in the short-ripple overshoot of the k^4 term, and in the law's own bookkeeping, where $c_s^2 = dp/d\varepsilon$ is measured against it. The last is why the weak-field condition of Section 11.5 and an equation of state

for Section 11.6 written against that speed contradict each other at first order: one is the on-the-spot reading of c_s, the other the reading from afar. With the law written on the Ocean's own rulers and clocks (Section 11.1), stiffness, $a \lesssim 10^{-23}$, is required rather than fatal: gravity is the change of the rulers and clocks seen from afar, not of the on-the-spot ripple speed. Writing that law out in full is the central research problem (Section 14).

11.5 The weak-field condition

Near a mass c_s is slower. Light bends because c_s varies along its path. Slow patterns fall because their internal rhythm is slower deeper in, and a pattern drifts toward where its rhythm, and so its energy, is lower: refraction for matter. The one-speed principle ties a pattern's rhythm and size together, rhythm times size equaling c_s, which makes the change invisible locally. It leaves open how the slowing is shared. The standard post-Newtonian account settles it: the equivalence principle alone gives half the observed deflection of starlight, and the curvature of space, measured by γ, gives the other half [54]. General Relativity's light bending therefore requires an even split,

$$\text{rhythm} \propto \sqrt{c_s}, \qquad \text{size} \propto \sqrt{c_s}. \tag{31}$$

The same scaling — a coordinate light speed falling as Φ^2, rods and clocks each scaling as Φ, inertia as Φ^{-3} — was used by Broekaert, after Sjödin, to reproduce the four classical tests at first post-Newtonian order [38]; Equation (31) is that condition stated for the Ocean. Large bodies follow automatically: a star is made of patterns, and if each scales alike the star falls as its parts do. The condition is a rule for the weak field: close to a mass it gives way, and Section 11.7 gives what replaces it. A route through the Ocean's inertia, by which a rising phase rate near a mass would produce the scaling through a laboratory-superfluid healing length, fails: that formula does not carry over to the Ocean, and combined with an equation of state written against a fixed speed it contradicts the Ocean's thermodynamics at first order (Section 11.4). Equation (31) enters only as one constant, $\gamma = 1$, and is otherwise not needed.

11.6 A candidate mechanism: the current through time

This subsection is a candidate, not a derivation. Ordinary sideways flow past a star cannot be the source of its gravity: the Sun travels at about $370\,\mathrm{km\,s^{-1}}$ relative to the microwave background, and gravity from flow past it would be lopsided, stronger on its flanks than ahead and behind, which the post-Newtonian preferred-frame parameters exclude at $\alpha_1 \lesssim 10^{-4}$ and $\alpha_2 \lesssim 2 \times 10^{-9}$ [54]. One current is the same whichever way a star travels: the flow that carries every pattern forward in time. On the one-speed principle every pattern moves through the Ocean at c_s in total, all of it along time when at rest.

(i) **The star slows the current.** A star exists at every moment, so it lies along the current through time rather than across it, and a frictionless flow along such a body is undisturbed. The star must therefore drag on the current, through the medium's viscosity. The slowed current spreads sideways through three space directions, so the deficit falls off as $u \propto M/r$: Newton's inverse distance, with a strength set by the drag per pattern relative to the Ocean's viscosity.

(ii) **Clocks slow by u.** Patterns are carried through time by the local current, so their rhythm slows by u. This is the half of gravity that makes things fall.

(iii) **The Ocean stays in step.** A river speeds up past an island because the island blocks the water and the same amount must squeeze past its flanks. A star blocks nothing: it is made of the Ocean, and it lies along the current, which has always flowed past it. What the Ocean must not do is fall out of step with itself. If the Ocean near a star turned more slowly, seen from afar, than the Ocean far away, the mismatch would grow at every moment, and a growing mismatch is a growing current: an upstream wake, which nothing observed supports. So, seen from afar, the Ocean's phase turns at one rate everywhere. Measured on the spot, where clocks run slow, it turns faster, by exactly the clocks' slowing, $\mu_{\mathrm{local}} = \mu_\infty / U$ with U the rate of clocks seen from afar. This is the relativistic

form of Bernoulli's relation for a fluid at rest in a gravitational field: Klein showed that its chemical potential, including the mass energy, times $\sqrt{|g_{00}|}$ is the same everywhere [55], and the Ocean's chemical potential is its phase rate (Section 2.2). On the spot the current through time is always at the carrying speed, so all change shows in the view from afar.

(iv) **The amount of Ocean stays the same.** The Ocean is stiff (Section 11.1): its amount ρ barely gives. Near a mass it is not the Ocean that is squeezed but its rulers and clocks. No pressure–density law is needed for this step, such as one of generalized Chaplygin form [56, 57].

(v) **Far from a mass: clocks and rulers in lockstep.** The three rules above fix clocks and the Ocean's turning, but not rulers. Far from a mass the lockstep of Equation (31) supplies them, and to first order

	General Relativity	Candidate
Clock rhythm	$1-u$	$1-u$
Light speed seen from afar	$1-2u$	$1-2u$
Pattern size ($=c_s/$rhythm)	$1-u$	$1-u$

Write the rate of clocks seen from afar as U, so that $|g_{00}| = U^2$, and the slowing as $u = GM/(rc_0^2)$. If each layer of the Ocean subtracted a fixed amount from the current, $U = 1-u$, the post-Newtonian parameter β would be $\frac{1}{2}$ and Mercury's perihelion would advance $(2+2\gamma-\beta)/3 = 7/6$ times too fast. If each layer slows the current by a fraction of what it receives, $U = e^{-u}$ to second order and $\beta = 1$. Rulers set instead so that the amount of Ocean in a region looks fixed from afar would fail: with the amount fixed on the spot (item (iv)) and the phase turning faster there by the clocks' slowing (item (iii)), the Ocean's amount counted from afar, the time part of its current (Equation (13)), changes by $(1+3\gamma)u$; holding it fixed gives $\gamma = -\frac{1}{3}$, and light would bend a third as much as it does.

(vi) **Close to a mass, lockstep gives way.** Taken all the way in, lockstep gives the exponential metric of Yilmaz, $|g_{00}| = e^{-2u}$, $g_{ij} = e^{2u}\delta_{ij}$, which Arminjon's isotropic ether theory has in full [35] and which underlies Broekaert's model [38]. In it outward travel slows without limit but never stops: there is no point of no return, and the shadow of a black hole would be 4.6 per cent larger than General Relativity's, outside one of the two ranges the Event Horizon Telescope allows for Sagittarius A* [26]. A horizon is a point of no return: before it escape is possible, after it, it is not. The Ocean needs one, so lockstep is a rule for the weak field only. Section 11.7 gives the rule that replaces it close in. Water's form of Bernoulli's relation, applied along the time direction, would put a horizon at $2.9\,GM/c_0^2$; the rule of item (iii) takes its place.

The candidate's inputs are the first-order lockstep constant, equivalent to the observed bending of light, and the strength of the drag, which sets G. The composition of the slowing in layers is not an input: it follows from the rules of Section 11.7. One cosmological requirement stands on its own: unified fluids whose pressure is a function of one density are excluded as the cosmic medium once $|d\ln p/d\ln n|$ exceeds about 10^{-5}, because their sound speed erases the growth of structure on galactic scales [58]. The Ocean has two state variables, the amount ρ and the phase rate μ (Section 2.2), so its cosmological pressure need not be such a function; that it is not is to be shown.

11.7 The strong field and the point of no return

Two principles the framework already holds fix the geometry around a non-spinning mass completely. Write it in isotropic form,

$$ds^2 = -U^2 c_0^2 dt^2 + \psi^4 \left(dr^2 + r^2 d\Omega^2\right), \tag{32}$$

with U the rate of clocks seen from afar and ψ^2 the number of the Ocean's own rulers that fit in a unit step of r.

(a) **Staying in step, counted with the Ocean's own rulers.** The slowing of clocks spreads from the mass by the inverse-distance law, with distance counted by the Ocean's own rulers, which shrink near the mass, so that more of them fit across a given span and $\psi^2 > 1$, and not by a flat ruler underneath (Section 11.1): $\nabla \cdot (\psi^2 \nabla U) = 0$.

(b) **Nothing counted twice.** Where there is no pattern, the Ocean around a mass does not squeeze its own rulers; the field is not its own source (Section 11.3): $\nabla^2 \psi = 0$.

With the first-order lockstep used only to fix one constant, $\gamma = 1$, the solution is

$$\psi = 1 + \frac{GM}{2rc_0^2}, \qquad U = \frac{1 - GM/(2rc_0^2)}{1 + GM/(2rc_0^2)}, \tag{33}$$

General Relativity's geometry outside a non-spinning mass. The consequences, each checked numerically:

- **The point of no return.** U reaches zero, seen from afar, at isotropic radius $GM/2c_0^2$, areal radius $2GM/c_0^2$: General Relativity's horizon, produced by the geometry rather than assumed. It is the same point for every pattern, since every pattern has the same carrying speed.
- **Mercury, derived.** To second order U agrees with e^{-u}, so $\beta = 1$ follows; the slowing composes in layers as a consequence, not an input.
- **The shadow.** Light orbits at areal radius $3GM/c_0^2$ and the shadow's radius is $3\sqrt{3}\,GM/c_0^2$, as in General Relativity. The Event Horizon Telescope finds the shadow of Sagittarius A* within about ten per cent of that value: a fractional deviation of $-0.04^{+0.09}_{-0.10}$ or -0.08 ± 0.09 for the two determinations of its mass and distance, and no evidence of a surface [26]. The orbit of the star S0-2 about the same black hole gives $(2 + 2\gamma - \beta)/3 = 1.1 \pm 0.19$ (reported there); the Ocean gives one.
- **What separates the Ocean from the nearest precedents.** The exponential metric of Section 11.6 satisfies rule (a) but not rule (b): where there is no pattern it squeezes the Ocean's own rulers, which counts the medium twice. Rule (b) is the split of Section 11.3, and it is what selects a point of no return.

The derivation assumes a static, spherically symmetric mass and the isotropic slicing, which spherical symmetry always allows. A spinning black hole needs the Ocean's account of frame dragging beyond first order in spin (Section 11.8), which is not given; the Kerr shadow varies by about eight per cent with spin and viewing angle [26].

11.8 Frame dragging

[Candidate] Near a spinning body a gyroscope slowly turns, and so does the plane of an orbit, as if the body carried the directions around it along with its spin. This is frame dragging, and it is measured around the Earth. Gravity Probe B's gyroscopes turned by -37.2 ± 7.2 milliarcseconds a year, against General Relativity's -39.2 [59]. The orbit planes of the laser-ranged satellites LAGEOS, LAGEOS 2 and LARES turn as General Relativity gives to about two per cent [60], and with LARES-2 to $1.001 \pm 0.001\,(\text{stat}) \pm 0.002\,(\text{syst})$ of its value [61]. The two principles of Section 11.7 are written for a mass that does not spin, and do not give it. This subsection gives frame dragging to first order in spin from inside the Ocean, with General Relativity used only for comparison.

The rule. A pattern's disturbance includes the slowed Ocean around it (Section 6). When the pattern travels, the whole disturbance is re-made with it, and the re-made surroundings carry the direction of travel: the current through time there is turned partly sideways, in proportion to the travel. What is re-made is the Ocean's state, not the Ocean carried along, as for the river picture of Section 8.1.

How far the current is turned. Travel tilts whatever is re-made with it, by the one speed (Section 11.1). Seen from Ocean that is not traveling, the surroundings of a part of mass m traveling at $\mathbf{V}$ carry the tilt of both their slowed clocks and their shrunk rulers:

$$\mathbf{s} = (2+2\gamma)\,u\mathbf{V}, \qquad u = \frac{Gm}{c_0^2 d}, \tag{34}$$

with d the distance from the part and u the slowing of Section 11.6. The clock share is $2u\mathbf{V}$, the ruler share $2\gamma u\mathbf{V}$, and $\gamma = 1$ (Section 11.5). In the geometry, $\mathbf{s}$ is the shift, $g_{0i} = -s_i/c_0$; it is what a resting mass's geometry becomes when seen from a frame in which the mass travels.

A spinning body. The Earth is Ocean, and its spin is its own patterns traveling round its axis; a black hole's spin is the net circulation of its own Ocean (Section 8.1). One rule serves both. The slowing alone does not change with spin, since the amount of energy stays where it is; the tilts of the traveling parts add up. Outside a body of spin $\mathbf{J}$,

$$\mathbf{s} = \frac{2G}{c_0^2}\,\frac{\mathbf{J}\times\mathbf{r}}{r^3}, \tag{35}$$

checked numerically by adding the tilts of the parts of a two-layer model Earth. The turning of this tilt, $\nabla\times\mathbf{s}$, has the pattern of a bar magnet's field: along the spin near the poles, against it near the equator, and falling off as the cube of distance.

What it does. In Equation (15) a pattern traveling through turning Ocean is turned sideways by $-\mathbf{w}\times(\nabla\times\mathbf{s})$. Applied to every part of a spinning rotor, that term turns the rotor's axis at half the local turning, $\frac{1}{2}\nabla\times\mathbf{s}$, whatever the rotor's own spin: Larmor's theorem, checked numerically. That is the turning of a gyroscope. Applied to an orbit, the same term turns the orbit's plane. For the Earth, with its measured spin:

	This rule	Measured
Gravity Probe B gyroscopes, toward the guide star	-39.2	-37.2 ± 7.2
LAGEOS, LAGEOS 2, LARES orbit planes	30.6, 31.4, 118.1	0.991 ± 0.02
LARES-2 with LAGEOS	—	1.001 ± 0.002

Rates are in milliarcseconds a year. For the orbit planes the measured turning is given as a fraction of General Relativity's, whose values for the three orbit planes are 30.68, 31.50 and 118.50 [60]; the rule's agree with them to within the rounding of the Earth's spin used here. General Relativity gives the same frame dragging; here it is a comparison, not an input.

Travel through the Ocean leaves no trace. The Earth also follows the Sun and travels through the Ocean, and every part of it carries that shared travel. The tilt it adds is, near the Earth, several thousand times the spin's, and would swamp the measurements if a gyroscope traveling with the Earth registered its turning. It does not. Such a gyroscope is also turned by its own travel through the slowed Ocean, and the two cancel exactly when the tilt carries both the clock share and the ruler share of Equation (34). Checked numerically, with a gyroscope that keeps its direction as the Ocean's rulers and clocks carry it: with both shares the travel leaves no turning; with the ruler share missing a turning of $|\mathbf{V}\times\nabla u|$ remains, and with no tilt $(1+\gamma)\,|\mathbf{V}\times\nabla u|$, for travel $\mathbf{V}$. What travel with the Earth around the Sun does leave, the turning of the lunar orbit by its travel through the Sun's slowed Ocean, is measured as General Relativity gives it, to 0.6 per cent [62]. Any trace of travel through the Ocean itself is bounded at about one part in 10^4 [54].

What it requires. The tilt around a spinning body turns everywhere outside it. The Ocean's flow, as Section 2.2 defines it, is the gradient of its phase, and a gradient does not turn. So the turning is not carried by the phase. It is carried by the re-made state: the direction in which each part of the surroundings is rebuilt is part of what the rebuild hands forward (Section 12.1), the motion being part of the build, and that direction turns continuously wherever the surroundings reach. The phase needs no whirl at all. Around a black hole spinning near its limit, the phase of Equation (36) (Section 11.9) extends as $\theta = \mu_\infty[t + F(r)]$ with $\Delta F' = 2Mr_+$, where $\Delta = r^2 - 2Mr + a^2$, a here is the hole's spin per unit mass, and r_+ is the outer point of no return, in the coordinates of General Relativity's spinning hole: zero circulation on every loop, smooth across the point of no return, while the Ocean is still carried around at the rate gyroscopes measure, from the hole's own rate at its edge to the first-order rate far away: the turning is entirely the re-made tilt.

Quantized lines of the phase could carry the turning instead, each with circulation hc_0^2/μ [63]. Read point by point, they do not turn a gyroscope; read over at least one line spacing, they carry it exactly. Checked numerically, a gyroscope whose parts read a field of lines over one spacing turns at half the average turning, as for smooth turning, whatever its own spin. But the lines are fine enough only if the phase turns fast, of order 10 keV or more, which no stiff Ocean allows, and they need a reading scale that is assumed. With the turning in the re-made tilt, neither is needed, and frame dragging is not carried by lines. In superfluid helium-3 A the flow is likewise the twist of an internal set of directions, its turning continuous, as the Mermin–Ho relation states, and that turning acts on the excitations as a gravimagnetic field [64].

What is assumed. Two things. That patterns read the Ocean's turning as Equation (15) says, assumed in Section 6 and not derived; the gyroscope's turning then follows. And that the surroundings are re-made with the pattern exactly, with no lag and no wake, below the critical speed of Section 14: this is the condition that travel leaves no trace, assumed here rather than derived. That the slowed surroundings obey the one speed as patterns do, so that travel tilts them, is not assumed: the first-order law of clocks, rulers and tilt, fixed by one speed and preservation, gives the tilt from travel with the coefficient $2 + 2\gamma$ of Equation (34) (Section 12.7).

Beyond first order. The two principles of Section 11.7 keep the Ocean's rulers shrunk equally in every direction. Around a spinning hole they cannot be: among axisymmetric slicings of General Relativity's spinning hole that reduce smoothly to the non-spinning case, none keeps them so beyond first order in spin [65]. At first order there is no obstruction, and this subsection stays there. Beyond it, the same rule must give a spinning hole whose outside depends only on its mass and spin (Section 8.1), and an Earth whose outside also carries the shape of its mass. The shape and shadow of a spinning hole (Section 11.7) and the ringdowns of mergers wait on it (Section 14).

11.9 At the edge and inside

Nothing happens at the edge. If the Ocean stayed in step while holding still all the way down to the point of no return, it would there have to turn infinitely fast, and a traveler falling in would meet it turning ever faster. It does not need to. In the stiff Ocean the phase obeys the wave equation on the geometry of Equation (33), and that equation has a solution that is in step far away and smooth across the point of no return. With r now the areal radius, the one a sphere's surface measures (not the isotropic radius of Equation (33)), and in units with $G = c_0 = 1$,

$$\theta = \mu_\infty\,[t + f(r)]\,, \qquad r^2\left(1 - \frac{2M}{r}\right) f'(r) = 4M^2. \tag{36}$$

Far away $f' \to 4M^2/r^2$ and the phase turns at one rate seen from afar; at the point of no return the phase gradient stays finite for a falling traveler. Near the edge the Ocean keeps in step by flowing gently inward, not by turning faster. The traveler falls with it and meets nothing there, as the equivalence principle

requires. The Ocean is whole and connected across the point of no return; what is cut off is outward travel. The inflow carries energy in at a rate proportional to $\mu_\infty^2 M^2$: the black hole takes in the Ocean's background very slowly, at a rate set by its pressure's departure from $w = -1$. The solution uses the phase equation of the minimal law (Equation (13), with ρ held); under the law Section 14 requires it must be redone, and the inflow stays slow only if the energy carried by the phase's turning stays small.

Inside. Past the point of no return, forward in time points inward, and the Ocean's own phase flows inward with whatever falls. A pattern's time is its rhythm; as patterns are stripped, their clocks go with them.

The core: flow-through. Nothing is gathered at the center. Recycling, for the Ocean, is flow-through: the inward flow does not stop or collect at the center, but passes through and returns to the Ocean, later. Time is recycled with it: inward on the way in, outward again on the way out. A black hole and a star are the two halves: a star sends energy outward from a dense core, and the return from a black hole runs the same way. Two results bear on how.

- **Stiffness cannot turn the flow around.** For any material with $\varepsilon + p \geq 0$, which includes the stiffest that one speed allows, the rate of change of a collapsing region's expansion is $dH/dt = -4\pi G(\varepsilon + p) \leq 0$; once collapse is under way inside the point of no return it cannot reverse. The Ocean's resistance to compression ($\lambda > 0$, Section 9.1) is not enough.
- **The Ocean's grip can.** Equation (1) carries $\kappa(\phi)$, how strongly the Ocean's rulers and clocks respond to what is in them. If that grip loosens under extreme compression, $G_{\text{eff}} = G(1 - \rho/\rho_c)$, the collapsing interior obeys $H^2 = (8\pi G/3)\rho(1 - \rho/\rho_c)$: collapse slows, stops at ρ_c and reverses. In a ball of patterns falling in from rest far away, the time from crossing the point of no return to the turnaround, on a clock falling with it, is about 10^{-4} s for a black hole of ten solar masses, 28 s for Sagittarius A* and 12 hours for M87*; nothing is crushed to a point. A loosening triggered by density would also act inside neutron stars, making them fall differently from white dwarfs; the triple pulsar PSR J0337+1715 shows they do not [66], which places $\rho_c \gtrsim 2.4 \times 10^{23}\,\text{kg}\,\text{m}^{-3}$. If ρ_c is set by a length ℓ through $\rho_c c_0^2 \sim \hbar c_0/\ell^4$, then $\ell \lesssim 0.035$ femtometers, some ten thousand times finer than the electron's localization length; with the dent near the Planck energy, ℓ would be near the Planck length (Section 11.10).

The return, seen from afar. On a falling clock the turnaround is quick; for an outside observer the delay is set by the region just outside the point of no return, through which the reversed flow must come back. That region depends on how the Ocean's grip loosens there, which is a property of what the Ocean is made of. Written as a function of the fine scale ℓ, the delay is $\tau \sim (GM/c_0^2)^{q+1}/(c_0\,\ell^q)$, with $q = 1$ if the loosening builds up quickly near the edge and $q = 2$ if slowly; both forms are borrowed scalings, to be derived. The $q = 1$ form is a random walk across the black hole's size in steps of ℓ, the same rule that keeps light made in the Sun's core inside the Sun for tens of thousands of years or more; what comes out is not what went in, as gamma rays made in the core leave the surface as sunlight. Black holes of a few solar masses are seen merging after billions of years; taking five solar masses and ten billion years, with $q = 1$ the step must be $\ell \lesssim 6 \times 10^{-19}$ m, close to the scale, about 10^{-18} m, down to which the electron shows no size; with $q = 2$ any ℓ below 0.035 femtometers passes. For Sagittarius A* the return then takes of order 10^{22} years or longer. Every known black hole returns what it holds far in the future, consistent with none having been seen to do so. The release, if a random walk, would leak out gradually and thermalized, like starlight, rather than as a single burst.

11.10 What the Ocean must be made of

The framework does not say what is beneath the Ocean. It does say what the Ocean is: its amount and phase, together with the re-made state, held by the plexus the dynamo keeps running (Section 12.7). Any substrate beneath it must satisfy four requirements: a single speed, so that no signal passes through it faster than c_s, which would return the stage one level down; a grain, if there is one, hidden below every tested scale; structure that can be twisted; and the ability to hold confined, circulating patterns.

The third follows from light. Light's oscillation is entirely transverse, in two independent directions; crossed polarizers extinguish it. A complex field is one complex number at each point, a size and a phase, and a number has no direction, so every free ripple of it is sound-like and could not be polarized. Light is therefore not a ripple of the Ocean's amount and phase; it is a wave of the plexus (Section 12.7), and the plexus must carry something with a direction that can be twisted, whose twisting and compressional ripples cannot travel at different speeds, as they do in a solid. The same demand appears twice more: gravitational waves are transverse, in two patterns, which the re-made state's pushes supply; and spin-$\frac{1}{2}$ patterns need a direction with an angle around it, which each unit's direction of time-making and hand supply (Section 13.2, item (7)). Frame dragging adds nothing further: it is carried by the re-made tilt (Section 11.8).

Black holes add two numbers (Section 11.9). The substrate must turn collapse around only at densities no known object reaches: the triple pulsar PSR J0337+1715, whose neutron star falls as its white dwarf does, places a density-triggered turnaround above about $2 \times 10^{23}\,\mathrm{kg\,m^{-3}}$, and with the dent near the Planck energy it lies near the Planck density. And if the Ocean's grip loosens quickly near a point of no return, the substrate must redirect flow at spacings of about 6×10^{-19} m or less. Its account will fix how long a black hole takes to return what it holds.

End of §11 · token 5ab9ee

12 The Rebuild and the Law

The sections above describe what the Ocean does to patterns in it: it slows their clocks, turns their travel, carries their spin. This section describes what the Ocean does at every moment to make those patterns at all, and states the law that follows. Travel was already defined as a pattern being re-made farther along (glossary); here the re-making itself becomes the account.

12.1 Building and reclaiming

Every pattern of the Ocean is rebuilt continuously. The rebuild has two sides, and nothing else.

- *Building* makes the pattern: three arrows and a left twist. It is the only side that makes anything.
- *Reclaiming* takes the pattern just made back down: three arrows and a right twist. It makes nothing; it only takes away.

There is one kind of build. What distinguishes one pattern from another is not a second kind of build but which of the Ocean's waves the build is made on (Section 12.7). Nothing in the Ocean switches. Every change, a decay included, is a reclaiming of what was and a building of something else, with the books balanced. The energy the reclaiming takes down is handed forward to the build, which is why a pattern carries straight through from one moment to the next and never vanishes and reappears. The motion is part of what is handed forward: each part of a pattern is rebuilt a little farther along its path. That is inertia. If the motion were not part of the build, every pattern would stop at every rebuild.

The Ocean itself is rebuilt in the same way. Each place makes its own time (Section 12.2), so the quiet Ocean between galaxies is rebuilt as surely as the Earth: every unit of it carries the build's forward direction and left hand, its re-made state is re-made at every place, and the plexus is built and reclaimed at every moment (Section 12.7). Three things differ. A pattern carries more than a place does: its kind, its version, its extent, its pairing and its loan (Section 12.6). A pattern's build is supplied from the Ocean

around it, while the Ocean's own build is supplied by its flow, through the dynamo; that is why the drive running out ends a beat (Section 12.5). And the reclaiming takes away time, structure and disorder, but never the Ocean's energy, which returns to the Ocean: the Ocean is what every build is made from and returned to.

12.2 Time is made in the build

There is no time already present for the Ocean to move through. Each build makes the moment, and time is the building itself, not a background against which it happens.

1. Each place makes its own time. How fast a place is rebuilt is its clock; there is no single clock for the Ocean.
2. Gravity and time are one act. The rebuild runs at the one speed. Where there is more energy to rebuild, the rebuild takes longer, so less time is made there; that is the slowing of clocks near a mass. A pattern beside it, its near side made more slowly than its far side, is rebuilt a little toward the slow side each time; that is falling (Section 2.5). Gravity is the uneven making of time.
3. Order comes from what is handed forward. Each build is made from the one before it, so there is a before and an after without any clock outside the Ocean.
4. Backward means unmade. The reclaiming does not travel back along a time that exists; it takes down time that was made.

For the law this has one consequence above all others: no time outside the Ocean appears in it. The rate at which the Ocean's phase turns (written μ in Section 2.2) is each place's clock, read off the Ocean, not a rate measured against some other clock. Time dilation (Section 11.1) and the slowing of clocks near a mass are the same reading: how much of the build's pace goes into travel, and how much the load at that place allows.

12.3 The rebuild of the Earth

The Earth, seen from inside the Ocean, is about 4×10^{51} confined patterns, each rebuilt of order 10^{23} times a second, pressed at 360 GPa at its center, spinning, orbiting the Sun at $30\,\mathrm{km\,s^{-1}}$ and traveling about $370\,\mathrm{km\,s^{-1}}$ relative to the frame in which the microwave background is uniform. Its rebuild, described at one moment (the rebuild is continuous; the order below is an order of description, not of ticks), runs as follows.

0. *What keeps it running.* The Ocean's flow powers a dynamo; the dynamo keeps the plexus running, the Ocean-wide waves on which every pattern is made; the flow's left twist is the twist every build carries.
1. *Reclaiming.* Every part of the Earth just made is taken down, with a right twist. This side is antimatter (Section 12.4).
2. *Handed forward.* What the reclaiming takes down supplies the build.
3. *Building.* Each part is built, with a left twist, on the waves that make it what it is. Its three arrows carry its travel and the three directions in which it is pressed or stretched.
4. *The build meets its own reclaiming.* The two match exactly, so the antimatter side stays hidden. Measurement puts any mismatch per rebuild below about one part in 10^{45}.
5. *The motion is part of the build.* Each part is rebuilt a little farther along its path.
6. *The scoop.* Rebuilding this much takes more than the Earth alone carries; the extra is borrowed from the surrounding Ocean and returned as the Earth moves on. What is out on loan is the Earth's binding energy, about four parts in 10^{10} of its mass.

7. *Slower where there is more to rebuild.* The rebuild uses a share of the Ocean's capacity for rebuilding, about 1.4×10^{-9} at the Earth's surface; a clock runs at $\sqrt{1 - \text{share}}$, so clocks there run slow by half that, about 7×10^{-10} (Section 12.7).

8. *The surroundings are rebuilt with everything the Earth carries.* Its energy sets the clocks' slowing; its motion sets the tilt that turns with it, which is frame dragging (Section 11.8); its pushes set the rulers' stretch and squeeze.

9. *Version changes lay down antimatter that escapes.* Uranium, thorium and potassium decay inside the Earth. In each decay the rebuild reclaims one version and builds the other, and the books force one electron and one antineutrino to be laid down (or, when potassium captures an electron, the electron is picked up and a neutrino laid down). The antineutrino has no build of its own to meet, and leaves. About 10^6 of them from uranium and thorium cross each square centimeter of the surface each second; they are detected coming up out of the ground [67].

10. *The record spreads at the one speed.* What the Earth has just been, its energy, motion and pushes, spreads outward through the plexus at c_s. It reaches the Moon about 1.28 s later and, because it carries the Earth's motion, arrives aimed at where the Earth is, not where it was.

11. *Others rebuild themselves reading it.* The Moon's own build reads that record across its body. Its near side is made slightly more slowly than its far side, so it is rebuilt a little toward the Earth each time; with the share at the Moon's distance, 2.3×10^{-11}, the turning $(c_0^2/2)\nabla(\text{share})$ is $2.70 \times 10^{-3}\,\text{m}\,\text{s}^{-2}$, the Moon's measured fall toward the Earth. Nothing reaches across to move it.

12. *The loan is returned,* and the Earth just built is the one reclaimed next.

12.4 Antimatter is the reclaiming

Antimatter is the reclaiming side of the rebuild: the same pattern, unmade. The reading of an antiparticle as a pattern whose own time runs counter to ours is Stueckelberg's, as used in [33]; here it is not a reading of individual objects but a side of every rebuild.

- *Why it stays hidden.* The reclaiming runs into a past that never existed as the present. The past we know is the built past: every moment of it was once a present and left records. The reclaiming's past was never anyone's present, so nothing sees it, and nothing it does reaches us.

- *Why the present is matter.* Every build has its reclaiming, so the books balance overall; the present holds the builds. We see more matter in the present because the antimatter is always in the past we cannot see.

- *Visible antimatter.* When energy alone starts a build and a reclaiming that do not belong to each other (pair creation, which needs twice the lowest energy of the kind made), the reclaiming has no build to meet. It travels through the present, where it is seen, until it meets a build of its kind; the two balance off, and the energy returns to light. Hidden antimatter met its own build; visible antimatter did not.

- *What it must reproduce.* At the start of the beat this happened in enormous numbers, and almost every pair met and returned to light, which is now the microwave background. What remained is about six matter patterns per ten billion of light: the baryon-to-photon ratio is $(6.104 \pm 0.058) \times 10^{-10}$ from the microwave background and $(6.148 \pm 0.191) \times 10^{-10}$ from the light elements, in agreement [68]. The rebuild must give that number. The measured difference between how builds and reclaimings behave [48] is where to look for it.

12.5 The beat's turn and the collapse

The same two sides carry the beat to its end (Section 5.2).

1. When the Ocean's energy has spread out and no high-to-low differences remain, the flow no longer drives the dynamo, and builds are no longer supplied.
2. Nothing switches. The reclaiming has been running at every moment; once the building is no longer supplied, the reclaiming carries the Ocean. Because the spent Ocean reads the same everywhere, a condition read at each place is met everywhere together, and the contraction is global. A partial contraction could not be stopped, and would be a black hole.
3. Running forward makes matter; running backward takes everything away. In the collapse nothing new forms. Time is reclaimed with everything else: the time the beat made, the structure built in it and its disorder. Black holes, if any remain, are reclaimed too.
4. Every place reads the same record, so nothing falls toward anything; the collapse is a common drawing-in, not a fall. The load rises together everywhere until the share of capacity in use reaches one everywhere at once: a point of no return with no outside.
5. The energy is not destroyed. It returns to the Ocean, compressed, and the next beat begins from it.

Here the Ocean parts from General Relativity, in which a recollapse keeps its matter and so keeps its disorder. Disorder is the arrangement of what was built; once the reclaiming has taken everything away, nothing remains to be disordered. That would give each beat the low-disorder start the arrow of time requires.

12.6 What a build knows

Walking the rebuild step by step gives what each pattern must have at hand. Four things are rules, the same everywhere and not variables: the one speed c_s; the Ocean's capacity for rebuilding; the dent's tuning (Section 12.7); and that the books must balance.

	Carried by each pattern	
1	mode	building (left twist) or reclaiming (right twist)
2	energy	
3	motion	direction and speed of travel
4	pushes	three directions and their sizes
5	kind	which of the Ocean's waves it is built on
6	version	which of two versions (what changes in a decay)
7	what must balance	what measurement calls charge
8	pairing	which reclaiming is its own, or none
9	loan	what it has borrowed from the surrounding Ocean
10	extent	how far across the Ocean it reaches
	Read where it is	
11	the making of time	which way time is being made, from what was handed forward
12	the plexus	the waves present
13	the load	the share of capacity in use
14	the record	what arrives from others, and how it differs across the pattern

At the beat's turn three more are read: whether a high-to-low difference is available, the balance of building and reclaiming at that place, and the twist of the plexus waves present. Everything a build needs is at its own place, what it carries and what arrives there. Nothing reaches across.

12.7 The law

What the Ocean is. Its amount and phase, each unit carrying the direction in which time is made and a left hand, together with the re-made state, ten numbers at each place (energy one, motion three, pushes six), held by the plexus. The plexus is Ocean-wide waves, made and kept running by the dynamo; it is not a structure. The flow powers the dynamo; the Ocean's own scale, the cost of a dent, tunes it, as a resonator sets the pitch of a steadily driven string. The substance is best described as a new superfluid: frictionless for what moves through it, one connected phase, a direction and a hand at every place, with no container and no outside, and a grain that nothing moving through it can feel. Its flow conducts: currents in the flow make and read the plexus. The leakage a dynamo needs in order to run is not friction but the plexus's own reclaiming, which the flow must keep supplying; nothing is dissipated, so nothing moving through the Ocean feels drag.

The rules.

Quantity	Rule
mode	building makes, with a left twist; reclaiming unmakes, with a right twist; nothing switches
energy	handed forward from the reclaiming; conserved; recycled, never consumed
motion	carried in each rebuild; turned by the record read across the pattern, at a rate $(c_0^2/2)\nabla(\text{share})$
clocks, rulers, tilt	the unique first-order law fixed by one speed and preservation (Section 11.5); outside a resting mass, General Relativity's exact geometry (Section 11.7); the tilt carries frame dragging (Section 11.8)
pushes	rebuilt with the pattern; ripple off as the two stretch-and-squeeze waves, at c_s
what must balance	kept in every rebuild; what is laid down in a change is forced by the books
pairing	a build meets its own reclaiming (hidden); an unmatched reclaiming travels until it meets one of its kind (visible)
loan	borrowed from the surrounding Ocean, returned as the pattern moves on; its size is the binding energy
load	$\text{share} = \text{energy}/(\text{capacity} \times \text{distance})$; a clock runs at $\sqrt{1-\text{share}}$; full use is a point of no return
record	spreads outward at c_s, carrying energy, motion and pushes, aimed at the source's present position
the dynamo	supplied by the flow; its leakage is the plexus's own reclaiming

The share's form is a reading of General Relativity's clock rate outside a resting mass, with capacity $c_0^4/2G$; what in the Ocean sets the capacity is the strength of gravity, below. The outward spreading of the record is measured: the binary pulsar's orbit decays at 0.9983 ± 0.0016 of the rate outward spreading gives, where inward spreading would reverse its sign.

The Ocean's own waves. Ripples of the settled amount and phase (Section 11.4) come in exactly two kinds. One is a sound wave with no lowest energy, the wave every medium carries when its settled state gives up a sameness of its rules (here, the freedom to shift the phase everywhere at once); it travels at $c_s^2 = (1+a)/(1+3a)$. The other is a heavy ripple of the amount, whose lowest energy is the cost of a dent. With the dent near the Planck energy (below), $a \lesssim 10^{-120}$: the sound wave runs at the one speed to one part in 10^{120} or better, and the heavy ripple is far beyond reach. Together with the re-made state's two stretch-and-squeeze waves, that is all the Ocean's own part carries. *Light is not among them.* Light is a wave of the plexus. Two requirements follow. The sound wave must barely touch patterns, or it would act as a second long-range force; Cassini's light-timing measurement [54] limits the coupling to about 10^{-5} in the square, and in the minimal law a ripple in the amount at rest carries no Ocean charge, so the coupling vanishes at leading order. And the settled Ocean must carry no charge under the plexus: a medium that did would act as a superconductor for it and give light a mass.

The strength of gravity. If the plexus is a sea of waves, its resistance to being bent has Einstein's form, and its strength is set by how many kinds of wave it carries and by their shortest wavelength [32]:

$$\frac{1}{G} \;\simeq\; -\frac{1}{2\pi}\,\mathrm{str}[k_1]\,\kappa^2, \tag{37}$$

where κ is the shortest-wave cutoff, here the cost of a dent, and $\mathrm{str}[k_1]$ counts each kind of wave with a weight fixed by its type (boson-like kinds with one sign, fermion-like kinds with the other). Gravity attracts only if $\mathrm{str}[k_1] < 0$. Counting the kinds the rebuild builds (Section 12.8), the twelve force carriers and the field the dynamo sustains to give mass, and taking that field not to respond to the overall scale of the rulers, gives $\mathrm{str}[k_1] = -\frac{1}{2}$[1]: gravity attracts, and the dent lies at about 3.5 times the Planck energy, about $4 \times 10^{28}\,\mathrm{eV}$ (a one-loop estimate; factors of order one depend on how the cutoff is taken). The Ocean's own vast energy does not gravitate, because nothing is outside it to press on it. The dent is the resonator, and the Planck scale is the dent seen through gravity, not a second scale.

Health. The Ocean's own part has no ghost and no imaginary speed, and its sound runs below c_s. The re-made state's first-order law has exactly two waves; any departure from it adds four more, which binary pulsars rule out. Beyond first order the law inherits Einstein's form, provided the plexus's grain picks out no frame, which the cosmic-ray bounds already require (Section 11.1). The plexus's own waves, those of a driven superfluid, are not yet checked.

12.8 The family the rebuild allows

The standard model organizes what is measured into a table of sixteen pieces per family. The table is bookkeeping; what the Ocean must reproduce is the measurements. Atoms are measured neutral to extraordinary precision; charge inside the proton and neutron comes in thirds while everything free comes in whole units; the width of the Z counts three kinds of neutrino; the Sun burns by changing one version into the other; and that change happens only to left-twisted patterns. The rebuild's books reproduce these. When a version changes, what is laid down must carry one unit of balance and leave the counts of builds unchanged; no single piece can, and the cheapest pair that can is the electron with an antineutrino, so the partner is forced. The balance the plexus must keep across the Ocean then sets the charges: the two versions differ by exactly one electron's worth, split as $+\frac{2}{3}$ and $-\frac{1}{3}$, with the neutrino at zero. The thirds are a known consequence of balance conditions of this kind; what the rebuild adds is a reason for the inputs.

Of the sixteen settings the table allows, one carries nothing and takes part in nothing: the right-handed neutrino in the standard model's terms. The rebuild does not build it. A build with nothing in it is not a build; the mathematics allows it, the Ocean does not make it. Equation (37) gives a second reason: with sixteen built per family, the known kinds give no gravity at all. Fifteen are built.

Two things the table records are not yet given by the rebuild: why the three-fold mixing of the strong force has three parts (they cannot be directions in space: three directions would give three carriers where eight are measured, and the color factors measured at LEP rule it out), and why there are three families.

[1] Weights k_1 summed over each kind's states, as tabulated in [32]: a massless carrier $-\frac{2}{3}$, a left-handed (Weyl) build $-\frac{1}{6}$, a scalar not responding to the rulers' scale 0; fermion-like kinds enter str with the opposite sign. Twelve carriers give -8; three families of fifteen builds give $+\frac{15}{2}$; the four states of the field that gives mass give 0. With sixteen builds per family the total is 0. Other conventions (per state, or with gauge-fixing modes counted separately) rescale the weights and the cutoff together, so the sign is what the comparison tests.

12.9 The committed difference

With the empty setting not built, the neutrino has no separate partner to join for its measured mass, and joins its own reclaiming. A rare change must then occur: two versions changed at once, two electrons laid down and no antineutrino, the antineutrino of one change taken up as its own reclaiming by the other. This is neutrinoless double-beta decay. The present limit is a half-life above 3.8×10^{26} years in ^{136}Xe, an effective neutrino mass below 28–122 meV [69]. If the neutrino masses are ordered so that the rate cannot be small, the next generation of searches must see it. If the empty setting were built, it would not occur.

End of §12 · token ca7ea3

13 Open Items

Everything the framework leaves open is gathered here. The research program that follows (Section 14) sets out the work that addresses it; the deficiencies against current practice and measurement are in Section 10.

13.1 What the law does not yet say

Which waves the dynamo makes, and so the kind, the version and the extent of a pattern: this is where the three parts of the strong force's mixing and the three families must come from, from the Ocean's own kinds of movement (its flow, its rivers and eddies, its own ripples) rather than from a table. The size of the matter that remained at the start of the beat. The plexus's own waves and their health. And these, stated elsewhere and gathered here:

- *Flow versus build.* How the Ocean's behaviors divide between its flow and the build; most are likely to be a bit of both. Several sections use the flow picture: gravity as a current slowed by drag (Section 11.6), against the uneven making of time (Section 12.2) and a law in which nothing feels drag; expansion and contraction as divergent and convergent flow (Section 5.1); the inflow and flow-through of black holes (Sections 8.1 and 11.9); the geometry built from the flow (Section 11.3); patterns carried by currents (Sections 11.2 and 8.2); the larger current that sets a galaxy's eddy (Section 6); the phase's gap that galaxies are asked to bound (Section 13.2, item (5)); and the same reading in the summary (Section 15).
- *The phase rate today.* What sets it. The slow rate used for the Ocean's own waves (Section 12.7), for the k^4 term (Section 13.2, item (3)), against quantized lines (Section 11.8) and for matter and antimatter (Section 10) is not derived; stiffness alone allows it up to about 10^{17} eV.
- *The eddy's tilt.* How the larger current puts it in place, since the first-order law takes tilt only from the momentum of patterns, and how patterns read it, Equation (15), which is assumed (Section 6).
- *The viscous stress.* Whether $\Pi_{\mu\nu}$ has any role (Section 2.3).
- *The turn.* What turns a beat's collapse into the next beat (Section 5.2).

13.2 Matter and its threshold

(1) **The gaps.** Each kind's lowest energy must be derived: the dent's from the Ocean's own law, and each plexus kind's from the dynamo, with the electron's 0.511 MeV from the strength of the field the dynamo sustains (Section 14, "The cost of a dent").

(2) **Stability.** Require $L_{,X} > 0$ and $\mathscr{K} = L_{,X} + 2XL_{,XX} > 0$; otherwise $c_s^2 < 0$ (gradient instability) or the kinetic term flips sign (ghost).

(3) **Lorentz-violation bounds.** Under the one-speed principle (Section 11.1) c_s is the local c_0 by construction, and the test becomes whether light and matter share it; their speed limits agree to

about one part in 10^{23} (Section 11.4). The k^4 term of the dispersion relation gave short-wavelength excitations a group velocity above c_s, which the one-speed principle does not allow for anything that carries a signal. With the law written on the Ocean's own rulers and clocks the Ocean must be stiff, and in the stiff Ocean the coefficient of the term falls with the shortfall of speed, in the minimal law to about $4a^2 \lesssim 4 \times 10^{-46}$ (Section 11.1), so it carries no signal at any accessible energy. With the dent near the Planck energy it is far smaller still. Photons from distant gamma-ray bursts show no energy-dependent arrival times, and the cosmic-ray bounds place any such effect beyond the Planck scale (Section 11.1). The matter–antimatter comparisons of Section 10, deficiency (5), are among the most precise of these tests.

(4) **Varying-m_e bounds.** Quasar absorption spectra constrain drift in m_p/m_e at roughly the 10^{-5}–10^{-6} level over cosmological time; the Oklo natural reactor and atomic clocks constrain it locally. The electron's mass is a constant of the beat (Section 7, Consequences), and the bounds confirm it.

(5) **The scale hierarchy.** Galactic-scale effects act across kiloparsecs, while the Ocean's own scale, the dent's width, lies near the Planck length. Read as properties of the medium, the two would differ by more than fifty orders of magnitude. They belong to different parts of the Ocean: the phase, whose potential vanishes by symmetry, and the amount, whose gap is the dent.

Route. On the single-structure reading of Section 6 the galactic scale is a property of the individual galactic solution, not of the medium, and the requirement on the medium is an upper bound on the phase's gap,

$$\Omega_{\text{gap}} \lesssim 10^{-26}\,\text{eV}, \tag{38}$$

which $\Omega_{\text{gap}} = 0$ satisfies. A vanishing gap is protected, rather than tuned, because the long-wavelength excitation is the phase θ of the complex field, whose $U(1)$ shift symmetry (5) forbids it a potential. The medium then carries one intrinsic length, the dent's width, and the size of a galaxy is the size of a solution, as a hurricane's is a property of the storm and not of air. The route carries three obligations: to show that the symmetry survives the couplings of equation (1); to say what fixes the size of a galactic solution; and to pass the vertical-acceleration test in the form stated in Section 6. Because the medium then carries a length and no intrinsic acceleration, the acceleration scale $g_\dagger$ of the radial acceleration relation [10] must arise elsewhere, most plausibly from the cosmological background state.

The scale gap has a twin. The strength of gravity at the electron's scale is the measured dimensionless ratio

$$\alpha_G \equiv \frac{G m_e^2}{\hbar c_0} = \left(\frac{\ell_P}{\xi_e}\right)^2 \approx 1.75 \times 10^{-45}, \tag{39}$$

against $\alpha \approx 1/137$ for electromagnetism. Written as a length, it places the Planck length $\ell_P = \sqrt{\hbar G/c_0^3} \approx 1.6 \times 10^{-35}$ m about 10^{22} below ξ_e. Here the Planck length is not a second scale: it is the dent's width seen through gravity (Equation (37)). Gravity is weak at the electron's scale because the electron's lowest energy sits nearly 10^{23} below the dent. The twin is therefore a question about the dynamo: why the field strength it sustains runs so far below the resonator that tunes it. This is the hierarchy problem in the Ocean's form.

(6) **Relation to the Standard Model.** The Higgs mechanism accounts for m_e through a Yukawa coupling to a background field. In this framework that background is the field the dynamo sustains, and the coupling is how strongly the electron's kind reads it. The framework must show that the dynamo supplies the field's measured strength, rather than taking it from the Standard Model.

(7) **Size and spin of the localized excitation.** ξ_e is the scale below which the electron localizes, not its radius. Quantum electrodynamics has the same separation: the electron's reduced Compton wavelength limits how far it can be localized without pair creation, while scattering and its magnetic moment show no internal structure down to about 10^{-18} m, several hundred thousand times smaller.

The framework must show that its localized excitation responds as a point to probes at that resolution.

Separately, the electron has spin $\frac{1}{2}$. Excitations of a scalar field, real or complex, carry integer spin. In this framework each unit of the Ocean carries the direction in which time is made and a hand (Section 12.7); a direction of travel together with an angle around it, counted as a half-angle, is the geometric content of a two-component spinor, which returns to itself only after two full turns. The Ocean's amount and phase stay bosonic, and the spin-$\frac{1}{2}$ order lives in the plexus the dynamo makes, as superfluid helium-3 A shows a medium can support [28]. Light's two transverse polarizations and the two tensor polarizations of gravitational waves are carried by the plexus and the re-made state (Section 11.10). That the dynamo makes exactly this order is not derived.

(8) **Matter and antimatter are not told apart by rotation.** A localized excitation and its antimatter partner are the same pattern, built and reclaimed (Section 12.4). Rotation acts on the three arrows' directions in space, not on whether time is being made or unmade, so no rotation turns matter into antimatter, as observation requires. No second internal phase is needed for this.

(9) **The conserved sense is shared across kinds.** Total lepton number has never been observed to change, while in beta decay an electron appears with an antineutrino. In the rebuild both are laid down together because the books require it (Section 12.8), and both are patterns on plexus waves, so whatever plays the role of lepton number passes between kinds as the books pass. Neutrinoless double-beta decay, which the framework requires (Section 12.9), would be the first change in it, by two units in one rare change.

End of §13 · token ad7aa2

14 Research Program and Testable Predictions

The items are in dependency order: each group needs the groups before it, and the items within a group can be worked on side by side. Items marked *(decisive)* could end the framework as written.

0. What the Ocean is made of.

- **Order parameter.** Show that the plexus the dynamo makes carries the direction-and-angle order that gives spin $\frac{1}{2}$ (Section 13.2, item (7)) and the transverse waves of light, with the Ocean's own amount and phase bosonic; the tensor polarizations of gravitational waves are the re-made state's (Section 11.10). Whether the dynamo can make a fermion-like order from a bosonic flow is the question; laboratory systems show that bosonic units can give fermion-like waves, but not yet the observed family.

1. The law.

- **Flow versus build.** Settle how the Ocean's behaviors divide between its flow and the build, most likely a bit of both (Section 13.1). In particular, show whether the current through time of Section 11.6 is an effective reading of the uneven making of time (Section 12.2) with no dissipative drag, and revise every passage listed there to match.
- **Write the one-speed law.** Derive the field equations from a covariant action rather than introducing stress terms phenomenologically: write the Ocean's law out in full on its own field of rulers and clocks, with no fixed speed anywhere, and derive how the Ocean's state sets those rulers and clocks (Sections 11.1 and 11.4). Show that scalar, gravitational, and electromagnetic perturbations all travel at the one carrying speed $c_s(\phi)$, as Section 11.1 requires: the refractive reading of $c(\phi)$ is adopted, and what remains is to derive it. The principle and its consequences for stiffness are in Section 11.1; the geometry around a non-spinning mass follows from two principles (Section 11.7);

the first-order law follows uniquely from one speed and preservation, and beyond first order the plexus gives Einstein's form (Section 12.7); deriving that form in full from the plexus's own waves remains. Volovik's helium-3 work is the standard example of a medium whose excitations share one emergent speed [28].

- **The waves the dynamo makes.** The law of Section 12.7 states what each pattern carries and how it is rebuilt, but not which waves the dynamo makes. From the Ocean's own kinds of movement (its flow, its rivers and eddies, its own ripples), derive the kinds of plexus wave, and with them a pattern's kind, version and extent. This is where the three parts of the strong force's mixing and the three families must come from. Check that the count reproduces the measured family with fifteen built per family, and that Equation (37) keeps the sign that makes gravity attract.
- **The plexus's own waves.** Derive the spectrum of the driven plexus, show it has the frame-free form (energy per wave proportional to its frequency) the absence of drag requires, and check its health.
- **The leftover.** From the rebuild, derive the matter remaining at the start of the beat, a baryon-to-photon ratio of 6.1×10^{-10} [68].
- **Neutrinoless double-beta decay** *(decisive).* The law requires it (Section 12.9). If the neutrino masses are shown to be ordered so that its rate cannot be small, and the next generation of searches does not see it, the account of the family in Section 12.8 fails.
- **Determine the sign of λ** *(decisive)* from the chosen $L(\phi, X)$ and verify the defocusing condition of Section 9.1. A framework whose potential turns out to be focusing about the present vacuum has no mechanism arresting concentration and is not viable as written. Map the states in which stiffness is lost, determine what the medium does there, and confront the result with domain-wall, microwave-background and relic-abundance bounds.

2. What the law gives.

- **Derive gravity from the medium** *(decisive).* Derive the effective geometry of Equation (1) from the dynamics of the medium, and show that the refractive account of Section 2.5 meets its three decisive requirements: every excitation responds in exactly the same proportion; time is slowed and space stretched in equal measure, giving the observed deflection of starlight; and gravitational waves travel at the speed of light. Then calculate the weak-field and post-Newtonian limits to verify the Solar-System observations, and extend the account to perihelion advance, frame dragging and gravitational radiation, or identify where it fails. Derive the principle of Section 5.1: a gradient of state refracts without net flow, and a uniform change of state produces divergent or convergent flow. Derive Equation (1) with the background defining the metric and only excitations as sources (Section 11.3), under the one-speed law, including how $P(\phi)$ enters c_s and the conservation conditions that follow.
- **Derive the candidate mechanism's inputs.** The current through time (Sections 11.6 and 11.7) needs two inputs: the first-order lockstep constant, equivalent to $\gamma = 1$, and the strength of the drag, which in that picture sets G; Equation (37) sets it instead from the plexus's kinds of wave (the next item), and how the two readings relate is the flow-versus-build question below. Derive the first under the one-speed law; meet the bounds $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ and $\beta - 1 = (-4.1 \pm 7.8) \times 10^{-5}$ [54]; and show that the Ocean's second state variable keeps its cosmological pressure from being a function of density alone, as the exclusion of single-density unified fluids requires [58].
- **The strength of gravity** *(decisive).* Equation (37) gives G from the plexus's kinds of wave and the dent, with the sign that makes gravity attract only for fifteen builds per family and a field giving mass that does not respond to the overall scale of the rulers (Section 12.7). Make the count exact: fix the order-one factors, derive from the dynamo that no further kinds exist (or count them), and show that the result is G's measured value, equivalently α_G of (39). The twin is the electron's

lowest energy, nearly 10^{23} below the dent (Section 13.2, item (5)). A separate numerical relation stands beside this: the relation of Section 10.3 gives the strength of gravity from the electron mass and the cosmological constant, $G = \alpha^3\hbar^2\sqrt{\Lambda}/m_e^3$, which with the measured Λ comes within about ten per cent of the measured G, and within a few per cent if the locally measured expansion rate is used. Determine whether the Ocean produces it or whether it is a numerical coincidence of the present epoch, and trace it to the source that first derived it.

- **Frame dragging.** Section 11.8 gives it to first order in spin, carried by the re-made tilt, on two assumptions (the nearest precedent lacks it [37]), and shows the phase needs no whirl even at a black hole spinning near its limit. Derive Equation (15); show that the surroundings are re-made with a traveling pattern exactly below the critical speed; and extend the rule beyond first order, where the outside of a spinning hole must depend only on its mass and spin.
- **Travel through the surroundings** *(decisive).* Determine the flow regimes of a disturbance traveling relative to the medium, including the superfluid critical speed. Travel through the Ocean below that speed is frictionless and leaves no trace; the microwave-background dipole shows the solar system traveling at about $370\,\mathrm{km\,s^{-1}}$ and the Milky Way at about $600\,\mathrm{km\,s^{-1}}$ relative to the frame in which that background is uniform, the natural candidate for the Ocean's local rest frame, and precision tests see nothing at those speeds, so the framework must produce a critical speed well above them. Determine whether the Ocean's rest frame is that frame. Test for residual wake offsets relative to the direction of travel; for residuals of the radial acceleration relation that track speed through the environment, a property not among those tested in [9]; and against colliding clusters.
- **The cost of a dent.** Derive the gap of the Ocean's amount, and with it the healing length of (20), from the Ocean's own law. In the stiff Ocean the gap is $\sqrt{\lambda}\,v$ and the healing length its inverse, $\xi = \hbar c_0/E_{\mathrm{dent}}$. The strength of gravity places the dent near the Planck energy (Equation (37)), so the healing length lies near the Planck length; it is not the electron's 386 femtometers, which is the electron's own localization length, set by the dynamo (Section 7). Derive the electron's 0.511 MeV, and each kind's lowest energy, from the field the dynamo sustains and the dent's tuning of it.
- **Reproduce the measured vacuum** *(decisive).* Its strengths would also fix the strength of the field the dynamo sustains, separately from the dent that tunes it. Compute the plexus's contribution to vacuum polarization, and hence to the Lamb shift and the anomalous magnetic moment, together with light-by-light scattering, vacuum birefringence and the Casimir force, and compare with quantum electrodynamics and with measurement (deficiency (4) of Section 10).
- **The beat's constant.** Determine how the dent tunes the dynamo so that the electron's mass is fixed for the beat while the flow that powers it falls (Section 7, Consequences), and what at a beat transition sets the tuning for the next beat.
- **Size of the localized excitation.** Compute the probe response of the localized excitation and show it is pointlike to about $10^{-18}\,\mathrm{m}$, and determine how the field that sets its lowest energy behaves at high concentration, consistently with the collapse floor. The structure that supplies spin $\frac{1}{2}$ is the order parameter's (group 0).
- **Antimatter and circulation sense.** Antimatter is the reclaiming (Section 12.4). Derive from the rebuild that partners have identical mass and magnetic moment and opposite electric charge (Section 10, deficiency (5)), and confront any predicted CPT and Lorentz violation with the limits there. Show that lepton number passes between kinds as the books of the rebuild require (Section 13.2, item (9)), and that the long-range field electric charge requires is the plexus's light.
- **Hearing the medium.** If the gapless phonon of the phase (Section 2.2) couples to matter, it implies a weak long-range force or an energy-loss channel. Compare with torsion-balance and atom-interferometer bounds.

3. The dissipative sector.

- **Whether $\Pi_{\mu\nu}$ has a role.** In the settled Ocean nothing is dissipated and $\Pi_{\mu\nu}$ carries nothing (Section 2.3). Determine whether it has any role, as at the beat's turn, and if so what carries it. A carrier that needs a normal component, an excitation gas distinct from the condensate, brings a second sound with a speed of its own [50], which must carry no signal patterns can use or must share c_s.
- **Reformulate the dissipative sector causally.** If $\Pi_{\mu\nu}$ has a role, replace the first-order constitutive relation of Section 2.3 with a second-order Israel–Stewart or BDNK form, and verify that the resulting system is causal and stable in the weak-field limit.
- **Establish whether matter coupling constitutes a forcing term.** The known finite-time blowup constructions for viscous media require an external force injecting momentum; the unforced problem remains open. In this framework the candidate forcing is the matter sector acting on the vacuum through $\kappa(\phi)T^{\text{matter}}_{\mu\nu}$ in (1). Determine whether that coupling can drive the medium to finite-time concentration, or whether the healing-length cutoff bounds it for any physically admissible matter distribution. This bears on the regularity of the medium.

4. Black holes, galaxies and the largest flows.

- **Black holes.** The geometric and medium horizons coincide (Section 8.1); determine what horizon-scale imaging and ringdown measurements constrain beyond the shadow. Explain why a black hole's capacity for state scales with horizon area. Compute the infall asymmetry and drag for a black hole traveling relative to the medium, including the critical-speed regime, and compare with high-velocity merger remnants. For a non-spinning mass the Ocean's geometry is General Relativity's, with the point of no return at $2GM/c_0^2$ and the observed shadow size (Section 11.7); extend it to spinning black holes, which needs frame dragging beyond first order (Section 11.8), and confront the ringdowns of mergers. Whether collapse turns around is a separate question from regularity: stiffness cannot do it, and the candidate is the loosening grip of Section 11.9; derive it, and show that it acts at the core of a black hole as at the end of a beat's contraction. Calculate the turnaround at the core and the time a black hole takes to return what it holds, which depend on what the Ocean is made of (Section 11.9). Revise the stages of Section 8.2 as these results arrive.
- **Resolve or explain the hierarchy** *(decisive)* between the mass gap Ω_{gap} required by galactic dynamics and the Ocean's own scale, the width of a dent near the Planck length; and its twin, the electron's lowest energy nearly 10^{23} below the dent. The candidate route of Section 13.2, item (5), divides the first into three tasks.
  - *Protection.* Write the scalar sector explicitly in the complex field (5), with P, κ, ξ, η, ζ and the quartic potential of Section 9.1 as functions of the modulus or of X, and determine whether the shift symmetry of the phase survives the matter coupling and the dissipative sector. If it does not, compute the gap that results.
  - *Size.* Determine what fixes the extent of a galactic solution, the galaxy and its eddy, given that the medium supplies no kiloparsec scale of its own: what in the current sets an eddy's size and capacity, and where their values originate (Section 6).
  - *Rotation curves without per-galaxy freedom.* Solve the weak-field, stationary, rotating-galaxy limit and derive $v_{\phi,\text{eff}}(r)$ from the baryonic distribution alone, with one parameter set for every galaxy and no independently chosen halo profile, and compare with the observed radial acceleration relation [10, 9], including its universal acceleration scale $g_\dagger \approx 1.2 \times 10^{-10}\,\text{m}\,\text{s}^{-2}$, which the framework must then account for, and extend the derivation to pressure-supported galaxies, which follow the same relation [9]. This must be done jointly with the next item, since a contribution fixed by the baryons is the kind most exposed to it.
- **Radial, not vertical.** A contribution parallel to the baryonic gravitational field, as a scalar-mediated MOND-like force is, enhances radial and vertical accelerations by a common factor and is excluded

by local Milky Way data [11, 12]. The eddy's tilt turns travel across its direction instead (Section 6). Confront the swirl's prediction, about 9 to 15 per cent added to the vertical pull within a kiloparsec of the midplane, together with the rotation curve, with the full Milky Way analysis; and derive the reading of Equation (15), which is assumed. For the swirl of Section 6, test the dependence of the swirl's turning on how a star travels through it against Gaia's velocity data, check the asymmetric deflection of light against galaxy lensing, and find what sets the swirl's strength from a galaxy's baryons. A star traveling against the swirl is turned outward by it, so retrograde and pressure-supported populations, stellar streams and elliptical galaxies test it directly.

- **Distinguish the galactic mechanism from its precedent.** Determine whether the turning of Section 6, carried by the eddy's tilt, is physically distinct from the vortex formation obtained by Huang and collaborators [1] from a nonlinear Klein–Gordon equation with galaxies as external sources, or whether the two are the same mechanism expressed in different formalisms.
- **Simulate a galactic disturbance.** Once the law is written out in full, solve for a galaxy-scale disturbance in its eddy: the tilt the eddy puts in place, its shape near the disk, and the galaxy's wake. Show how the larger current puts the tilt in place, which the first-order law, taking tilt only from the momentum of patterns, does not contain.
- **Flow on the largest scales.** A smooth flow is invisible; only shear, gradients and eddies can be seen. Formulate source-free currents of the Ocean, driven by differences of chemical potential and shaped by their circulation, and determine what a beat transition leaves of them. Within the solar system they are bounded by preferred-frame tests; on large scales they would appear as coherent streams of galaxies that the mass distribution does not account for. Compare the measured large-scale flow of galaxies with the flow predicted from their mass, and test for coherent rotation beyond what the mass predicts.

5. Cosmology and the beats.

- **The heartbeat mechanism.** Derive the effective equation of state $P(\phi)$ and determine whether a stable oscillatory cosmological solution exists. Model the exhaustion of the gradients that drive expansion; show that, as μ falls, the Ocean's state stops driving the divergence of its flow and begins opposing it, so that the divergence reaches zero and turns; model the contraction as the rebuild describes it (Section 12.5), the reclaiming carrying the Ocean everywhere together, and show that it stays smooth. Determine what sets μ today, and whether the fall of μ as the Ocean expands is what ends expansion, with the fall of μ marking a beat's progress and ρ as what resets at the transition (Section 2.2). What turns a contraction around is the loosening grip of the black-hole item (above).
- **Cosmological tests.** Test cosmological expansion, structure growth, gravitational lensing, and background-radiation constraints.
- **The redshift budget.** Show where the energy lost by radiation crossing an expanding region goes, consistently with preservation, and confront any exchange term with the microwave background and light-element abundances (Section 9.3).
- **Contraction-phase contact and the age of the Ocean.** Show that contact during the previous contraction produces the uniformity and fluctuation spectrum of the microwave background, including the baryon acoustic scale; and determine whether the Heartbeat has net expansion, on which the Ocean's having an age depends (Section 5.2; disorder does not accumulate, Section 12.5).
- **The matter excess across a beat.** In the collapse the reclaiming takes everything away and returns the energy (Section 12.5). Show how the next beat's building, with the measured difference between building and reclaiming, leaves the measured excess, and that the reset of disorder holds.

End of §14 · token 873db2

15 Summary of the Revised Formalism

The revised Ocean framework rests on a single premise: there is no empty space. The Ocean is one continuous medium without edges or outside, and everything in it, matter and light included, is a pattern of that medium. Gravity is proposed to be the medium's response to concentrated energy. The mathematical formulation keeps Einstein's equation as the effective description of that response, with a scalar-field stress sector, a viscous constitutive sector that carries nothing in the settled Ocean, a variable pressure contribution and a field-dependent effective coupling as the medium's contributions to it.

The framework extends that idea to the matter sector itself. Each kind of wave the Ocean carries either has a lowest energy or has none. Kinds with one can be held in place, as confined patterns, which are matter; kinds without one, light among them, cannot. The distinction between matter and vacuum then becomes a statement about the Ocean's own waves rather than an imposed energy cut, and the electron's 0.511 MeV is the lowest energy of its kind, set by the field the Ocean's flow sustains.

Three principles frame the account. The Ocean has one speed, at which every disturbance travels, and gravity is where that speed varies from place to place; no speed is fixed. One change of state, in two shapes, makes both gravity and expansion: a gradient of state refracts, and a uniform change of state flows. And the state of the Ocean is never destroyed, only its patterns scrambled; black holes recycle energy and keep no structure.

Beneath all three is the rebuild. Every pattern is built and reclaimed continuously; building makes matter and time, and reclaiming unmakes them and is antimatter. Gravity is the uneven making of time. The law that follows is stated in Section 12, with the one prediction it commits to.

The central hypothesis is that phenomena conventionally divided into dark energy, dark matter, and localized matter may have a common origin in different dynamical regimes of one vacuum field. Establishing that claim requires deriving the relevant limits and demonstrating quantitative agreement with observations.

End of §15 · token ef5186

References

[1] K. Huang, C. Xiong and X. Zhao, *Scalar-field theory of dark matter*, Int. J. Mod. Phys. A **29**, 1450074 (2014); doi:10.1142/S0217751X14500742; arXiv:1304.1595.

[2] L. Berezhiani and J. Khoury, *Theory of dark matter superfluidity*, Phys. Rev. D **92**, 103510 (2015); arXiv:1507.01019.

[3] T. M. Davis and C. H. Lineweaver, *Expanding confusion: common misconceptions of cosmological horizons and the superluminal expansion of the universe*, Publ. Astron. Soc. Aust. **21**, 97 (2004); arXiv:astro-ph/0310808.

[4] A. G. Riess *et al.*, *Observational evidence from supernovae for an accelerating universe and a cosmological constant*, Astron. J. **116**, 1009 (1998); arXiv:astro-ph/9805201.

[5] S. Perlmutter *et al.*, *Measurements of* Ω *and* Λ *from 42 high-redshift supernovae*, Astrophys. J. **517**, 565 (1999); arXiv:astro-ph/9812133.

[6] DESI Collaboration, *DESI DR2 results II: Measurements of baryon acoustic oscillations and cosmological constraints*, arXiv:2503.14738 (2025).

[7] P. J. Steinhardt and N. Turok, *A cyclic model of the universe*, Science **296**, 1436 (2002); doi:10.1126/science.1070462; arXiv:hep-th/0111030.

[8] Planck Collaboration, *Planck 2018 results. X. Constraints on inflation*, Astron. Astrophys. **641**, A10 (2020); doi:10.1051/0004-6361/201833887; arXiv:1807.06211.

[9] F. Lelli, S. S. McGaugh, J. M. Schombert and M. S. Pawlowski, *One law to rule them all: The radial acceleration relation of galaxies*, Astrophys. J. **836**, 152 (2017); doi:10.3847/1538-4357/836/2/152; arXiv:1610.08981.

[10] S. S. McGaugh, F. Lelli and J. M. Schombert, *Radial acceleration relation in rotationally supported galaxies*, Phys. Rev. Lett. **117**, 201101 (2016); arXiv:1609.05917.

[11] M. Lisanti, M. Moschella, N. J. Outmezguine and O. Slone, *Testing dark matter and modifications to gravity using local Milky Way observables*, Phys. Rev. D **100**, 083009 (2019); arXiv:1812.08169.

[12] M. Lisanti, M. Moschella, N. J. Outmezguine and O. Slone, *A preference for cold dark matter over superfluid dark matter in local Milky Way data*, arXiv:1911.12365v2 (2023).

[13] W. Ubachs, J. Bagdonaite, E. J. Salumbides, M. T. Murphy and L. Kaper, *Search for a drifting proton–electron mass ratio from H_2*, Rev. Mod. Phys. **88**, 021003 (2016); arXiv:1511.04476.

[14] R. Lange, N. Huntemann, J. M. Rahm, C. Sanner, H. Shao, B. Lipphardt, Chr. Tamm, S. Weyers and E. Peik, *Improved limits for violations of local position invariance from atomic clock comparisons*, Phys. Rev. Lett. **126**, 011102 (2021); arXiv:2010.06620.

[15] L. Biskupek, J. Müller and J.-M. Torre, *Benefit of new high-precision LLR data for the determination of relativistic parameters*, Universe **7**, 34 (2021); arXiv:2012.12032.

[16] J. Steinhauer, *Observation of quantum Hawking radiation and its entanglement in an analogue black hole*, Nature Phys. **12**, 959 (2016); arXiv:1510.00621.

[17] A. J. S. Hamilton and J. P. Lisle, *The river model of black holes*, Am. J. Phys. **76**, 519 (2008); doi:10.1119/1.2830526; arXiv:gr-qc/0411060.

[18] T. Torres, S. Patrick, A. Coutant, M. Richartz, E. W. Tedford and S. Weinfurtner, *Rotational superradiant scattering in a vortex flow*, Nature Phys. **13**, 833 (2017); doi:10.1038/nphys4151; arXiv:1612.06180.

[19] W. G. Unruh, *Experimental black-hole evaporation?*, Phys. Rev. Lett. **46**, 1351 (1981); doi:10.1103/PhysRevLett.46.1351.

[20] D. N. Page, *Particle emission rates from a black hole. II. Massless particles from a rotating hole*, Phys. Rev. D **14**, 3260 (1976); doi:10.1103/PhysRevD.14.3260.

[21] Ya. B. Zel'dovich, *Amplification of cylindrical electromagnetic waves reflected from a rotating body*, Zh. Eksp. Teor. Fiz. **62**, 2076 (1972) [Sov. Phys. JETP **35**, 1085 (1972)].

[22] S. B. Giddings, *Hawking radiation, the Stefan–Boltzmann law, and unitarization*, Phys. Lett. B **754**, 39 (2016); arXiv:1511.08221.

[23] D. N. Page, *Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole*, Phys. Rev. D **13**, 198 (1976).

[24] R. D. Blandford and R. L. Znajek, *Electromagnetic extraction of energy from Kerr black holes*, Mon. Not. R. Astron. Soc. **179**, 433 (1977); doi:10.1093/mnras/179.3.433.

[25] M. Su, T. R. Slatyer and D. P. Finkbeiner, *Giant gamma-ray bubbles from Fermi-LAT: AGN activity or bipolar Galactic wind?*, Astrophys. J. **724**, 1044 (2010); arXiv:1005.5480.

[26] Event Horizon Telescope Collaboration, *First Sagittarius A* Event Horizon Telescope results. VI. Testing the black hole metric*, Astrophys. J. Lett. **930**, L17 (2022); doi:10.3847/2041-8213/ac6756; arXiv:2311.09484.

[27] E. L. Hahn, *Spin echoes*, Phys. Rev. **80**, 580 (1950); doi:10.1103/PhysRev.80.580.

[28] G. E. Volovik, *The Universe in a Helium Droplet*, International Series of Monographs on Physics **117**, Clarendon Press / Oxford University Press (2003).

[29] K. G. Zloshchastiev, *An alternative to dark matter and dark energy: Scale-dependent gravity in superfluid vacuum theory*, Universe **6**, 180 (2020); doi:10.3390/universe6100180; arXiv:2011.12565.

[30] X. Dou and X.-H. Meng, *Bulk viscous cosmology: unified dark matter*, Advances in Astronomy **2011**, 829340 (2011); arXiv:1012.3045.

[31] P.-H. Chavanis, *Polytropes, logotropes, the universal value of the surface density of dark matter halos, and the value of the cosmological constant*, arXiv:2603.21302.

[32] M. Visser, *Sakharov's induced gravity: a modern perspective*, Mod. Phys. Lett. A **17**, 977 (2002); doi:10.1142/S0217732302006886; arXiv:gr-qc/0204062.

[33] L. Boyle, K. Finn and N. Turok, *CPT-symmetric universe*, Phys. Rev. Lett. **121**, 251301 (2018); arXiv:1803.08928.

[34] M. Arminjon, *Gravity as Archimedes' thrust and a bifurcation in that theory*, Found. Phys. **34**, 1703 (2004); arXiv:physics/0404103.

[35] M. Arminjon, *Space isotropy and weak equivalence principle in a scalar theory of gravity*, Braz. J. Phys. **36**, 177 (2006); arXiv:gr-qc/0412085.

[36] M. Arminjon, *Ether theory of gravitation: why and how?*, in M. C. Duffy and J. Levy (eds.), *Ether, Space-Time and Cosmology*, Vol. 1 (PD Publications, Liverpool, 2008), pp. 139–201; arXiv:gr-qc/0401021.

[37] M. Arminjon, *Continuum dynamics and the electromagnetic field in the scalar ether theory of gravitation*, Open Phys. **14**, 395 (2016); arXiv:1612.07996.

[38] J. Broekaert, *A modified Lorentz-transformation based gravity model confirming basic GRT experiments*, Found. Phys. **35**, 839 (2005); arXiv:gr-qc/0309023.

[39] W. A. Hiscock and L. Lindblom, *Generic instabilities in first-order dissipative relativistic fluid theories*, Phys. Rev. D **31**, 725 (1985).

[40] W. Israel and J. M. Stewart, *Transient relativistic thermodynamics and kinetic theory*, Annals of Physics **118**, 341 (1979).

[41] ATLAS Collaboration, *Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC*, Nature Phys. **13**, 852 (2017); doi:10.1038/nphys4208; arXiv:1702.01625.

[42] R. P. Mignani, V. Testa, D. González Caniulef, R. Taverna, R. Turolla, S. Zane and K. Wu, *Evidence for vacuum birefringence from the first optical-polarimetry measurement of the isolated neutron star RX J1856.5−3754*, Mon. Not. R. Astron. Soc. **465**, 492 (2017); arXiv:1610.08323.

[43] C. D. Anderson, *The positive electron*, Phys. Rev. **43**, 491 (1933).

[44] C. J. Baker *et al.* (ALPHA Collaboration), *Laser cooling of antihydrogen atoms*, Nature **592**, 35 (2021); doi:10.1038/s41586-021-03289-6.

[45] M. Ahmadi *et al.* (ALPHA Collaboration), *Characterization of the 1S–2S transition in antihydrogen*, Nature **557**, 71 (2018); doi:10.1038/s41586-018-0017-2.

[46] M. J. Borchert *et al.* (BASE Collaboration), *A 16-parts-per-trillion measurement of the antiproton-to-proton charge–mass ratio*, Nature **601**, 53 (2022); doi:10.1038/s41586-021-04203-w.

[47] E. K. Anderson *et al.* (ALPHA Collaboration), *Observation of the effect of gravity on the motion of antimatter*, Nature **621**, 716 (2023); doi:10.1038/s41586-023-06527-1.

[48] LHCb Collaboration, *Observation of charge–parity symmetry breaking in baryon decays*, Nature **643**, 1223 (2025); arXiv:2503.16954.

[49] P. Touboul *et al.* (MICROSCOPE Collaboration), *MICROSCOPE mission: final results of the test of the equivalence principle*, Phys. Rev. Lett. **129**, 121102 (2022); arXiv:2209.15487.

[50] M. Visser, *Acoustic black holes: horizons, ergospheres, and Hawking radiation*, Class. Quantum Grav. **15**, 1767 (1998); arXiv:gr-qc/9712010.

[51] E. Babichev, V. Mukhanov and A. Vikman, *k-Essence, superluminal propagation, causality and emergent geometry*, JHEP **0802**, 101 (2008); arXiv:0708.0561.

[52] S. Coleman and S. L. Glashow, *High-energy tests of Lorentz invariance*, Phys. Rev. D **59**, 116008 (1999); arXiv:hep-ph/9812418.

[53] F. W. Stecker and S. L. Glashow, *New tests of Lorentz invariance following from observations of the highest energy cosmic gamma rays*, Astropart. Phys. **16**, 97 (2001); arXiv:astro-ph/0102226.

[54] C. M. Will, *The confrontation between general relativity and experiment*, Living Rev. Relativ. **17**, 4 (2014); arXiv:1403.7377.

[55] O. Klein, *On the thermodynamical equilibrium of fluids in gravitational fields*, Rev. Mod. Phys. **21**, 531 (1949); doi:10.1103/RevModPhys.21.531.

[56] A. Kamenshchik, U. Moschella and V. Pasquier, *An alternative to quintessence*, Phys. Lett. B **511**, 265 (2001); arXiv:gr-qc/0103004.

[57] M. C. Bento, O. Bertolami and A. A. Sen, *Generalized Chaplygin gas, accelerated expansion and dark energy–matter unification*, Phys. Rev. D **66**, 043507 (2002); arXiv:gr-qc/0202064.

[58] H. B. Sandvik, M. Tegmark, M. Zaldarriaga and I. Waga, *The end of unified dark matter?*, Phys. Rev. D **69**, 123524 (2004); arXiv:astro-ph/0212114.

[59] C. W. F. Everitt *et al.*, *Gravity Probe B: Final results of a space experiment to test general relativity*, Phys. Rev. Lett. **106**, 221101 (2011); arXiv:1105.3456.

[60] I. Ciufolini, A. Paolozzi, E. C. Pavlis, G. Sindoni, J. Ries, R. Matzner, R. Koenig, C. Paris, V. Gurzadyan and R. Penrose, *An improved test of the general relativistic effect of frame-dragging using the LARES and LAGEOS satellites*, Eur. Phys. J. C **79**, 872 (2019); arXiv:1910.09908.

[61] I. Ciufolini, A. Paolozzi, E. C. Pavlis, J. C. Ries, C. Paris, E. Ortore, R. Matzner, M. Kuzmicz-Cieslak, D. Deka, D. E. Pavlis, P. Schreiner, W.-T. Ni, R. Penrose and V. Gurzadyan, *LARES-2 satellite measures frame-dragging effect around the Earth*, Nature **655**, 332 (2026); doi:10.1038/s41586-026-10715-0.

[62] J. G. Williams, S. G. Turyshev and D. H. Boggs, *Progress in lunar laser ranging tests of relativistic gravity*, Phys. Rev. Lett. **93**, 261101 (2004); doi:10.1103/PhysRevLett.93.261101; arXiv:gr-qc/0411113.

[63] N. Andersson and G. L. Comer, *Relativistic fluid dynamics: physics for many different scales*, Living Rev. Relativ. **24**, 3 (2021); doi:10.1007/s41114-021-00031-6; arXiv:2008.12069.

[64] G. E. Volovik, *Superfluid analogies of cosmological phenomena*, Phys. Rep. **351**, 195 (2001); doi:10.1016/S0370-1573(00)00139-3; arXiv:gr-qc/0005091.

[65] A. Garat and R. H. Price, *Nonexistence of conformally flat slices of the Kerr spacetime*, Phys. Rev. D **61**, 124011 (2000); doi:10.1103/PhysRevD.61.124011; arXiv:gr-qc/0002013.

[66] A. M. Archibald *et al.*, *Universality of free fall from the orbital motion of a pulsar in a stellar triple system*, Nature **559**, 73 (2018); arXiv:1807.02059.

[67] M. Agostini *et al.* (Borexino Collaboration), *Comprehensive geoneutrino analysis with Borexino*, Phys. Rev. D **101**, 012009 (2020); arXiv:1909.02257.

[68] B. D. Fields, K. A. Olive, T.-H. Yeh and C. Young, *Big-bang nucleosynthesis after Planck*, J. Cosmol. Astropart. Phys. **03**, 010 (2020); arXiv:1912.01132.

[69] S. Abe *et al.* (KamLAND-Zen Collaboration), *Search for Majorana neutrinos with the complete KamLAND-Zen dataset*, Phys. Rev. Lett. **135**, 262501 (2025); doi:10.1103/jkf6-48j8; arXiv:2406.11438.

Reference basis. Revised from an earlier formalism supplied by the author, which defined the Ocean metric, the vacuum viscosity tensor, the scalar pressure term, the effective coupling $\kappa(\phi)$, a variable-speed-of-light reading of $c(\phi)$, and a 0.511 MeV matter threshold. Two of these are superseded: the threshold by Section 7, and the variable-speed-of-light reading by Section 11.1, where $c(\phi)$ is the Ocean's one carrying speed rather than a second causal speed. The viscosity tensor is kept but carries nothing in the settled Ocean (Section 2.3).

End of References · token 5045e9

Reading manifest

Every part of this document, the page on which it begins and its checksum token. Each token is the first six hexadecimal digits of the SHA-256 hash of that part's LaTeX source; the document digest is the SHA-256 of the document name and all part hashes in order. The same digest is given in the repository README and the archive record, so a reading can be checked without the document.

Part	Title	Page	Token
Preface	Abstract, reader's note and preface	1	`a70279`
Glossary	Conventions and glossary	8	`692b89`
§1	The Field Equation	10	`c6c077`
§2	Tensor Definitions	11	`dd4ba8`
§3	Scalar-Field Dynamics	14	`d37c4e`
§4	The Ocean Interpretation	14	`bf999b`
§5	Cosmological Heartbeat	15	`cd0a4a`
§6	Galactic Dynamics	17	`824d10`
§7	The Localization Threshold	19	`f72705`
§8	Limiting Regimes	22	`e2cf88`
§9	Conservation and Consistency Conditions	26	`e0ec99`
§10	Relation to Existing Work	29	`b62bd0`
§11	One Speed and the Split	33	`5ab9ee`
§12	The Rebuild and the Law	42	`ca7ea3`
§13	Open Items	48	`ad7aa2`
§14	Research Program and Testable Predictions	50	`873db2`
§15	Summary of the Revised Formalism	55	`ef5186`
References	References and reference basis	55	`5045e9`

Document digest: `ca71-872a-82db`

Version 3.12 · 18 parts · 59 pages

End of document