

# Duality (đối ngẫu)

---

Hoàng Nam Dũng

Khoa Toán - Cơ - Tin học, Đại học Khoa học Tự nhiên, Đại học Quốc gia Hà Nội

# Lagrangian

Consider general minimization problem

$$\begin{array}{ll}\min_x & f(x) \\ \text{subject to} & h_i(x) \leq 0, \ i = 1, \dots, m \\ & l_j(x) = 0, \ j = 1, \dots, r.\end{array}$$

Need not be convex, but of course we will pay special attention to convex case.

We define the Lagrangian as

$$L(x, u, v) = f(x) + \sum_{i=1}^m u_i h_i(x) + \sum_{j=1}^r v_j l_j(x).$$


New variables  $u \in \mathbb{R}^m, v \in \mathbb{R}^r$ , with  $u \geq 0$  (implicitly, we define  $L(x, u, v) = -\infty$  for  $u < 0$ ).

Important property: for any  $u \geq 0$  and  $v$ ,

$$f(x) \geq L(x, u, v) \quad \text{at each feasible } x.$$

Why? For feasible  $x$ ,

$$L(x, u, v) = f(x) + \sum_{i=1}^m u_i \underbrace{h_i(x)}_{\leq 0} + \sum_{j=1}^r v_j \underbrace{l_j(x)}_{=0} \leq f(x).$$



- ▶ Solid line is  $f$ .
- ▶ Dashed line is  $h$ , hence feasible set  $\approx [-0.46, 0.46]$ .
- ▶ Each dotted line shows  $L(x, u, v)$  for different choice of  $u \geq 0$ .

(From B & V page 127).


## Lagrange dual function

Let  $C$  denote primal feasible set,  $f^*$  denote primal optimal value.  
Minimizing  $L(x, u, v)$  over all  $x$  gives a lower bound:

$$f^* \geq \min_{x \in C} L(x, u, v) \geq \min_x L(x, u, v) := g(u, v).$$

We call  $g(u, v)$  the **Lagrange dual function**, and it gives a lower bound on  $f^*$  for any  $u \geq 0$  and  $v$ , called dual feasible  $u, v$ .

- ▶ Dashed horizontal line is  $f^*$ .
  - ▶ Dual variables  $\lambda$  is (our  $u$ ).
  - ▶ Solid line shows  $g(\lambda)$ .
- (From B & V page 127).



## Example: quadratic program

Consider quadratic program:

$$\begin{array}{ll}\min_x & \frac{1}{2}x^T Qx + c^T x \\ \text{subject to} & Ax = b, x \geq 0,\end{array}$$

where  $Q \succ 0$ . Lagrangian:

$$L(x, u, v) = \frac{1}{2}x^T Qx + c^T x - u^T x + v^T (Ax - b).$$

Lagrange dual function:

$$g(u, v) = \min_x L(x, u, v) = -\frac{1}{2}(c - u + A^T v)^T Q^{-1}(c - u + A^T v) - b^T v.$$

For any  $u \geq 0$  and any  $v$ , this is lower a bound on primal optimal value  $f^*$ .

Same problem

$$\begin{array}{ll}\min_x & \frac{1}{2}x^T Qx + c^T x \\ \text{subject to} & Ax = b, x \geq 0,\end{array}$$

but now  $Q \succeq 0$ . Lagrangian:

$$L(x, u, v) = \frac{1}{2}x^T Qx + c^T x - u^T x + v^T (Ax - b).$$

Lagrange dual function:


$$g(u, v) = \begin{cases} -\frac{1}{2}(c - u + A^T v)^T Q^+ (c - u + A^T v) - b^T v & \text{if } c - u + A^T v \perp \text{null}(Q) \\ -\infty & \text{otherwise,} \end{cases}$$

where  $Q^+$  denotes generalized inverse of  $Q$ . For any  $u \geq 0$ ,  $v$ , and  $c - u + A^T v \perp \text{null}(Q)$ ,  $g(u, v)$  is a nontrivial lower bound on  $f^*$ .

## Example: quadratic program in 2D

We choose  $f(x)$  to be quadratic in 2 variables, subject to  $x \geq 0$ .

Dual function  $g(u)$  is also quadratic in 2 variables, also subject to  $u \geq 0$ .



Dual function  $g(u)$  provides a bound on  $f^*$  for every  $u \geq 0$ .

Largest bound this gives us: turns out to be exactly  $f^*$ ... coincidence?

More on this later, via KKT conditions.

## Lagrange dual problem

Given primal problem

$$\begin{array}{ll}\min_x & f(x) \\ \text{subject to} & h_i(x) \leq 0, \quad i = 1, \dots, m \\ & l_j(x) = 0, \quad j = 1, \dots, r.\end{array}$$

Our constructed dual function  $g(u, v)$  satisfies  $f^* \geq g(u, v)$  for all  $u \geq 0$  and  $v$ . Hence best lower bound is given by maximizing  $g(u, v)$  over all dual feasible  $u, v$ , yielding **Lagrange dual problem**:

$$\begin{array}{ll}\max_{u, v} & g(u, v) \\ \text{subject to} & u \geq 0.\end{array}$$

Key property, called **weak duality**: if dual optimal value is  $g^*$ , then

$$f^* \geq g^*.$$

Note that this always holds (even if primal problem is nonconvex).



Another key property: the dual problem is a **convex optimization** problem (as written, it is a concave maximization problem).

Again, this is always true (even when primal problem is not convex).



By definition:

$$\begin{aligned} g(u, v) &= \min_x \left\{ f(x) + \sum_{i=1}^m u_i h_i(x) + \sum_{j=1}^r v_j l_j(x) \right\} \\ &= - \max_x \left\{ \underbrace{-f(x) - \sum_{i=1}^m u_i h_i(x) - \sum_{j=1}^r v_j l_j(x)}_{\text{pointwise maximum of convex functions in } (u, v)} \right\} \end{aligned}$$

I.e.,  $g$  is concave in  $(u, v)$ , and  $u \geq 0$  is a convex constraint, hence dual problem is a concave maximization problem.

## Example: nonconvex quartic minimization

Define  $f(x) = x^4 - 50x^2 + 100x$  (nonconvex), minimize subject to constraints  $x \geq -4.5$ .



Dual function  $g$  can be derived explicitly, via closed-form equation for roots of a cubic equation.

Form of  $g$  is rather complicated:

$$g(u) = \min_{i=1,2,3} \{ F_i^4(u) - 50F_i^2(u) + 100F_i(u) \},$$

where for  $i = 1, 2, 3$ ,

$$F_i(u) = \frac{-a_i}{12 \cdot 2^{1/3}} \left( 432(100 - u) - (432^2(100 - u)^2 - 4 \cdot 1200^3)^{1/2} \right)^{1/3} \\ - 100 \cdot 2^{1/3} \frac{1}{(432(100 - u) - (432^2(100 - u)^2 - 4 \cdot 1200^3)^{1/2})^{1/3}},$$

and  $a_1 = 1, a_2 = (-1 + i\sqrt{3})/2, a_3 = (-1 - i\sqrt{3})/2$ .

Without the context of duality it would be difficult to tell whether or not  $g$  is concave ... but we know it must be!

## Strong duality

Recall that we always have  $f^* \geq g^*$  (weak duality). On the other hand, in some problems we have observed that actually

$$f^* = g^*,$$

which is called **strong duality**.

**Slater's condition**: if the primal is a convex problem (i.e.,  $f$  and  $h_1, \dots, h_m$  are convex,  $\ell_1, \dots, \ell_r$  are affine), and there exists at least one strictly feasible  $x \in \mathbb{R}^n$ , meaning

$$h_1(x) < 0, \dots, h_m(x) < 0 \quad \text{and} \quad \ell_1(x) = 0, \dots, \ell_r(x) = 0,$$

then strong duality holds.

This is a pretty weak condition. An important **refinement**: strict inequalities only need to hold over functions  $h_i$  that are not affine.

## Example: support vector machine dual

Given  $y \in \{-1, 1\}^n$ ,  $X \in \mathbb{R}^{n \times p}$ , rows  $x_1, \dots, x_n$ , recall the **support vector machine** problem:

$$\begin{aligned} \min_{\beta, \beta_0, \xi} \quad & \frac{1}{2} \|\beta\|_2^2 + C \sum_{i=1}^n \xi_i \\ \text{subject to} \quad & \xi_i \geq 0, \quad i = 1, \dots, n \\ & y_i(x_i^T \beta + \beta_0) \geq 1 - \xi_i, \quad i = 1, \dots, n. \end{aligned}$$

Introducing dual variables  $v, w \geq 0$ , we form the Lagrangian:

$$\begin{aligned} L(\beta, \beta_0, \xi, v, w) = & \frac{1}{2} \|\beta\|_2^2 + C \sum_{i=1}^n \xi_i - \sum_{i=1}^n v_i \xi_i + \\ & + \sum_{i=1}^n w_i (1 - \xi_i - y_i(x_i^T \beta + \beta_0)). \end{aligned}$$

Minimizing over  $\beta, \beta_0, \xi$  gives Lagrange dual function:

$$g(v, w) = \begin{cases} -\frac{1}{2}w^T \tilde{X} \tilde{X}^T w + 1^T w & \text{if } w = C1 - v, w^T y = 0 \\ -\infty & \text{otherwise,} \end{cases}$$

where  $\tilde{X} = \text{diag}(y)X$ . Thus SVM dual problem, eliminating slack variable  $v$ , becomes

$$\begin{aligned} \max_w \quad & -\frac{1}{2}w^T \tilde{X} \tilde{X}^T w + 1^T w \\ \text{subject to} \quad & 0 \leq w \leq C1, w^T y = 0. \end{aligned}$$

Check: Slater's condition is satisfied, and we have strong duality. Further, from study of SVMs, might recall that at optimality

$$\beta = \tilde{X}^T w.$$

This is not a coincidence, as we'll later via the KKT conditions.

## Duality gap

Given primal feasible  $x$  and dual feasible  $u, v$ , the quantity

$$f(x) - g(u, v),$$

is called the duality gap between  $x$  and  $u, v$ . Note that



$$f(x) - f^* \leq f(x) - g(u, v),$$

so if the duality gap is zero, then  $x$  is primal optimal (and similarly,  $u, v$  are dual optimal).

From an algorithmic viewpoint, provides a stopping criterion: if  $f(x) - g(u, v) \leq \epsilon$ , then we are guaranteed that  $f(x) - f^* \leq \epsilon$ .

Very useful, especially in conjunction with iterative methods ...  
more dual uses in coming lectures.

## References

-  S. Boyd and L. Vandenberghe (2004), *Convex optimization*, Chapter 5.
-  R. T. Rockafellar (1970), *Convex analysis*, Chapters 28–30.