

Hunter Lybbert
Student ID: 2426454
04-03-25
AMATH 503

HOMework 1

Exercises come from *Introduction to Partial Differential Equations by Peter J. Olver* as well as supplemented by instructor provided exercises.

1: Olver 1.1

Solution:

- (a) $\frac{du}{dx} + xu = 1$: Ordinary equilibrium differential equation of the first order.
- (b) $\frac{\partial u}{\partial t} + u \frac{\partial u}{\partial x} = x$: Partial dynamic differential equation of the first order.
- (c) $u_{tt} = 9u_{xx}$: Partial dynamic differential equation of the second order.
- (d) $\frac{\partial u}{\partial t} = \frac{\partial^2 u}{\partial x^2} + \frac{\partial u}{\partial x}$: Partial dynamic differential equation of the second order.
- (e) $-\frac{\partial^2 u}{\partial x^2} - \frac{\partial^2 u}{\partial y^2} = x^2 + y^2$: Partial equilibrium differential equation of the second order.
- (f) $\frac{\partial^2 u}{\partial t^2} + 3u = \sin t$: Ordinary equilibrium differential equation of the second order.
- (g) $u_{xx} + u_{yy} + u_{zz} + (x^2 + y^2 + z^2)u = 0$: Partial equilibrium differential equation of the second order.
- (h) $u_{xx} = x + u^2$: Ordinary equilibrium differential equation of the second order.
- (i) $\frac{\partial u}{\partial t} + \frac{\partial^3 u}{\partial x^3} + u \frac{\partial u}{\partial x} = 0$: Partial dynamic differential equation of the third order.
- (j) $\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y \partial z} = u$: Partial equilibrium differential equation of the second order.
- (k) $u_{tt} = u_{xxxx} + 2u_{xxyy} + u_{yyyy}$: Partial dynamic differential equation of the fourth order.

□

2: Olver 1.17

Solution:

- (a) $u_t = x^2 u_{xx} + 2xu_x$: homogeneous linear
- (b) $-u_{xx} = u_{yy} = \sin u$: nonlinear
- (c) $u_{xx} + 2yu_{yy} = 3$: inhomogeneous linear
- (d) $u_t + uu_x = 3u$: nonlinear
- (e) $e^y u_x = e^x u_y$: homogeneous linear
- (f) $u_t = 5u_{xxx} + x^2 u + x$: inhomogeneous linear

□

3: Olver 1.22

- (a) Prove that the Laplacian $\Delta = \partial_x^2 + \partial_y^2$ defines a linear differential operator.

Solution: We need to show that for some appropriate functions u, v and two scalars $a, b \in \mathbb{R}$

$$\Delta[au + bv] = a\Delta[u] + b\Delta[v].$$

We will do this directly,

$$\begin{aligned}\Delta[au + bv] &= (\partial_x^2 + \partial_y^2)(au + bv) = (\partial_x^2 + \partial_y^2)au + (\partial_x^2 + \partial_y^2)bv \\ &= \partial_x^2 au + \partial_y^2 au + \partial_x^2 bv + \partial_y^2 bv \\ &= a\partial_x^2 u + a\partial_y^2 u + b\partial_x^2 v + b\partial_y^2 v \\ &= au_{xx} + au_{yy} + bv_{xx} + bv_{yy} \\ &= a(u_{xx} + u_{yy}) + b(v_{xx} + v_{yy}) \\ &= a(\partial_x^2 u + \partial_y^2 u) + b(\partial_x^2 v + \partial_y^2 v) \\ &= a(\partial_x^2 + \partial_y^2)u + b(\partial_x^2 + \partial_y^2)v \\ &= a\Delta[u] + b\Delta[v].\end{aligned}$$

□

- (b) Write out the Laplace equation $\Delta[u] = 0$ and the Poisson equation $-\Delta[u] = f$.

Solution: The Laplace equation is

$$\Delta[u] = (\partial_x^2 + \partial_y^2)u = u_{xx} + u_{yy} = 0$$

and the Poisson equation is

$$-\Delta[u] = -(\partial_x^2 + \partial_y^2)u = -u_{xx} - u_{yy} = f.$$

□

4: We derive the advection-diffusion equation from the microscopic view. Define $u(x, t)$ as the density of the particles at location x and time t . Define the probability of jumping from the left as $p(x - \Delta x \rightarrow x, t) \approx \frac{1}{2} + \Delta x$ when Δx is small, and the probability of jumping from the right as $q(x + \Delta x \rightarrow x, t) \approx \frac{1}{2} - \Delta x$ with small Δx . Assume $D := \lim_{\Delta x, \Delta t \rightarrow 0} \frac{(\Delta x)^2}{\Delta t}$. Establish the equation of $u(x, t)$ in the continuum limit.

Solution:

TODO

5: (a) Consider the following boundary value problem (BVP).

$$\begin{cases} X''(x) + \lambda X = 0, & x \in (0, L) \\ X(0) = X(L) = 0, \end{cases}$$

where $L > 0$ is a constant. Solve the eigenpair:

$$(X_k, \lambda_k) = \left\{ \sin\left(\frac{k\pi x}{L}\right), \left(\frac{k\pi}{L}\right)^2 \right\}_{k=1}^{\infty}$$

Solution:

TODO

(b) Consider the following boundary value problem (BVP).

$$\begin{cases} X''(x) + \lambda X = 0, & x \in (0, L) \\ X'(0) = X'(L) = 0, \end{cases}$$

where $L > 0$ is a constant. Solve the eigenpair:

$$(X_k, \lambda_k) = \left\{ \cos\left(\frac{k\pi x}{L}\right), \left(\frac{k\pi}{L}\right)^2 \right\}_{k=0}^{\infty}$$

Solution:

TODO

6: Consider the following IBVP in a rectangle:

$$\begin{cases} u_t = \Delta u, & (x, y) \in (0, L_1) \times (0, L_2), t > 0 \\ \partial_{\mathbf{n}} u(x, y, t) = 0, & (x, y) \in \partial((0, L_1) \times (0, L_2)), t > 0 \\ u(x, y, 0) = u_0(x, y) \geq 0, \neq 0 & (x, y) \in (0, L_1) \times (0, L_2) \end{cases}$$

where \mathbf{n} denotes the unit outer normal derivative and $L_1, L_2 > 0$ are given constants. Solve to get the general solution. Recall that $\Delta = \partial_{xx} + \partial_{yy}$.

Solution:

TODO