

# §1. Set Theory

Math 4341 (Topology)

# Sets and subsets


- ▶ Informally, we consider a *set* as a “collection of things”. These “things” will be referred to as the *elements* of the set.

# Sets and subsets

- ▶ Informally, we consider a *set* as a “collection of things”. These “things” will be referred to as the *elements* of the set.
- ▶ If  $A$  is a set and  $a$  is an element of  $A$ , we write  $a \in A$ . If  $b$  is not an element of  $A$ , then we write  $b \notin A$ .

# Sets and subsets

- ▶ Informally, we consider a *set* as a “collection of things”. These “things” will be referred to as the *elements* of the set.
- ▶ If  $A$  is a set and  $a$  is an element of  $A$ , we write  $a \in A$ . If  $b$  is not an element of  $A$ , then we write  $b \notin A$ .
- ▶ If  $B$  is another set which contains all the elements of  $A$  (that is, if  $a \in A$  implies that  $a \in B$ ), then we say that  $A$  is a *subset* of  $B$  and write  $A \subset B$ .



# Sets and subsets (contd)

- ▶ We say that two sets  $A$  and  $B$  are equal (and write  $A = B$ ), if they satisfy  $A \subset B$  and  $B \subset A$ .

# Sets and subsets (contd)

- ▶ We say that two sets  $A$  and  $B$  are equal (and write  $A = B$ ), if they satisfy  $A \subset B$  and  $B \subset A$ .
- ▶ If  $A \subset B$  but  $A \neq B$ , we write  $A \subsetneq B$  and say that  $A$  is a *proper subset* of  $B$ .

# Sets and subsets (contd)

- ▶ We say that two sets  $A$  and  $B$  are equal (and write  $A = B$ ), if they satisfy  $A \subset B$  and  $B \subset A$ .
- ▶ If  $A \subset B$  but  $A \neq B$ , we write  $A \subsetneq B$  and say that  $A$  is a *proper subset* of  $B$ .
- ▶ Example 1.1: When a set contains only a few elements, we simply list them. For example if  $A$  contains only  $a$ ,  $b$  and  $c$ , we write  $A = \{a, b, c\}$ . Then if  $B = \{a, b, c, d\}$ , we see that  $A \subset B$  and since  $d \in B$  but  $d \notin A$ , we have  $A \subsetneq B$ .

# Sets and subsets (contd)

- ▶ We say that two sets  $A$  and  $B$  are equal (and write  $A = B$ ), if they satisfy  $A \subset B$  and  $B \subset A$ .
- ▶ If  $A \subset B$  but  $A \neq B$ , we write  $A \subsetneq B$  and say that  $A$  is a *proper subset* of  $B$ .
- ▶ Example 1.1: When a set contains only a few elements, we simply list them. For example if  $A$  contains only  $a$ ,  $b$  and  $c$ , we write  $A = \{a, b, c\}$ . Then if  $B = \{a, b, c, d\}$ , we see that  $A \subset B$  and since  $d \in B$  but  $d \notin A$ , we have  $A \subsetneq B$ .
- ▶ Example 1.2: Sets are often given by the properties of their elements. For example, the set consisting of all odd numbers is written as

$$\{x \mid x \text{ is an odd integer}\},$$

and is read as “ $x$  such that  $x$  is an odd integer”.



# Empty set and power set

- ▶ We also consider the set which contains no elements at all; this set is called *the empty set* and is denoted  $\emptyset$ . Note that  $\emptyset \subset X$  for any set  $X$ . Moreover,  $\emptyset \neq \{\emptyset\}$ .

# Empty set and power set

- ▶ We also consider the set which contains no elements at all; this set is called *the empty set* and is denoted  $\emptyset$ . Note that  $\emptyset \subset X$  for any set  $X$ . Moreover,  $\emptyset \neq \{\emptyset\}$ .
- ▶ Let  $X$  be any set. The *power set* of  $X$ , denoted  $\mathcal{P}(X)$ , is the set of all subsets of  $X$ . That is

$$\mathcal{P}(X) = \{U \mid U \subset X\}.$$

Note that elements of power sets are themselves sets.

# Empty set and power set

- ▶ We also consider the set which contains no elements at all; this set is called *the empty set* and is denoted  $\emptyset$ . Note that  $\emptyset \subset X$  for any set  $X$ . Moreover,  $\emptyset \neq \{\emptyset\}$ .
- ▶ Let  $X$  be any set. The *power set* of  $X$ , denoted  $\mathcal{P}(X)$ , is the set of all subsets of  $X$ . That is

$$\mathcal{P}(X) = \{U \mid U \subset X\}.$$

Note that elements of power sets are themselves sets.

- ▶ Example 1.3: Some examples:

$$\mathcal{P}(\emptyset) = \{\emptyset\},$$

$$\mathcal{P}(\{a\}) = \{\emptyset, \{a\}\},$$

$$\mathcal{P}(\{a, b\}) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\},$$



$$\mathcal{P}(\{a, b, c\}) = \{\emptyset, \{a\}, \{b\}, \{c\}, \{a, b\}, \{b, c\}, \{a, c\}, \{a, b, c\}\}.$$

# Finite union and intersection

- Given two sets  $A$  and  $B$ , we define their *union*  $A \cup B$  and *intersection*  $A \cap B$  by

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\},$$

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}.$$





# Finite union and intersection

- ▶ Given two sets  $A$  and  $B$ , we define their *union*  $A \cup B$  and *intersection*  $A \cap B$  by

$$A \cup B = \{x \mid x \in A \text{ or } x \in B\},$$

$$A \cap B = \{x \mid x \in A \text{ and } x \in B\}.$$




- ▶ The sets  $A$  and  $B$  are called *disjoint* if  $A \cap B = \emptyset$ .

# Difference and complement

- ▶ Given two sets  $A$  and  $B$ , we define the *difference*  $A \setminus B$  by


$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}.$$



# Difference and complement


- ▶ Given two sets  $A$  and  $B$ , we define the *difference*  $A \setminus B$  by

$$A \setminus B = \{x \mid x \in A \text{ and } x \notin B\}.$$



- ▶ If  $A \subset X$ , we define the *complement* of  $A$  in  $X$ , written  $A^c$ , as

$$A^c = X \setminus A.$$



# De Morgan's laws

► Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$



# De Morgan's laws

- ▶ Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- ▶ Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .

# De Morgan's laws

► Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

► Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .

- Let  $a \in A \cap (B \cup C)$ . Then  $a \in A$  and  $a \in B \cup C$ . The latter means that  $a \in B$  or  $a \in C$ . Since  $a \in A$ , we have either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ .

# De Morgan's laws

- ▶ Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- ▶ Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .
  - ▶ Let  $a \in A \cap (B \cup C)$ . Then  $a \in A$  and  $a \in B \cup C$ . The latter means that  $a \in B$  or  $a \in C$ . Since  $a \in A$ , we have either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ .
  - ▶ Written in set notation, this says that  $a \in (A \cap B) \cup (A \cap C)$ .

# De Morgan's laws

- ▶ Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

- ▶ Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .
  - ▶ Let  $a \in A \cap (B \cup C)$ . Then  $a \in A$  and  $a \in B \cup C$ . The latter means that  $a \in B$  or  $a \in C$ . Since  $a \in A$ , we have either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ .
  - ▶ Written in set notation, this says that  $a \in (A \cap B) \cup (A \cap C)$ .
- ▶ We now prove  $(A \cap B) \cup (A \cap C) \subset A \cap (B \cup C)$ .

# De Morgan's laws

► Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

► Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .

► Let  $a \in A \cap (B \cup C)$ . Then  $a \in A$  and  $a \in B \cup C$ . The latter means that  $a \in B$  or  $a \in C$ . Since  $a \in A$ , we have either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ .

► Written in set notation, this says that  $a \in (A \cap B) \cup (A \cap C)$ .

► We now prove  $(A \cap B) \cup (A \cap C) \subset A \cap (B \cup C)$ .

► Let  $a \in (A \cap B) \cup (A \cap C)$ . Then  $a \in A \cap B$  or  $a \in A \cap C$ . This means that either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ . In both cases we have  $a \in A$  and either  $a \in B$  or  $a \in C$ .

# De Morgan's laws

► Proposition 1.1 (De Morgan's laws):

$$(1) \quad A \cap (B \cup C) = (A \cap B) \cup (A \cap C),$$

$$(2) \quad A \cup (B \cap C) = (A \cup B) \cap (A \cup C).$$

► Proof of (1): We first prove  $A \cap (B \cup C) \subset (A \cap B) \cup (A \cap C)$ .

► Let  $a \in A \cap (B \cup C)$ . Then  $a \in A$  and  $a \in B \cup C$ . The latter means that  $a \in B$  or  $a \in C$ . Since  $a \in A$ , we have either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ .

► Written in set notation, this says that  $a \in (A \cap B) \cup (A \cap C)$ .

► We now prove  $(A \cap B) \cup (A \cap C) \subset A \cap (B \cup C)$ .

► Let  $a \in (A \cap B) \cup (A \cap C)$ . Then  $a \in A \cap B$  or  $a \in A \cap C$ . This means that either  $a \in A$  and  $a \in B$ , or  $a \in A$  and  $a \in C$ . In both cases we have  $a \in A$  and either  $a \in B$  or  $a \in C$ .

► Written in set notation, this says that  $a \in A \cap (B \cup C)$ .

# Arbitrary union and intersection

- ▶ Notation:  $\forall$  means “for all”, and  $\exists$  means “there exists”.

# Arbitrary union and intersection

- ▶ Notation:  $\forall$  means “for all”, and  $\exists$  means “there exists”.
- ▶ Let  $I$  be any set. Then a collection  $\{A_i\}_{i \in I}$  of sets  $A_i$  is called a family of sets parametrized by  $I$ .



# Arbitrary union and intersection

- ▶ Notation:  $\forall$  means “for all”, and  $\exists$  means “there exists”.
- ▶ Let  $I$  be any set. Then a collection  $\{A_i\}_{i \in I}$  of sets  $A_i$  is called a family of sets parametrized by  $I$ .
- ▶ For such a family, define the union and the intersection by

$$\bigcup_{i \in I} A_i = \{x \mid \exists i \in I \text{ such that } x \in A_i\},$$

$$\bigcap_{i \in I} A_i = \{x \mid x \in A_i \forall i \in I\}.$$

# De Morgan's laws

- Proposition 1.2 (De Morgan's laws):

$$(1) \quad X \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} X \setminus A_i,$$

$$(2) \quad X \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} X \setminus A_i.$$

# De Morgan's laws

- ▶ Proposition 1.2 (De Morgan's laws):

$$(1) \quad X \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} X \setminus A_i,$$

$$(2) \quad X \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} X \setminus A_i.$$

- ▶ Proof of (1): We first prove  $X \setminus \bigcup_{i \in I} A_i \subset \bigcap_{i \in I} X \setminus A_i$ .

# De Morgan's laws

- ▶ Proposition 1.2 (De Morgan's laws):

$$(1) \quad X \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} X \setminus A_i,$$

$$(2) \quad X \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} X \setminus A_i.$$

- ▶ Proof of (1): We first prove  $X \setminus \bigcup_{i \in I} A_i \subset \bigcap_{i \in I} X \setminus A_i$ .

- ▶ Let  $a \in X \setminus \bigcup_{i \in I} A_i$ . Then  $a \in X$  and  $a \notin \bigcup_{i \in I} A_i$ . The latter means that  $a$  is not in any of the  $A_i$ . Since  $a \in X$ ,  $\forall i \in I$  we have  $a \in X \setminus A_i$ . This exactly means that  $a \in \bigcap_{i \in I} X \setminus A_i$ .

# De Morgan's laws

- ▶ Proposition 1.2 (De Morgan's laws):

$$(1) \quad X \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} X \setminus A_i,$$

$$(2) \quad X \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} X \setminus A_i.$$

- ▶ Proof of (1): We first prove  $X \setminus \bigcup_{i \in I} A_i \subset \bigcap_{i \in I} X \setminus A_i$ .
  - ▶ Let  $a \in X \setminus \bigcup_{i \in I} A_i$ . Then  $a \in X$  and  $a \notin \bigcup_{i \in I} A_i$ . The latter means that  $a$  is not in any of the  $A_i$ . Since  $a \in X$ ,  $\forall i \in I$  we have  $a \in X \setminus A_i$ . This exactly means that  $a \in \bigcap_{i \in I} X \setminus A_i$ .
- ▶ We now prove  $\bigcap_{i \in I} X \setminus A_i \subset X \setminus \bigcup_{i \in I} A_i$ .

# De Morgan's laws

- ▶ Proposition 1.2 (De Morgan's laws):

$$(1) \quad X \setminus \bigcup_{i \in I} A_i = \bigcap_{i \in I} X \setminus A_i,$$

$$(2) \quad X \setminus \bigcap_{i \in I} A_i = \bigcup_{i \in I} X \setminus A_i.$$

- ▶ Proof of (1): We first prove  $X \setminus \bigcup_{i \in I} A_i \subset \bigcap_{i \in I} X \setminus A_i$ .
  - ▶ Let  $a \in X \setminus \bigcup_{i \in I} A_i$ . Then  $a \in X$  and  $a \notin \bigcup_{i \in I} A_i$ . The latter means that  $a$  is not in any of the  $A_i$ . Since  $a \in X$ ,  $\forall i \in I$  we have  $a \in X \setminus A_i$ . This exactly means that  $a \in \bigcap_{i \in I} X \setminus A_i$ .
- ▶ We now prove  $\bigcap_{i \in I} X \setminus A_i \subset X \setminus \bigcup_{i \in I} A_i$ .
  - ▶ Let  $a \in \bigcap_{i \in I} X \setminus A_i$ . Then  $\forall i \in I$  we have  $a \in X \setminus A_i$ . This means that  $\forall i \in I$  we have  $a \in X$  and  $a \notin A_i$ . The latter implies  $a \notin \bigcup_{i \in I} A_i$ . Since  $a \in X$ , we obtain  $a \in X \setminus \bigcup_{i \in I} A_i$ .

# Cartesian products

- ▶ If  $A$  and  $B$  are two sets, then the *Cartesian product*  $A \times B$  is the set of all pairs  $(a, b)$ , where  $a \in A$  and  $b \in B$ . That is

$$A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}.$$

# Cartesian products

- ▶ If  $A$  and  $B$  are two sets, then the *Cartesian product*  $A \times B$  is the set of all pairs  $(a, b)$ , where  $a \in A$  and  $b \in B$ . That is

$$A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}.$$

- ▶ If  $A_1, \dots, A_n$ , the Cartesian product  $A_1 \times \dots \times A_n$  is the set of all  $n$ -tuples  $(a_1, \dots, a_n)$ , where  $a_i \in A_i$  for all  $i = 1, \dots, n$ .

$$A_1 \times \dots \times A_n = \{(a_1, \dots, a_n) \mid a_i \in A_i \ \forall i = 1, \dots, n\}.$$



# Cartesian products

- ▶ If  $A$  and  $B$  are two sets, then the *Cartesian product*  $A \times B$  is the set of all pairs  $(a, b)$ , where  $a \in A$  and  $b \in B$ . That is

$$A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}.$$

- ▶ If  $A_1, \dots, A_n$ , the Cartesian product  $A_1 \times \dots \times A_n$  is the set of all  $n$ -tuples  $(a_1, \dots, a_n)$ , where  $a_i \in A_i$  for all  $i = 1, \dots, n$ .

$$A_1 \times \dots \times A_n = \{(a_1, \dots, a_n) \mid a_i \in A_i \ \forall i = 1, \dots, n\}.$$

- ▶ Example 1.4:  $\mathbb{R}^n = \mathbb{R} \times \dots \times \mathbb{R}$  ( $n$  times).

# Cartesian products

- ▶ If  $A$  and  $B$  are two sets, then the *Cartesian product*  $A \times B$  is the set of all pairs  $(a, b)$ , where  $a \in A$  and  $b \in B$ . That is

$$A \times B = \{(a, b) \mid a \in A \text{ and } b \in B\}.$$

- ▶ If  $A_1, \dots, A_n$ , the Cartesian product  $A_1 \times \dots \times A_n$  is the set of all  $n$ -tuples  $(a_1, \dots, a_n)$ , where  $a_i \in A_i$  for all  $i = 1, \dots, n$ .

$$A_1 \times \dots \times A_n = \{(a_1, \dots, a_n) \mid a_i \in A_i \ \forall i = 1, \dots, n\}.$$

- ▶ Example 1.4:  $\mathbb{R}^n = \mathbb{R} \times \dots \times \mathbb{R}$  ( $n$  times).
- ▶ In general, if  $\{A_i\}_{i \in I}$  is a family of sets then the Cartesian product  $\prod_{i \in I} A_i$  is the set of all functions  $a : I \rightarrow \bigcup_{i \in I} A_i$  such that  $a(i) \in A_i$  for all  $i \in I$ .

$$\prod_{i \in I} A_i = \left\{ a : I \rightarrow \bigcup_{i \in I} A_i \mid a(i) \in A_i \ \forall i \in I \right\}.$$

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .

# Relations

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,
  - ▶ *symmetric* if  $xCy$  implies that  $yCx$  for all  $x, y \in A$ ,

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,
  - ▶ *symmetric* if  $xCy$  implies that  $yCx$  for all  $x, y \in A$ ,
  - ▶ *anti-symmetric* if  $xCy$  and  $yCx$  implies  $x = y$  for all  $x, y \in A$ ,



- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,
  - ▶ *symmetric* if  $xCy$  implies that  $yCx$  for all  $x, y \in A$ ,
  - ▶ *anti-symmetric* if  $xCy$  and  $yCx$  implies  $x = y$  for all  $x, y \in A$ ,
  - ▶ *transitive* if  $xCy$  and  $yCz$  implies that  $xCz$  for all  $x, y, z \in A$ ,

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,
  - ▶ *symmetric* if  $xCy$  implies that  $yCx$  for all  $x, y \in A$ ,
  - ▶ *anti-symmetric* if  $xCy$  and  $yCx$  implies  $x = y$  for all  $x, y \in A$ ,
  - ▶ *transitive* if  $xCy$  and  $yCz$  implies that  $xCz$  for all  $x, y, z \in A$ ,
  - ▶ *total* if either  $xCy$  or  $yCx$  when  $x, y \in A$ .

- ▶ A *binary relation*  $C$  on a set  $A$  is a subset  $C \subset A \times A$ . When  $(x, y) \in C$ , we will often write  $xCy$ .
- ▶ Example 1.5: The subset of  $\mathbb{R} \times \mathbb{R}$  given by  $C = \{(x, y) \mid x \leq y\}$  is a relation on  $\mathbb{R}$ , and  $xCy$  iff  $x \leq y$ .
- ▶ A relation  $C \subset A \times A$  is called
  - ▶ *reflexive* if  $xCx$  for all  $x \in A$ ,
  - ▶ *symmetric* if  $xCy$  implies that  $yCx$  for all  $x, y \in A$ ,
  - ▶ *anti-symmetric* if  $xCy$  and  $yCx$  implies  $x = y$  for all  $x, y \in A$ ,
  - ▶ *transitive* if  $xCy$  and  $yCz$  implies that  $xCz$  for all  $x, y, z \in A$ ,
  - ▶ *total* if either  $xCy$  or  $yCx$  when  $x, y \in A$ .
- ▶ Example 1.6: The relation in Example 1.5 is reflexive, anti-symmetric, transitive, and total, but it is not symmetric.

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .
  - ▶ If the partial order relation is also total, then it is called a *total order*, and  $(A, C)$  is called a *totally ordered set*.

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .
  - ▶ If the partial order relation is also total, then it is called a *total order*, and  $(A, C)$  is called a *totally ordered set*.
- ▶ Example 1.7: The pair  $(\mathbb{R}, \leq)$  is a totally ordered set.

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .
  - ▶ If the partial order relation is also total, then it is called a *total order*, and  $(A, C)$  is called a *totally ordered set*.
- ▶ Example 1.7: The pair  $(\mathbb{R}, \leq)$  is a totally ordered set.
- ▶ An *equivalence relation* is a relation which is reflexive, symmetric, and transitive.



# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .
  - ▶ If the partial order relation is also total, then it is called a *total order*, and  $(A, C)$  is called a *totally ordered set*.
- ▶ Example 1.7: The pair  $(\mathbb{R}, \leq)$  is a totally ordered set.
- ▶ An *equivalence relation* is a relation which is reflexive, symmetric, and transitive.
  - ▶ When  $C$  is an equivalence relation, we use the notation  $x \sim y$  for  $xCy$  and say that  $x$  is equivalent to  $y$ .

# Partial order and equivalence relation

- ▶ A relation  $C$  on a set  $A$  is called a *partial order* if it is reflexive, anti-symmetric, and transitive. The pair  $(A, C)$  is called a *poset*.
  - ▶ We often denote partial orders by the symbol  $\preceq$ .
  - ▶ If the partial order relation is also total, then it is called a *total order*, and  $(A, C)$  is called a *totally ordered set*.
- ▶ Example 1.7: The pair  $(\mathbb{R}, \leq)$  is a totally ordered set.
- ▶ An *equivalence relation* is a relation which is reflexive, symmetric, and transitive.
  - ▶ When  $C$  is an equivalence relation, we use the notation  $x \sim y$  for  $xCy$  and say that  $x$  is equivalent to  $y$ .
- ▶ Example 1.8: Fix  $p \in \mathbb{N}$ . Let  $C \subset \mathbb{Z} \times \mathbb{Z}$  be the subset of pairs  $(m, n)$  such that  $m - n$  is a multiple of  $p$ , i.e.  $m - n = kp$  for some  $k \in \mathbb{Z}$ . Then  $C$  is an equivalence relation on  $\mathbb{Z}$ .

# Equivalence classes

- ▶ Given any equivalence relation on a set  $A$ , it is possible to partition  $A$  into smaller sets consisting of elements that are equivalent to each other. More precisely, for  $x \in A$  let

$$[x] = \{y \mid y \sim x\}$$

be the *equivalence class* of  $x$ .

# Equivalence classes

- ▶ Given any equivalence relation on a set  $A$ , it is possible to partition  $A$  into smaller sets consisting of elements that are equivalent to each other. More precisely, for  $x \in A$  let

$$[x] = \{y \mid y \sim x\}$$

be the *equivalence class* of  $x$ .

- ▶ Note that  $x \sim x$  by reflexivity so  $x \in [x]$  for all  $x \in A$ .

# Equivalence classes

- ▶ Given any equivalence relation on a set  $A$ , it is possible to partition  $A$  into smaller sets consisting of elements that are equivalent to each other. More precisely, for  $x \in A$  let

$$[x] = \{y \mid y \sim x\}$$

be the *equivalence class* of  $x$ .

- ▶ Note that  $x \sim x$  by reflexivity so  $x \in [x]$  for all  $x \in A$ .
- ▶ Lemma 1.3: Let  $\sim$  denote an equivalence relation on a set  $A$ . Then for two elements  $x, x' \in A$ , the equivalence classes  $[x]$  and  $[x']$  are either disjoint or equal.

# Equivalence classes

- ▶ Given any equivalence relation on a set  $A$ , it is possible to partition  $A$  into smaller sets consisting of elements that are equivalent to each other. More precisely, for  $x \in A$  let

$$[x] = \{y \mid y \sim x\}$$

be the *equivalence class* of  $x$ .

- ▶ Note that  $x \sim x$  by reflexivity so  $x \in [x]$  for all  $x \in A$ .
- ▶ Lemma 1.3: Let  $\sim$  denote an equivalence relation on a set  $A$ . Then for two elements  $x, x' \in A$ , the equivalence classes  $[x]$  and  $[x']$  are either disjoint or equal.
- ▶ Proof: It is equivalent to show that if  $[x]$  and  $[x']$  are not disjoint then  $[x] = [x']$ .

# Equivalence classes (contd)

- ▶ Suppose  $[x]$  and  $[x']$  are not disjoint. Then there is a  $z \in A$  such that  $z \in [x]$  and  $z \in [x']$ . That is,  $z \sim x$  and  $z \sim x'$ .

# Equivalence classes (contd)

- ▶ Suppose  $[x]$  and  $[x']$  are not disjoint. Then there is a  $z \in A$  such that  $z \in [x]$  and  $z \in [x']$ . That is,  $z \sim x$  and  $z \sim x'$ .
- ▶ Showing  $[x] \subset [x']$ : Let  $y \in [x]$ . Then  $y \sim x$ , so  $x \sim y$  (by symmetry). Since  $z \sim x$ , by transitivity,  $z \sim y$ . Thus  $y \sim z$  (by symmetry). Since  $z \sim x'$ , by transitivity,  $y \sim x'$ . This means that  $y \in [x']$ . Hence  $[x] \subset [x']$ .



# Equivalence classes (contd)

- ▶ Suppose  $[x]$  and  $[x']$  are not disjoint. Then there is a  $z \in A$  such that  $z \in [x]$  and  $z \in [x']$ . That is,  $z \sim x$  and  $z \sim x'$ .
- ▶ Showing  $[x] \subset [x']$ : Let  $y \in [x]$ . Then  $y \sim x$ , so  $x \sim y$  (by symmetry). Since  $z \sim x$ , by transitivity,  $z \sim y$ . Thus  $y \sim z$  (by symmetry). Since  $z \sim x'$ , by transitivity,  $y \sim x'$ . This means that  $y \in [x']$ . Hence  $[x] \subset [x']$ .
- ▶ Showing  $[x'] \subset [x]$ : similar.

# Equivalence classes (contd)

- ▶ The set of equivalence classes on a set  $A$  with respect to an equivalence relation  $\sim$  is denoted  $A/\sim$ . That is,

$$A/\sim = \{[x] \mid x \in A\}.$$

# Equivalence classes (contd)

- ▶ The set of equivalence classes on a set  $A$  with respect to an equivalence relation  $\sim$  is denoted  $A/\sim$ . That is,

$$A/\sim = \{[x] \mid x \in A\}.$$

- ▶ Example 1.9: Consider the relation  $\sim$  from Example 1.8. The equivalence class of an integer  $n \in \mathbb{Z}$  is the set of integers

$$[n] = \{\dots, n - 2p, n - p, n, n + p, n + 2p, \dots\}.$$

and we can write  $\mathbb{Z}$  as the union of  $p$  equivalence classes:

$$\mathbb{Z} = [0] \cup [1] \cup [2] \cup \dots \cup [p-1].$$

Similarly,

$$\mathbb{Z}/\sim = \{[0], [1], \dots, [p-1]\}.$$