

## HOMEWORK 4 SOLUTIONS – MATH 4341

**Problem 1.** (a) Suppose  $\mathcal{T}_1$  and  $\mathcal{T}_2$  are two different topologies on a set  $X$ . When is the identity map  $id : X \rightarrow X$  given by  $id(x) = x$  a continuous map from  $(X, \mathcal{T}_1)$  to  $(X, \mathcal{T}_2)$ ?

(b) Show that the subspace topology  $\mathcal{T}_Y$  is the smallest topology on  $Y \subset X$  for which the inclusion  $\iota : Y \rightarrow X$  is a continuous map.

*Proof.* (a)  $id : (X, \mathcal{T}_1) \rightarrow (X, \mathcal{T}_2)$  is continuous  $\Leftrightarrow id^{-1}(U) \in \mathcal{T}_1$  for all  $U \in \mathcal{T}_2 \Leftrightarrow U \in \mathcal{T}_1$  for all  $U \in \mathcal{T}_2 \Leftrightarrow \mathcal{T}_2 \subset \mathcal{T}_1$ .

(b) We first prove that  $\iota : (Y, \mathcal{T}_Y) \rightarrow X$  is continuous. Let  $V \subset X$  be an open set. Then  $\iota^{-1}(V) = Y \cap V \in \mathcal{T}_Y$ . Hence  $\iota : (Y, \mathcal{T}_Y) \rightarrow X$  is continuous.

Suppose  $\iota : (Y, \mathcal{T}) \rightarrow X$  is continuous. We will show that  $\mathcal{T}_Y \subset \mathcal{T}$ . Let  $U \in \mathcal{T}_Y$ . Then  $U = Y \cap V$  for some open set  $V \subset X$ . Since  $\iota^{-1}(V) = Y \cap V = U$  and  $\iota : (Y, \mathcal{T}) \rightarrow X$  is continuous, we have  $U \in \mathcal{T}$ . Hence  $\mathcal{T}_Y \subset \mathcal{T}$ .  $\square$

**Problem 2.** (a) Let  $Y \subset X$  be an open subset of a topological space  $X$ . Show that a set  $U \subset Y$  is open in the subspace topology on  $Y$  if and only if  $U$  is open in  $X$ .

(b) Let  $Y \subset X$  be a closed subset of a topological space  $X$ . Show that a set  $U \subset Y$  is closed in the subspace topology on  $Y$  if and only if  $U$  is closed in  $X$ .

*Proof.* (a) Suppose  $U \subset Y$  is open in  $X$ . Since  $U = Y \cap U$ , we have  $U$  is open in  $Y$ .

Suppose  $U \subset Y$  is open. Then  $U = Y \cap V$  for some open set  $V \subset X$ . Since both  $Y$  and  $V$  are open in  $X$ ,  $U = Y \cap V$  is open in  $X$ .

(b) Suppose  $U \subset Y$  is closed in  $X$ . Then  $X \setminus U$  is open in  $X$ . Since  $Y \setminus U = Y \cap (X \setminus U)$ , we have  $Y \setminus U$  is open in  $Y$ . This means that  $U$  is closed in  $Y$ .

Suppose  $U \subset Y$  is closed. Then  $Y \setminus U$  is open in  $Y$ , and hence  $Y \setminus U = Y \cap V$  for some open set  $V \subset X$ . We have

$$U = Y \setminus (Y \setminus U) = Y \setminus (Y \cap V) = (Y \setminus Y) \cup (Y \setminus V) = Y \setminus V = Y \cap (X \setminus V).$$

Since both  $Y$  and  $X \setminus V$  are closed in  $X$ ,  $U = Y \cap (X \setminus V)$  is closed in  $X$ .  $\square$

**Problem 3.** Let  $(X_1, d_1)$  and  $(X_2, d_2)$  be metric spaces. Define a function on  $X_1 \times X_2$  by

$$d((x_1, x_2), (y_1, y_2)) = \max(d_1(x_1, y_1), d_2(x_2, y_2)).$$

(a) Show that  $d$  is a metric on  $X_1 \times X_2$ .

(b) Show that the metric topology on  $X_1 \times X_2$  induced by  $d$  is the product topology, where  $X_1$  and  $X_2$  have the metric topologies from  $d_1$  and  $d_2$  respectively.

*Proof.* (a) (positivity)  $d((x_1, x_2), (y_1, y_2)) = \max(d_1(x_1, y_1), d_2(x_2, y_2)) \geq 0$  and equality holds iff  $d_1(x_1, y_1) = 0$  and  $d_2(x_2, y_2) = 0$ , i.e.  $x_1 = y_1$  and  $x_2 = y_2$ , so  $(x_1, x_2) = (y_1, y_2)$ .

(symmetry)  $d((x_1, x_2), (y_1, y_2)) = d((y_1, y_2), (x_1, x_2))$ , since  $d_1(x_1, y_1) = d_1(y_1, x_1)$  and  $d_2(x_2, y_2) = d_2(y_2, x_2)$ .

(triangle inequality) We have  $d((x_1, x_2), (y_1, y_2)) + d((y_1, y_2), (z_1, z_2)) \geq d_1(x_1, y_1) + d_1(y_1, z_1) \geq d_1(x_1, z_1)$ . Similarly,  $d((x_1, x_2), (y_1, y_2)) + d((y_1, y_2), (z_1, z_2)) \geq d_2(x_2, y_2) + d_2(y_2, z_2) \geq d_2(x_2, z_2)$ . Hence

$$d((x_1, x_2), (y_1, y_2)) + d((y_1, y_2), (z_1, z_2)) \geq \max(d_1(x_1, z_1), d_2(x_2, z_2)) = d((x_1, x_2), (z_1, z_2)).$$

(b) We will apply Lemma 2.4 in the lecture notes. Let  $B_d((y_1, y_2), r)$  be an open ball in the topology  $\mathcal{T}_d$  and  $(x_1, x_2) \in B_d((y_1, y_2), r)$ . Then

$$r > d((x_1, x_2), (y_1, y_2)) = \max(d_1(x_1, y_1), d_2(x_2, y_2))$$

and  $(x_1, x_2) \in B_{d_1}(y_1, r) \times B_{d_2}(y_2, r) \subset B_d((y_1, y_2), r)$ . Indeed, if  $(z_1, z_2) \in B_{d_1}(y_1, r) \times B_{d_2}(y_2, r)$ , then  $d_1(y_1, z_1) < r$  and  $d_2(y_2, z_2) < r$ . This implies that  $d((y_1, y_2), (z_1, z_2)) = \max(d_1(y_1, z_1), d_2(y_2, z_2)) < r$ . Hence  $(z_1, z_2) \in B_d((y_1, y_2), r)$ .

Conversely, let  $U_1 \times U_2$  be a basis element for the product topology on  $X_1 \times X_2$ , and  $(x_1, x_2) \in U_1 \times U_2$ . Since  $U_1 \subset X_1$  is open and  $x_1 \in U_1$ , there exists an open ball  $B_{d_1}(y_1, r_1)$  such that  $x_1 \in B_{d_1}(y_1, r_1) \subset U_1$ . Similarly, there exists an open ball  $B_{d_2}(y_2, r_2)$  such that  $x_2 \in B_{d_2}(y_2, r_2) \subset U_2$ . Let

$$r = \min(r_1 - d(x_1, y_1), r_2 - d(x_2, y_2)) > 0.$$

Then  $(x_1, x_2) \in B_d((x_1, x_2), r) \subset B_{d_1}(y_1, r_1) \times B_{d_2}(y_2, r_2) \subset U_1 \times U_2$ . Indeed, if  $(z_1, z_2) \in B_d((x_1, x_2), r)$  then  $d((x_1, x_2), (z_1, z_2)) < r$ , i.e.

$$\max(d_1(x_1, z_1), d_2(x_2, z_2)) < \min(r_1 - d(x_1, y_1), r_2 - d(x_2, y_2)).$$

This implies that  $d_1(x_1, z_1) < r_1 - d(x_1, y_1)$  and  $d_1(x_2, z_2) < r_2 - d(x_2, y_2)$ , i.e.  $d_1(x_1, z_1) + d(x_1, y_1) < r_1$  and  $d_1(x_2, z_2) + d(x_2, y_2) < r_2$ . Together with the triangle inequality, we obtain  $d(y_1, z_1) < r_1$  and  $d(y_2, z_2) < r_2$ . Hence  $z_1 \in B_{d_1}(y_1, r_1)$  and  $z_2 \in B_{d_2}(y_2, r_2)$ , so  $(z_1, z_2) \in B_{d_1}(y_1, r_1) \times B_{d_2}(y_2, r_2)$ .  $\square$