

4. Closeness

Math 4341 (Topology)

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .
 - ▶ The *closure* of Y , denoted \overline{Y} , is the intersection of all closed subsets of X containing Y .

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .
 - ▶ The *closure* of Y , denoted \overline{Y} , is the intersection of all closed subsets of X containing Y .
 - ▶ The subset Y is called *dense* in X if $\overline{Y} = X$.

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .
 - ▶ The *closure* of Y , denoted \overline{Y} , is the intersection of all closed subsets of X containing Y .
 - ▶ The subset Y is called *dense* in X if $\overline{Y} = X$.
- ▶ $\text{Int } Y$ is open, since it is a union of open sets. Likewise, \overline{Y} is closed, and we have

$$\text{Int } Y \subset Y \subset \overline{Y}.$$

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .
 - ▶ The *closure* of Y , denoted \overline{Y} , is the intersection of all closed subsets of X containing Y .
 - ▶ The subset Y is called *dense* in X if $\overline{Y} = X$.
- ▶ $\text{Int } Y$ is open, since it is a union of open sets. Likewise, \overline{Y} is closed, and we have

$$\text{Int } Y \subset Y \subset \overline{Y}.$$

- ▶ Y is open iff $Y = \text{Int } Y$ and Y is closed iff $Y = \overline{Y}$. Moreover, $\text{Int } Y$ is the largest open subset contained in Y , and \overline{Y} is the smallest closed subset containing Y .

Interior and closure

- ▶ Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$.
 - ▶ The *interior* of Y , denoted $\overset{\circ}{Y}$ or $\text{Int } Y$, is the union of all open subsets of X contained in Y .
 - ▶ The *closure* of Y , denoted \overline{Y} , is the intersection of all closed subsets of X containing Y .
 - ▶ The subset Y is called *dense* in X if $\overline{Y} = X$.
- ▶ $\text{Int } Y$ is open, since it is a union of open sets. Likewise, \overline{Y} is closed, and we have

$$\text{Int } Y \subset Y \subset \overline{Y}.$$

- ▶ Y is open iff $Y = \text{Int } Y$ and Y is closed iff $Y = \overline{Y}$. Moreover, $\text{Int } Y$ is the largest open subset contained in Y , and \overline{Y} is the smallest closed subset containing Y .
- ▶ Note that $\text{Int}(X \setminus Y) = X \setminus \overline{Y}$ and $\overline{X \setminus Y} = X \setminus \text{Int}(Y)$.

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
 - ▶ (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
 - ▶ (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - ▶ (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,

- **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
- (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,
 - (iii) $\text{Int} Y \cup \text{Int} Z \subset \text{Int}(Y \cup Z)$, and

- **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
- (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,
 - (iii) $\text{Int} Y \cup \text{Int} Z \subset \text{Int}(Y \cup Z)$, and
 - (iv) $\text{Int} Y \cap \text{Int} Z = \text{Int}(Y \cap Z)$.

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
 - ▶ (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - ▶ (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,
 - ▶ (iii) $\text{Int} Y \cup \text{Int} Z \subset \text{Int}(Y \cup Z)$, and
 - ▶ (iv) $\text{Int} Y \cap \text{Int} Z = \text{Int}(Y \cap Z)$.
- ▶ *Proof.* We will show (i). The remaining ones are similar.

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
 - ▶ (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - ▶ (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,
 - ▶ (iii) $\text{Int} Y \cup \text{Int} Z \subset \text{Int}(Y \cup Z)$, and
 - ▶ (iv) $\text{Int} Y \cap \text{Int} Z = \text{Int}(Y \cap Z)$.
- ▶ *Proof.* We will show (i). The remaining ones are similar.
 - ▶ Since $\overline{Y} \cup \overline{Z}$ is a closed subset containing $Y \cup Z$, we have $\overline{Y \cup Z} \subset \overline{Y} \cup \overline{Z}$.

- ▶ **Proposition 4.1.** Let Y and Z be subsets of a topological space X . Then
 - ▶ (i) $\overline{Y \cup Z} = \overline{Y} \cup \overline{Z}$,
 - ▶ (ii) $\overline{Y \cap Z} \subset \overline{Y} \cap \overline{Z}$,
 - ▶ (iii) $\text{Int} Y \cup \text{Int} Z \subset \text{Int}(Y \cup Z)$, and
 - ▶ (iv) $\text{Int} Y \cap \text{Int} Z = \text{Int}(Y \cap Z)$.
- ▶ *Proof.* We will show (i). The remaining ones are similar.
 - ▶ Since $\overline{Y \cup Z}$ is a closed subset containing $Y \cup Z$, we have $\overline{Y \cup Z} \subset \overline{Y \cup Z}$.
 - ▶ Since $Y \subset \overline{Y \cup Z}$ and the latter set is closed, we have $\overline{Y} \subset \overline{Y \cup Z}$. For the same reason $\overline{Z} \subset \overline{Y \cup Z}$. Hence $\overline{Y} \cup \overline{Z} \subset \overline{Y \cup Z}$.

Boundary and limit points

- ▶ In a topological space X , we say that U is a *neighborhood* (nbhd) of a point x if U is open and $x \in U$.

Boundary and limit points

- ▶ In a topological space X , we say that U is a *neighborhood* (nbhd) of a point x if U is open and $x \in U$.
- ▶ **Definition.** Let $Y \subset X$. Then

Boundary and limit points

- ▶ In a topological space X , we say that U is a *neighborhood* (nbhd) of a point x if U is open and $x \in U$.
- ▶ **Definition.** Let $Y \subset X$. Then
 - ▶ The *boundary* of Y , denoted ∂Y , is the set

$$\partial Y = \{x \in X \mid U \cap Y \neq \emptyset \text{ and } U \cap Y^c \neq \emptyset \text{ for all nbhds } U \text{ of } x\}.$$

That is, $x \in \partial Y$ iff all nbhds of x intersect both Y and Y^c .

Boundary and limit points

- ▶ In a topological space X , we say that U is a *neighborhood* (nbhd) of a point x if U is open and $x \in U$.

- ▶ **Definition.** Let $Y \subset X$. Then

- ▶ The *boundary* of Y , denoted ∂Y , is the set

$$\partial Y = \{x \in X \mid U \cap Y \neq \emptyset \text{ and } U \cap Y^c \neq \emptyset \text{ for all nbhds } U \text{ of } x\}.$$

That is, $x \in \partial Y$ iff all nbhds of x intersect both Y and Y^c .

- ▶ A *limit point* of Y is a point $x \in X$ with the property that all its nbhds intersect Y in a point which is not x itself. Let Y' denote the set of all limit points of Y .

Boundary and limit points

- ▶ In a topological space X , we say that U is a *neighborhood* (nbhd) of a point x if U is open and $x \in U$.

- ▶ **Definition.** Let $Y \subset X$. Then

- ▶ The *boundary* of Y , denoted ∂Y , is the set

$$\partial Y = \{x \in X \mid U \cap Y \neq \emptyset \text{ and } U \cap Y^c \neq \emptyset \text{ for all nbhds } U \text{ of } x\}.$$

That is, $x \in \partial Y$ iff all nbhds of x intersect both Y and Y^c .

- ▶ A *limit point* of Y is a point $x \in X$ with the property that all its nbhds intersect Y in a point which is not x itself. Let Y' denote the set of all limit points of Y .
- ▶ **Example.** Let $Y = [0, 1) \cup \{2\} \subset \mathbb{R}$. Then $\text{Int } Y = (0, 1)$, $\bar{Y} = [0, 1] \cup \{2\}$, $\partial Y = \{0, 1, 2\}$, and $Y' = [0, 1]$.

Boundary and limit points

- ▶ **Theorem 4.2.** Let $Y \subset X$. Then
 - ▶ (i) $\partial Y = X \setminus (\text{Int } Y \cup \text{Int}(X \setminus Y)) = \overline{Y} \cap \overline{X \setminus Y}$,
 - ▶ (ii) $\overline{Y} = Y \cup \partial Y$, and
 - ▶ (iii) $\overline{Y} = Y \cup Y'$.

Boundary and limit points

- ▶ **Theorem 4.2.** Let $Y \subset X$. Then
 - ▶ (i) $\partial Y = X \setminus (\text{Int } Y \cup \text{Int}(X \setminus Y)) = \overline{Y} \cap \overline{X \setminus Y}$,
 - ▶ (ii) $\overline{Y} = Y \cup \partial Y$, and
 - ▶ (iii) $\overline{Y} = Y \cup Y'$.
- ▶ *Proof.* (i) is equivalent to $X \setminus \partial Y = \text{Int } Y \cup \text{Int}(X \setminus Y)$.
 - ▶ Let $x \in X \setminus \partial Y$. Then there is a nbhd U of x so that $U \subset Y$ or $U \subset X \setminus Y$. Hence $x \in \text{Int } Y$ or $x \in \text{Int}(X \setminus Y)$.
 - ▶ Suppose $x \in \text{Int } Y$. Then there is an open set U such that $x \in U \subset Y$. Since $U \cap Y^c = \emptyset$, we have $x \notin \partial Y$. Similarly, if $x \in \text{Int}(X \setminus Y)$, there is an open nbhd U of x with $U \cap Y = \emptyset$, so once again $x \notin \partial Y$. This shows (i).

Boundary and limit points

- ▶ **Theorem 4.2.** Let $Y \subset X$. Then
 - ▶ (i) $\partial Y = X \setminus (\text{Int } Y \cup \text{Int}(X \setminus Y)) = \overline{Y} \cap \overline{X \setminus Y}$,
 - ▶ (ii) $\overline{Y} = Y \cup \partial Y$, and
 - ▶ (iii) $\overline{Y} = Y \cup Y'$.
- ▶ *Proof.* (i) is equivalent to $X \setminus \partial Y = \text{Int } Y \cup \text{Int}(X \setminus Y)$.
 - ▶ Let $x \in X \setminus \partial Y$. Then there is a nbhd U of x so that $U \subset Y$ or $U \subset X \setminus Y$. Hence $x \in \text{Int } Y$ or $x \in \text{Int}(X \setminus Y)$.
 - ▶ Suppose $x \in \text{Int } Y$. Then there is an open set U such that $x \in U \subset Y$. Since $U \cap Y^c = \emptyset$, we have $x \notin \partial Y$. Similarly, if $x \in \text{Int}(X \setminus Y)$, there is an open nbhd U of x with $U \cap Y = \emptyset$, so once again $x \notin \partial Y$. This shows (i).
- ▶ (ii) follows from (i):

$$Y \cup \partial Y = Y \cup (\overline{Y} \cap \overline{X \setminus Y}) = \overline{Y} \cap X = \overline{Y}.$$

Boundary and limit points

- ▶ Finally, we use (ii) and show that $Y \cup \partial Y = Y \cup Y'$. To see this, it suffices to show that $\partial Y \setminus Y = Y' \setminus Y$.

Boundary and limit points

- ▶ Finally, we use (ii) and show that $Y \cup \partial Y = Y \cup Y'$. To see this, it suffices to show that $\partial Y \setminus Y = Y' \setminus Y$.
 - ▶ Let $x \in \partial Y \setminus Y$. Then any nbhd U of x intersects Y , and since $x \notin Y$, this necessarily means that U intersects Y in a point which is not x itself. Hence $x \in Y' \setminus Y$.

Boundary and limit points

- ▶ Finally, we use (ii) and show that $Y \cup \partial Y = Y \cup Y'$. To see this, it suffices to show that $\partial Y \setminus Y = Y' \setminus Y$.
 - ▶ Let $x \in \partial Y \setminus Y$. Then any nbhd U of x intersects Y , and since $x \notin Y$, this necessarily means that U intersects Y in a point which is not x itself. Hence $x \in Y' \setminus Y$.
 - ▶ Let $x \in Y' \setminus Y$. Then any nbhd U of x will intersect Y ; it will also intersect $X \setminus Y$, since x belongs to that set. Hence $x \in \partial Y \setminus Y$. This completes the proof.

Boundary and limit points

- ▶ Finally, we use (ii) and show that $Y \cup \partial Y = Y \cup Y'$. To see this, it suffices to show that $\partial Y \setminus Y = Y' \setminus Y$.
 - ▶ Let $x \in \partial Y \setminus Y$. Then any nbhd U of x intersects Y , and since $x \notin Y$, this necessarily means that U intersects Y in a point which is not x itself. Hence $x \in Y' \setminus Y$.
 - ▶ Let $x \in Y' \setminus Y$. Then any nbhd U of x will intersect Y ; it will also intersect $X \setminus Y$, since x belongs to that set. Hence $x \in \partial Y \setminus Y$. This completes the proof.
- ▶ **Remark.** The above theorem provides us with the following useful characterization of the closure: we see that $x \in \overline{Y}$ if and only if every nbhd of x intersects Y .

- **Example.** We claim that \mathbb{Q} is dense in \mathbb{R} , i.e. $\overline{\mathbb{Q}} = \mathbb{R}$. To see this, it suffices to show that $\partial\mathbb{Q} = \mathbb{R}$.

Boundary and limit points

- ▶ **Example.** We claim that \mathbb{Q} is dense in \mathbb{R} , i.e. $\overline{\mathbb{Q}} = \mathbb{R}$. To see this, it suffices to show that $\partial\mathbb{Q} = \mathbb{R}$.
 - ▶ Let $x \in \mathbb{R}$ and U a nbhd of x . Since U contains some open interval and any interval contains both rational and irrational numbers, we have $U \cap \mathbb{Q} \neq \emptyset$ and $U \cap (\mathbb{R} \setminus \mathbb{Q}) \neq \emptyset$. This is exactly the condition that $x \in \partial\mathbb{Q}$.

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called
 - ▶ T_0 if for every pair $x \neq y$ in X , there exists a nbhd of x that does not contain y or there exists a nbhd of y that does not contain x .

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called
 - ▶ T_0 if for every pair $x \neq y$ in X , there exists a nbhd of x that does not contain y *or* there exists a nbhd of y that does not contain x .
 - ▶ T_1 if for every pair $x \neq y$ in X , x has a nbhd not containing y *and* y has a nbhd not containing x .

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called
 - ▶ T_0 if for every pair $x \neq y$ in X , there exists a nbhd of x that does not contain y or there exists a nbhd of y that does not contain x .
 - ▶ T_1 if for every pair $x \neq y$ in X , x has a nbhd not containing y and y has a nbhd not containing x .
 - ▶ T_2 (or *Hausdorff*) if for every pair $x \neq y$ in X , there exists nbhds U_x and U_y of x and y respectively s.t. $U_x \cap U_y = \emptyset$.

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called
 - ▶ T_0 if for every pair $x \neq y$ in X , there exists a nbhd of x that does not contain y or there exists a nbhd of y that does not contain x .
 - ▶ T_1 if for every pair $x \neq y$ in X , x has a nbhd not containing y and y has a nbhd not containing x .
 - ▶ T_2 (or *Hausdorff*) if for every pair $x \neq y$ in X , there exists nbhds U_x and U_y of x and y respectively s.t. $U_x \cap U_y = \emptyset$.
- ▶ **Proposition 4.4.** X is T_1 iff $\{x\}$ is closed for all $x \in X$.

Separation axioms and Hausdorff spaces

- ▶ **Definition.** A topological space (X, \mathcal{T}) is called
 - ▶ T_0 if for every pair $x \neq y$ in X , there exists a nbhd of x that does not contain y or there exists a nbhd of y that does not contain x .
 - ▶ T_1 if for every pair $x \neq y$ in X , x has a nbhd not containing y and y has a nbhd not containing x .
 - ▶ T_2 (or *Hausdorff*) if for every pair $x \neq y$ in X , there exists nbhds U_x and U_y of x and y respectively s.t. $U_x \cap U_y = \emptyset$.
- ▶ **Proposition 4.4.** X is T_1 iff $\{x\}$ is closed for all $x \in X$.
- ▶ *Proof.* (\Leftarrow) Let $x \neq y$ in X . Then $X \setminus \{x\}$ is a nbhd of y not containing x , and $X \setminus \{y\}$ is a nbhd of x not containing y .
(\Rightarrow) Every $y \in X$ has a nbhd U_y not containing x . Since $X \setminus \{x\} = \bigcup_{y \neq x} U_y$ is open, $\{x\}$ is closed.

Separation axioms and Hausdorff spaces

- ▶ **Example.** Suppose X contains at least two points. With the trivial topology X is not T_0 (or T_1 or T_2), since the only nbhd of a point x is X .

Separation axioms and Hausdorff spaces

- ▶ **Example.** Suppose X contains at least two points. With the trivial topology X is not T_0 (or T_1 or T_2), since the only nbhd of a point x is X .
- ▶ **Example.** If X has the discrete topology, then X is Hausdorff (and T_1 and T_0).

Separation axioms and Hausdorff spaces

- ▶ **Example.** Suppose X contains at least two points. With the trivial topology X is not T_0 (or T_1 or T_2), since the only nbhd of a point x is X .
- ▶ **Example.** If X has the discrete topology, then X is Hausdorff (and T_1 and T_0).
- ▶ **Example.** Let $X = \{a, b, c\}$ with the topology

$$\mathcal{T} = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}.$$

Then X is T_0 but not T_1 .

Separation axioms and Hausdorff spaces

- ▶ **Example.** Suppose X contains at least two points. With the trivial topology X is not T_0 (or T_1 or T_2), since the only nbhd of a point x is X .
- ▶ **Example.** If X has the discrete topology, then X is Hausdorff (and T_1 and T_0).
- ▶ **Example.** Let $X = \{a, b, c\}$ with the topology

$$\mathcal{T} = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}.$$

Then X is T_0 but not T_1 .

Separation axioms and Hausdorff spaces

- ▶ **Example.** Suppose X contains at least two points. With the trivial topology X is not T_0 (or T_1 or T_2), since the only nbhd of a point x is X .
- ▶ **Example.** If X has the discrete topology, then X is Hausdorff (and T_1 and T_0).
- ▶ **Example.** Let $X = \{a, b, c\}$ with the topology

$$\mathcal{T} = \{\emptyset, \{b\}, \{c\}, \{b, c\}, X\}.$$

Then X is T_0 but not T_1 .

- ▶ **Example.** All metric spaces (with the metric topology) are Hausdorff.

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .
 - ▶ We say that a sequence $\{x_n\}_{n \in \mathbb{N}}$ *converges* to a point $x \in X$, if for any nbhd U of x , there is an $N \in \mathbb{N}$ s.t. $x_n \in U$ for all $n > N$. In which case, we write $x_n \rightarrow x$.

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .
 - ▶ We say that a sequence $\{x_n\}_{n \in \mathbb{N}}$ *converges* to a point $x \in X$, if for any nbhd U of x , there is an $N \in \mathbb{N}$ s.t. $x_n \in U$ for all $n > N$. In which case, we write $x_n \rightarrow x$.
 - ▶ A *subsequence* $\{y_n\}_{n \in \mathbb{N}}$ is a sequence such that $y_i = x_{n_i}$ for some $n_1 < n_2 < \dots$.

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .
 - ▶ We say that a sequence $\{x_n\}_{n \in \mathbb{N}}$ *converges* to a point $x \in X$, if for any nbhd U of x , there is an $N \in \mathbb{N}$ s.t. $x_n \in U$ for all $n > N$. In which case, we write $x_n \rightarrow x$.
 - ▶ A *subsequence* $\{y_n\}_{n \in \mathbb{N}}$ is a sequence such that $y_i = x_{n_i}$ for some $n_1 < n_2 < \dots$.
- ▶ **Proposition 4.5.** If $\{y_n\}$ is a subsequence of $\{x_n\}$ and $x_n \rightarrow x$, then $y_n \rightarrow x$.

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .
 - ▶ We say that a sequence $\{x_n\}_{n \in \mathbb{N}}$ *converges* to a point $x \in X$, if for any nbhd U of x , there is an $N \in \mathbb{N}$ s.t. $x_n \in U$ for all $n > N$. In which case, we write $x_n \rightarrow x$.
 - ▶ A *subsequence* $\{y_n\}_{n \in \mathbb{N}}$ is a sequence such that $y_i = x_{n_i}$ for some $n_1 < n_2 < \dots$.
- ▶ **Proposition 4.5.** If $\{y_n\}$ is a subsequence of $\{x_n\}$ and $x_n \rightarrow x$, then $y_n \rightarrow x$.
- ▶ **Example.** In the trivial topology, all sequences converge to any given point. In the discrete topology, for a sequence $\{x_n\}$ to converge to a point x , it has to be constantly equal to x for all large enough n .

Sequences and Convergence

- ▶ **Definition.** Let X be a topological space. A *sequence* in X is a family $\{x_n\}_{n \in \mathbb{N}}$ of points in X .
 - ▶ We say that a sequence $\{x_n\}_{n \in \mathbb{N}}$ *converges* to a point $x \in X$, if for any nbhd U of x , there is an $N \in \mathbb{N}$ s.t. $x_n \in U$ for all $n > N$. In which case, we write $x_n \rightarrow x$.
 - ▶ A *subsequence* $\{y_n\}_{n \in \mathbb{N}}$ is a sequence such that $y_i = x_{n_i}$ for some $n_1 < n_2 < \dots$.
- ▶ **Proposition 4.5.** If $\{y_n\}$ is a subsequence of $\{x_n\}$ and $x_n \rightarrow x$, then $y_n \rightarrow x$.
- ▶ **Example.** In the trivial topology, all sequences converge to any given point. In the discrete topology, for a sequence $\{x_n\}$ to converge to a point x , it has to be constantly equal to x for all large enough n .
- ▶ **Example.** The constant sequence is convergent, regardless of the topology on the space.

Sequences and Convergence

- **Proposition 4.6.** Let (X, d) be a metric space with the metric topology. Then a sequence $\{x_n\}$ in X converges to $x \in X$ if and only if

$$\forall \epsilon > 0, \exists N > 0 : n > N \Rightarrow d(x_n, x) < \epsilon.$$

Sequences and Convergence

- **Proposition 4.6.** Let (X, d) be a metric space with the metric topology. Then a sequence $\{x_n\}$ in X converges to $x \in X$ if and only if

$$\forall \epsilon > 0, \exists N > 0 : n > N \Rightarrow d(x_n, x) < \epsilon.$$

- *Proof.* There are two things to prove.

Sequences and Convergence

- ▶ **Proposition 4.6.** Let (X, d) be a metric space with the metric topology. Then a sequence $\{x_n\}$ in X converges to $x \in X$ if and only if

$$\forall \epsilon > 0, \exists N > 0 : n > N \Rightarrow d(x_n, x) < \epsilon.$$

- ▶ *Proof.* There are two things to prove.
 - ▶ Suppose $x_n \rightarrow x$ and let $\epsilon > 0$. Let $U = B_d(x, \epsilon)$. Then there exists an $N > 0$ s.t. $x_n \in U$ for all $n > N$, but this says that $d(x_n, x) < \epsilon$ for all $n > N$.

Sequences and Convergence

- ▶ **Proposition 4.6.** Let (X, d) be a metric space with the metric topology. Then a sequence $\{x_n\}$ in X converges to $x \in X$ if and only if

$$\forall \epsilon > 0, \exists N > 0 : n > N \Rightarrow d(x_n, x) < \epsilon.$$

- ▶ *Proof.* There are two things to prove.
 - ▶ Suppose $x_n \rightarrow x$ and let $\epsilon > 0$. Let $U = B_d(x, \epsilon)$. Then there exists an $N > 0$ s.t. $x_n \in U$ for all $n > N$, but this says that $d(x_n, x) < \epsilon$ for all $n > N$.
 - ▶ Let U be a nbhd of x . Then there exists an $\epsilon > 0$ s.t. $B_d(x, \epsilon) \subset U$. Now by assumption there is an $N > 0$ s.t. $x_n \in B_d(x, \epsilon) \subset U$ for all $n > N$.

Sequences and Convergence

- **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.

Sequences and Convergence

- ▶ **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.
- ▶ *Proof.* Assume $x \neq y$. Choose U and V disjoint nbhds of x and y respectively. By definition of convergence, we get $N_U, N_V > 0$ such that $x_n \in U$ for all $n > N_U$ and $x_n \in V$ for all $n > N_V$. For $n > \max(N_U, N_V)$ we therefore have $x_n \in U \cap V = \emptyset$, which is a contradiction.

Sequences and Convergence

- ▶ **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.
- ▶ *Proof.* Assume $x \neq y$. Choose U and V disjoint nbhds of x and y respectively. By definition of convergence, we get $N_U, N_V > 0$ such that $x_n \in U$ for all $n > N_U$ and $x_n \in V$ for all $n > N_V$. For $n > \max(N_U, N_V)$ we therefore have $x_n \in U \cap V = \emptyset$, which is a contradiction.
- ▶ **Definition.** Let X be a topological space.

Sequences and Convergence

- ▶ **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.
- ▶ *Proof.* Assume $x \neq y$. Choose U and V disjoint nbhds of x and y respectively. By definition of convergence, we get $N_U, N_V > 0$ such that $x_n \in U$ for all $n > N_U$ and $x_n \in V$ for all $n > N_V$. For $n > \max(N_U, N_V)$ we therefore have $x_n \in U \cap V = \emptyset$, which is a contradiction.
- ▶ **Definition.** Let X be a topological space.
 - ▶ We say that X has a *countable basis at* $x \in X$ if there is a collection of nbhds $\{B_n\}_{n \in \mathbb{N}}$ of x s.t. if U is any nbhd of x there exists an $n \in \mathbb{N}$ s.t. $B_n \subset U$.

Sequences and Convergence

- ▶ **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.
- ▶ *Proof.* Assume $x \neq y$. Choose U and V disjoint nbhds of x and y respectively. By definition of convergence, we get $N_U, N_V > 0$ such that $x_n \in U$ for all $n > N_U$ and $x_n \in V$ for all $n > N_V$. For $n > \max(N_U, N_V)$ we therefore have $x_n \in U \cap V = \emptyset$, which is a contradiction.
- ▶ **Definition.** Let X be a topological space.
 - ▶ We say that X has a *countable basis at* $x \in X$ if there is a collection of nbhds $\{B_n\}_{n \in \mathbb{N}}$ of x s.t. if U is any nbhd of x there exists an $n \in \mathbb{N}$ s.t. $B_n \subset U$.
 - ▶ X is called *first-countable* if it has a countable basis at x for all $x \in X$.

Sequences and Convergence

- ▶ **Proposition 4.7.** Let X be Hausdorff. If $x_n \rightarrow x$ and $x_n \rightarrow y$ in X , then $x = y$.
- ▶ *Proof.* Assume $x \neq y$. Choose U and V disjoint nbhds of x and y respectively. By definition of convergence, we get $N_U, N_V > 0$ such that $x_n \in U$ for all $n > N_U$ and $x_n \in V$ for all $n > N_V$. For $n > \max(N_U, N_V)$ we therefore have $x_n \in U \cap V = \emptyset$, which is a contradiction.
- ▶ **Definition.** Let X be a topological space.
 - ▶ We say that X has a *countable basis* at $x \in X$ if there is a collection of nbhds $\{B_n\}_{n \in \mathbb{N}}$ of x s.t. if U is any nbhd of x there exists an $n \in \mathbb{N}$ s.t. $B_n \subset U$.
 - ▶ X is called *first-countable* if it has a countable basis at x for all $x \in X$.
- ▶ **Example.** All metric spaces are first-countable.

Sequences and Convergence

- ▶ **Lemma 4.8.** (The sequence lemma) Let $A \subset X$. If there is a sequence in A that converges to x then $x \in \bar{A}$. The converse holds if X is first-countable.

Sequences and Convergence

- ▶ **Lemma 4.8.** (The sequence lemma) Let $A \subset X$. If there is a sequence in A that converges to x then $x \in \bar{A}$. The converse holds if X is first-countable.
- ▶ *Proof.* Suppose $x_n \rightarrow x$ and $x_n \in A$ for all n . Let U be a nbhd of x ; then there is an $N > 0$ s.t. $x_n \in U$ for all $n > N$. This implies that $U \cap A \neq \emptyset$. Hence $x \in \bar{A}$.

Sequences and Convergence

- ▶ **Lemma 4.8.** (The sequence lemma) Let $A \subset X$. If there is a sequence in A that converges to x then $x \in \bar{A}$. The converse holds if X is first-countable.
- ▶ *Proof.* Suppose $x_n \rightarrow x$ and $x_n \in A$ for all n . Let U be a nbhd of x ; then there is an $N > 0$ s.t. $x_n \in U$ for all $n > N$. This implies that $U \cap A \neq \emptyset$. Hence $x \in \bar{A}$.
- ▶ Suppose X is first-countable and $x \in \bar{A}$. Let us show that there is a sequence $\{x_n\}$ in A with $x_n \rightarrow x$.

Sequences and Convergence

- ▶ **Lemma 4.8.** (The sequence lemma) Let $A \subset X$. If there is a sequence in A that converges to x then $x \in \bar{A}$. The converse holds if X is first-countable.
- ▶ *Proof.* Suppose $x_n \rightarrow x$ and $x_n \in A$ for all n . Let U be a nbhd of x ; then there is an $N > 0$ s.t. $x_n \in U$ for all $n > N$. This implies that $U \cap A \neq \emptyset$. Hence $x \in \bar{A}$.
- ▶ Suppose X is first-countable and $x \in \bar{A}$. Let us show that there is a sequence $\{x_n\}$ in A with $x_n \rightarrow x$.
 - ▶ Let $\{B_n\}$ be a countable basis at x and define for every $n \in \mathbb{N}$ an open nbhd $U_n = \bigcap_{k=1}^n B_k$ of x . Since $x \in \bar{A}$, $U_n \cap A \neq \emptyset$.

Sequences and Convergence

- ▶ **Lemma 4.8.** (The sequence lemma) Let $A \subset X$. If there is a sequence in A that converges to x then $x \in \bar{A}$. The converse holds if X is first-countable.
- ▶ *Proof.* Suppose $x_n \rightarrow x$ and $x_n \in A$ for all n . Let U be a nbhd of x ; then there is an $N > 0$ s.t. $x_n \in U$ for all $n > N$. This implies that $U \cap A \neq \emptyset$. Hence $x \in \bar{A}$.
- ▶ Suppose X is first-countable and $x \in \bar{A}$. Let us show that there is a sequence $\{x_n\}$ in A with $x_n \rightarrow x$.
 - ▶ Let $\{B_n\}$ be a countable basis at x and define for every $n \in \mathbb{N}$ an open nbhd $U_n = \bigcap_{k=1}^n B_k$ of x . Since $x \in \bar{A}$, $U_n \cap A \neq \emptyset$.
 - ▶ Choose $x_n \in U_n \cap A$ for every n . We claim that $x_n \rightarrow x$. To see this, let U be any nbhd of x . Since X is first-countable, there is an $N \in \mathbb{N}$ s.t. $B_N \subset U$. For all $n > N$, we have $x_n \in U_n \subset U_N \subset B_N \subset U$ which means that $x_n \rightarrow x$.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
 - ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
 - ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

- ▶ Let $\mathbf{0}$ be the origin in \mathbb{R}^ω , that is, the point $(0, 0, \dots)$ each of whose coordinates is 0.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
 - ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

- ▶ Let $\mathbf{0}$ be the origin in \mathbb{R}^ω , that is, the point $(0, 0, \dots)$ each of whose coordinates is 0.
 - ▶ Claim 1: In the box topology $\mathbf{0} \in \overline{A}$.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
- ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

- ▶ Let $\mathbf{0}$ be the origin in \mathbb{R}^ω , that is, the point $(0, 0, \dots)$ each of whose coordinates is 0.
- ▶ Claim 1: In the box topology $\mathbf{0} \in \overline{A}$.
 - ▶ Let U be any neighborhood of $\mathbf{0}$. Then there exists a basis element B such that $\mathbf{0} \in B \subset U$.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
- ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

- ▶ Let $\mathbf{0}$ be the origin in \mathbb{R}^ω , that is, the point $(0, 0, \dots)$ each of whose coordinates is 0.
- ▶ Claim 1: In the box topology $\mathbf{0} \in \overline{A}$.
 - ▶ Let U be any neighborhood of $\mathbf{0}$. Then there exists a basis element B such that $\mathbf{0} \in B \subset U$.
 - ▶ Without loss of generality we can assume that

$$B = (-a_1, a_1) \times (-a_2, a_2) \times \cdots \times (-a_n, a_n) \times \cdots$$

where $a_i > 0$.

Sequences and Convergence

- ▶ **Remark.** We will show that the sequence lemma does not hold for \mathbb{R}^ω in the box topology.
- ▶ Let A be the subset of \mathbb{R}^ω consisting of those points all of whose coordinates are positive:

$$A = \{(x_1, x_2, \dots) \mid x_i > 0 \quad \forall i \in \mathbb{Z}_+\}.$$

- ▶ Let $\mathbf{0}$ be the origin in \mathbb{R}^ω , that is, the point $(0, 0, \dots)$ each of whose coordinates is 0.
- ▶ Claim 1: In the box topology $\mathbf{0} \in \overline{A}$.
 - ▶ Let U be any neighborhood of $\mathbf{0}$. Then there exists a basis element B such that $\mathbf{0} \in B \subset U$.
 - ▶ Without loss of generality we can assume that

$$B = (-a_1, a_1) \times (-a_2, a_2) \times \cdots \times (-a_n, a_n) \times \cdots$$

where $a_i > 0$.

- ▶ The sequence $(\frac{1}{2}a_1, \frac{1}{2}a_2, \dots, \frac{1}{2}a_n, \dots)$ belongs to $A \cap B$, so B intersects A . Hence U also intersects A .

Sequences and Convergence

- ▶ Claim 2: There is no sequence of points in A converging to $\mathbf{0}$.

Sequences and Convergence

- ▶ Claim 2: There is no sequence of points in A converging to $\mathbf{0}$.
 - ▶ Let (\mathbf{a}_n) be a sequence of points in A , where

$$\mathbf{a}_n = (x_{1n}, x_{2n}, \dots, x_{in}, \dots).$$

Sequences and Convergence

- ▶ Claim 2: There is no sequence of points in A converging to $\mathbf{0}$.
 - ▶ Let (\mathbf{a}_n) be a sequence of points in A , where

$$\mathbf{a}_n = (x_{1n}, x_{2n}, \dots, x_{in}, \dots).$$

- ▶ Consider a basis element B for the box topology on \mathbb{R}^ω :

$$B = (-x_{11}, x_{11}) \times (-x_{22}, x_{22}) \times \dots$$

Sequences and Convergence

- ▶ Claim 2: There is no sequence of points in A converging to $\mathbf{0}$.
 - ▶ Let (\mathbf{a}_n) be a sequence of points in A , where

$$\mathbf{a}_n = (x_{1n}, x_{2n}, \dots, x_{in}, \dots).$$

- ▶ Consider a basis element B for the box topology on \mathbb{R}^ω :

$$B = (-x_{11}, x_{11}) \times (-x_{22}, x_{22}) \times \dots$$

- ▶ Then B contains $\mathbf{0}$, but it contains no member of the sequence (\mathbf{a}_n) since the n th coordinate x_{nn} of \mathbf{a}_n does not belong to the interval $(-x_{nn}, x_{nn})$.

Sequences and Convergence

- **Theorem 4.9.** Let X and Y be topological spaces. If $f : X \rightarrow Y$ be continuous, then $x_n \rightarrow x$ in X implies that $f(x_n) \rightarrow f(x)$ in Y . The converse holds if X is first-countable; that is, if $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$ for all convergent sequences $\{x_n\}$, then f is continuous.

Sequences and Convergence

- ▶ **Theorem 4.9.** Let X and Y be topological spaces. If $f : X \rightarrow Y$ be continuous, then $x_n \rightarrow x$ in X implies that $f(x_n) \rightarrow f(x)$ in Y . The converse holds if X is first-countable; that is, if $x_n \rightarrow x$ implies that $f(x_n) \rightarrow f(x)$ for all convergent sequences $\{x_n\}$, then f is continuous.
- ▶ *Proof.* Suppose f is continuous and $\{x_n\}$ is a sequence with $x_n \rightarrow x$. Let us show that $f(x_n) \rightarrow f(x)$. Let $U \subset Y$ be a nbhd of $f(x)$. Then $f^{-1}(U)$ is a nbhd of x . Since $x_n \rightarrow x$, we can choose an $N > 0$ such that $x_n \in f^{-1}(U)$ for all $n > N$. Thus $f(x_n) \in U$ for all $n > N$, so $f(x_n) \rightarrow f(x)$.

Sequences and Convergence

- Suppose that X is first-countable and that $f(x_n) \rightarrow f(x)$ whenever $x_n \rightarrow x$.

Sequences and Convergence

- ▶ Suppose that X is first-countable and that $f(x_n) \rightarrow f(x)$ whenever $x_n \rightarrow x$.
 - ▶ Let $B \subset Y$ be a closed set and $A = f^{-1}(B)$. Let us show that $\overline{A} = A$ (i.e. A is closed), which means that f is continuous.

Sequences and Convergence

- ▶ Suppose that X is first-countable and that $f(x_n) \rightarrow f(x)$ whenever $x_n \rightarrow x$.
 - ▶ Let $B \subset Y$ be a closed set and $A = f^{-1}(B)$. Let us show that $\overline{A} = A$ (i.e. A is closed), which means that f is continuous.
 - ▶ Let $x \in \overline{A}$ be arbitrary. Then by Lemma 4.8 (the sequence lemma), there is a sequence $\{x_n\}$ with $x_n \in A$ such that $x_n \rightarrow x$. Since $f(x_n) \in B$ and $f(x_n) \rightarrow f(x)$, we have $f(x) \in \overline{B} = B$. Hence $x \in f^{-1}(B) = A$.