

## §2. Topological Spaces

Math 4341 (Topology)

## 2.1. Definitions & Examples

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .
- ▶ If  $\mathcal{T}$  is a topology on  $X$ , then the pair  $(X, \mathcal{T})$  is called a *topological space*. A set  $U \in \mathcal{T}$  is called *open*.

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .
- ▶ If  $\mathcal{T}$  is a topology on  $X$ , then the pair  $(X, \mathcal{T})$  is called a *topological space*. A set  $U \in \mathcal{T}$  is called *open*.
- ▶ **Remark.** By induction (T3) is equivalent to “If  $U, V \in \mathcal{T}$  then  $U \cap V \in \mathcal{T}$ .”



## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .
- ▶ If  $\mathcal{T}$  is a topology on  $X$ , then the pair  $(X, \mathcal{T})$  is called a *topological space*. A set  $U \in \mathcal{T}$  is called *open*.
- ▶ **Remark.** By induction (T3) is equivalent to “If  $U, V \in \mathcal{T}$  then  $U \cap V \in \mathcal{T}$ .”
- ▶ Any set  $X$  can be given a topology in at least two ways:

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .
- ▶ If  $\mathcal{T}$  is a topology on  $X$ , then the pair  $(X, \mathcal{T})$  is called a *topological space*. A set  $U \in \mathcal{T}$  is called *open*.
- ▶ **Remark.** By induction (T3) is equivalent to “If  $U, V \in \mathcal{T}$  then  $U \cap V \in \mathcal{T}$ .”
- ▶ Any set  $X$  can be given a topology in at least two ways:
  - ▶ Let  $\mathcal{T} = \{\emptyset, X\} \subset \mathcal{P}(X)$ . Then  $\mathcal{T}$  is a topology called the *trivial topology*.

## 2.1. Definitions & Examples

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{T} \subset \mathcal{P}(X)$  be a collection of subsets of  $X$ . Then  $\mathcal{T}$  is called a *topology* if
  - ▶ (T1):  $\emptyset \in \mathcal{T}$  and  $X \in \mathcal{T}$ ,
  - ▶ (T2): If  $U_i \in \mathcal{T}$  for all  $i \in I$ , then  $\bigcup_{i \in I} U_i \in \mathcal{T}$ ,
  - ▶ (T3): If  $U_1, \dots, U_n \in \mathcal{T}$ , then  $U_1 \cap U_2 \cap \dots \cap U_n \in \mathcal{T}$ .
- ▶ If  $\mathcal{T}$  is a topology on  $X$ , then the pair  $(X, \mathcal{T})$  is called a *topological space*. A set  $U \in \mathcal{T}$  is called *open*.
- ▶ **Remark.** By induction (T3) is equivalent to “If  $U, V \in \mathcal{T}$  then  $U \cap V \in \mathcal{T}$ .”
- ▶ Any set  $X$  can be given a topology in at least two ways:
  - ▶ Let  $\mathcal{T} = \{\emptyset, X\} \subset \mathcal{P}(X)$ . Then  $\mathcal{T}$  is a topology called *the trivial topology*.
  - ▶ Let  $\mathcal{T} = \mathcal{P}(X)$ . Then  $\mathcal{T}$  is a topology called *the discrete topology*.

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\}$ ,

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\},$
  - ▶  $\mathcal{T}_2 = \{\emptyset, \{a\}, X\},$

## 2.1. Definitions & Examples

- **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
- $\mathcal{T}_1 = \{\emptyset, X\},$
  - $\mathcal{T}_2 = \{\emptyset, \{a\}, X\},$
  - $\mathcal{T}_3 = \{\emptyset, \{b\}, X\},$

## 2.1. Definitions & Examples

- **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
- $\mathcal{T}_1 = \{\emptyset, X\},$
  - $\mathcal{T}_2 = \{\emptyset, \{a\}, X\},$
  - $\mathcal{T}_3 = \{\emptyset, \{b\}, X\},$
  - $\mathcal{T}_4 = \{\emptyset, \{a\}, \{b\}, X\}.$



## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\}$ ,
  - ▶  $\mathcal{T}_2 = \{\emptyset, \{a\}, X\}$ ,
  - ▶  $\mathcal{T}_3 = \{\emptyset, \{b\}, X\}$ ,
  - ▶  $\mathcal{T}_4 = \{\emptyset, \{a\}, \{b\}, X\}$ .
- ▶ **Definition.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be topologies on a set  $X$ .
  - ▶ If  $\mathcal{T} \subset \mathcal{T}'$  then we say that  $\mathcal{T}$  is *coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *finer* than  $\mathcal{T}$ .

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\}$ ,
  - ▶  $\mathcal{T}_2 = \{\emptyset, \{a\}, X\}$ ,
  - ▶  $\mathcal{T}_3 = \{\emptyset, \{b\}, X\}$ ,
  - ▶  $\mathcal{T}_4 = \{\emptyset, \{a\}, \{b\}, X\}$ .
- ▶ **Definition.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be topologies on a set  $X$ .
  - ▶ If  $\mathcal{T} \subset \mathcal{T}'$  then we say that  $\mathcal{T}$  is *coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *finer* than  $\mathcal{T}$ .
  - ▶ If  $\mathcal{T} \subsetneq \mathcal{T}'$ , we say that  $\mathcal{T}$  is *strictly coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *strictly finer* than  $\mathcal{T}$ .

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\}$ ,
  - ▶  $\mathcal{T}_2 = \{\emptyset, \{a\}, X\}$ ,
  - ▶  $\mathcal{T}_3 = \{\emptyset, \{b\}, X\}$ ,
  - ▶  $\mathcal{T}_4 = \{\emptyset, \{a\}, \{b\}, X\}$ .
- ▶ **Definition.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be topologies on a set  $X$ .
  - ▶ If  $\mathcal{T} \subset \mathcal{T}'$  then we say that  $\mathcal{T}$  is *coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *finer* than  $\mathcal{T}$ .
  - ▶ If  $\mathcal{T} \subsetneq \mathcal{T}'$ , we say that  $\mathcal{T}$  is *strictly coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *strictly finer* than  $\mathcal{T}$ .
  - ▶ If either  $\mathcal{T} \subset \mathcal{T}'$  or  $\mathcal{T}' \subset \mathcal{T}$ , we say that  $\mathcal{T}$  and  $\mathcal{T}'$  are *comparable*.

## 2.1. Definitions & Examples

- ▶ **Example.** Let  $X = \{a, b\}$ . There are 4 topologies on  $X$ :
  - ▶  $\mathcal{T}_1 = \{\emptyset, X\},$
  - ▶  $\mathcal{T}_2 = \{\emptyset, \{a\}, X\},$
  - ▶  $\mathcal{T}_3 = \{\emptyset, \{b\}, X\},$
  - ▶  $\mathcal{T}_4 = \{\emptyset, \{a\}, \{b\}, X\}.$
- ▶ **Definition.** Let  $\mathcal{T}$  and  $\mathcal{T}'$  be topologies on a set  $X$ .
  - ▶ If  $\mathcal{T} \subset \mathcal{T}'$  then we say that  $\mathcal{T}$  is *coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *finer* than  $\mathcal{T}$ .
  - ▶ If  $\mathcal{T} \subsetneq \mathcal{T}'$ , we say that  $\mathcal{T}$  is *strictly coarser* than  $\mathcal{T}'$ , and that  $\mathcal{T}'$  is *strictly finer* than  $\mathcal{T}$ .
  - ▶ If either  $\mathcal{T} \subset \mathcal{T}'$  or  $\mathcal{T}' \subset \mathcal{T}$ , we say that  $\mathcal{T}$  and  $\mathcal{T}'$  are *comparable*.
- ▶ In the example above,  $\mathcal{T}_2$  is strictly coarser than  $\mathcal{T}_4$ , but  $\mathcal{T}_2$  and  $\mathcal{T}_3$  are not comparable.

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,
  - ▶ (T2'): If  $C_i$  is closed for all  $i \in I$ , then  $\bigcap_{i \in I} C_i$  is also closed,



## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,
  - ▶ (T2'): If  $C_i$  is closed for all  $i \in I$ , then  $\bigcap_{i \in I} C_i$  is also closed,
  - ▶ (T3'): If  $C_1, \dots, C_n$  are closed, then  $C_1 \cup C_2 \cup \dots \cup C_n$  is also closed.

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,
  - ▶ (T2'): If  $C_i$  is closed for all  $i \in I$ , then  $\bigcap_{i \in I} C_i$  is also closed,
  - ▶ (T3'): If  $C_1, \dots, C_n$  are closed, then  $C_1 \cup C_2 \cup \dots \cup C_n$  is also closed.
- ▶ The proof is based on De Morgan's laws

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,
  - ▶ (T2'): If  $C_i$  is closed for all  $i \in I$ , then  $\bigcap_{i \in I} C_i$  is also closed,
  - ▶ (T3'): If  $C_1, \dots, C_n$  are closed, then  $C_1 \cup C_2 \cup \dots \cup C_n$  is also closed.
- ▶ The proof is based on De Morgan's laws
  - ▶  $X \setminus \bigcap_{i \in I} C_i = \bigcup_{i \in I} (X \setminus C_i),$

## 2.1. Definitions & Examples

- ▶ **Definition.** A subset  $A \subset X$  of a topological space is called *closed* if  $A^c := X \setminus A$  is open.
- ▶ **Proposition 2.1.** In a topological space  $X$ , we have
  - ▶ (T1'):  $\emptyset$  and  $X$  are closed,
  - ▶ (T2'): If  $C_i$  is closed for all  $i \in I$ , then  $\bigcap_{i \in I} C_i$  is also closed,
  - ▶ (T3'): If  $C_1, \dots, C_n$  are closed, then  $C_1 \cup C_2 \cup \dots \cup C_n$  is also closed.
- ▶ The proof is based on De Morgan's laws
  - ▶  $X \setminus \bigcap_{i \in I} C_i = \bigcup_{i \in I} (X \setminus C_i),$
  - ▶  $X \setminus \bigcup_{i \in I} C_i = \bigcap_{i \in I} (X \setminus C_i).$

## 2.2. Basis for a topology

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{B} \subset \mathcal{P}(X)$  be any collection of subsets of  $X$ . Then  $\mathcal{B}$  is called a *basis* for a topology on  $X$  if

## 2.2. Basis for a topology

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{B} \subset \mathcal{P}(X)$  be any collection of subsets of  $X$ . Then  $\mathcal{B}$  is called a *basis* for a topology on  $X$  if
  - ▶ (B1): For each  $x \in X$ , there is a  $B \in \mathcal{B}$  such that  $x \in B$ ,

## 2.2. Basis for a topology

- **Definition.** Let  $X$  be a set, and let  $\mathcal{B} \subset \mathcal{P}(X)$  be any collection of subsets of  $X$ . Then  $\mathcal{B}$  is called a *basis* for a topology on  $X$  if
- (B1): For each  $x \in X$ , there is a  $B \in \mathcal{B}$  such that  $x \in B$ ,
  - (B2): If  $x \in B_1 \cap B_2$  for  $B_1, B_2 \in \mathcal{B}$ , then there is a  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subset B_1 \cap B_2$ .

## 2.2. Basis for a topology

- ▶ **Definition.** Let  $X$  be a set, and let  $\mathcal{B} \subset \mathcal{P}(X)$  be any collection of subsets of  $X$ . Then  $\mathcal{B}$  is called a *basis* for a topology on  $X$  if
  - ▶ (B1): For each  $x \in X$ , there is a  $B \in \mathcal{B}$  such that  $x \in B$ ,
  - ▶ (B2): If  $x \in B_1 \cap B_2$  for  $B_1, B_2 \in \mathcal{B}$ , then there is a  $B_3 \in \mathcal{B}$  such that  $x \in B_3 \subset B_1 \cap B_2$ .
- ▶ If  $\mathcal{B}$  is a basis, we define  $\mathcal{T}_{\mathcal{B}}$ , the *topology generated by  $\mathcal{B}$* , by declaring that  $U \in \mathcal{T}_{\mathcal{B}}$  if for every  $x \in U$ , there is a basis element  $B \in \mathcal{B}$  such that  $x \in B \subset U$ .



## 2.2. Basis for a topology

- ▶ **Lemma 2.2.**  $\mathcal{T}_\mathcal{B} \subset \mathcal{P}(X)$  is a topology.

## 2.2. Basis for a topology

- ▶ **Lemma 2.2.**  $\mathcal{T}_{\mathcal{B}} \subset \mathcal{P}(X)$  is a topology.
- ▶ *Proof.* We show that  $\mathcal{T}_{\mathcal{B}}$  satisfies the properties (T1)–(T3).

## 2.2. Basis for a topology

- ▶ **Lemma 2.2.**  $\mathcal{T}_\mathcal{B} \subset \mathcal{P}(X)$  is a topology.
- ▶ *Proof.* We show that  $\mathcal{T}_\mathcal{B}$  satisfies the properties (T1)–(T3).
  - ▶ (T1) follows from (B1).

## 2.2. Basis for a topology

- ▶ **Lemma 2.2.**  $\mathcal{T}_{\mathcal{B}} \subset \mathcal{P}(X)$  is a topology.
- ▶ *Proof.* We show that  $\mathcal{T}_{\mathcal{B}}$  satisfies the properties (T1)–(T3).
  - ▶ (T1) follows from (B1).
  - ▶ (T2): Suppose  $U_i \in \mathcal{T}_{\mathcal{B}}$  for all  $i \in I$ . Let  $x \in \bigcup_{i \in I} U_i$ . Then there exists an  $i \in I$  so that  $x \in U_i$ , and since  $U_i \in \mathcal{T}_{\mathcal{B}}$  we get a basis element  $B \in \mathcal{B}$  so that  $x \in B \subset U_i \subset \bigcup_{i \in I} U_i$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.2.**  $\mathcal{T}_\mathcal{B} \subset \mathcal{P}(X)$  is a topology.
- ▶ *Proof.* We show that  $\mathcal{T}_\mathcal{B}$  satisfies the properties (T1)–(T3).
  - ▶ (T1) follows from (B1).
  - ▶ (T2): Suppose  $U_i \in \mathcal{T}_\mathcal{B}$  for all  $i \in I$ . Let  $x \in \bigcup_{i \in I} U_i$ . Then there exists an  $i \in I$  so that  $x \in U_i$ , and since  $U_i \in \mathcal{T}_\mathcal{B}$  we get a basis element  $B \in \mathcal{B}$  so that  $x \in B \subset U_i \subset \bigcup_{i \in I} U_i$ .
  - ▶ (T3): Suppose  $U_1, U_2 \in \mathcal{T}_\mathcal{B}$ . Let  $x \in U_1 \cap U_2$ . Then  $x \in U_1$  and  $x \in U_2$ , so we get  $B_1, B_2 \in \mathcal{B}$  so that  $x \in B_1 \subset U_1$  and  $x \in B_2 \subset U_2$ . Now, by (B2) we get a  $B_3 \in \mathcal{B}$  so that  $x \in B_3 \subset B_1 \cap B_2 \subset U_1 \cap U_2$ .

## 2.2. Basis for a topology

- **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .
- ▶ *Proof.* There are two things to show.

## 2.2. Basis for a topology

- ▶ **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .
- ▶ *Proof.* There are two things to show.
  - ▶ Suppose  $U = \bigcup_{i \in I} B_i$  for some  $B_i \in \mathcal{B}$ . Let  $x \in U$ . Then there is an  $i \in I$  so that  $x \in B_i \subset U$ . This shows that  $U \in \mathcal{T}_{\mathcal{B}}$ .



## 2.2. Basis for a topology

- ▶ **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .
- ▶ *Proof.* There are two things to show.
  - ▶ Suppose  $U = \bigcup_{i \in I} B_i$  for some  $B_i \in \mathcal{B}$ . Let  $x \in U$ . Then there is an  $i \in I$  so that  $x \in B_i \subset U$ . This shows that  $U \in \mathcal{T}_{\mathcal{B}}$ .
  - ▶ Suppose  $U \in \mathcal{T}_{\mathcal{B}}$ . For every  $x \in U$ , choose a basis element  $B_x$  so that  $x \in B_x \subset U$ . We now claim that  $U = \bigcup_{x \in U} B_x$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .
- ▶ *Proof.* There are two things to show.
  - ▶ Suppose  $U = \bigcup_{i \in I} B_i$  for some  $B_i \in \mathcal{B}$ . Let  $x \in U$ . Then there is an  $i \in I$  so that  $x \in B_i \subset U$ . This shows that  $U \in \mathcal{T}_{\mathcal{B}}$ .
  - ▶ Suppose  $U \in \mathcal{T}_{\mathcal{B}}$ . For every  $x \in U$ , choose a basis element  $B_x$  so that  $x \in B_x \subset U$ . We now claim that  $U = \bigcup_{x \in U} B_x$ .
    - ▶ Take any  $y \in U$ . Then  $y \in B_y$  and  $B_y \subset \bigcup_{x \in U} B_x$ , so  $y$  is an element of the union.

## 2.2. Basis for a topology

- ▶ **Lemma 2.3.** Let  $\mathcal{B}$  be the basis for a topology on a set  $X$ . Then  $U \in \mathcal{T}_{\mathcal{B}}$  if and only if  $U = \bigcup_{i \in I} B_i$  for some sets  $B_i \in \mathcal{B}$ . That is,  $\mathcal{T}_{\mathcal{B}}$  consists of all unions of elements from  $\mathcal{B}$ .
- ▶ *Proof.* There are two things to show.
  - ▶ Suppose  $U = \bigcup_{i \in I} B_i$  for some  $B_i \in \mathcal{B}$ . Let  $x \in U$ . Then there is an  $i \in I$  so that  $x \in B_i \subset U$ . This shows that  $U \in \mathcal{T}_{\mathcal{B}}$ .
  - ▶ Suppose  $U \in \mathcal{T}_{\mathcal{B}}$ . For every  $x \in U$ , choose a basis element  $B_x$  so that  $x \in B_x \subset U$ . We now claim that  $U = \bigcup_{x \in U} B_x$ .
    - ▶ Take any  $y \in U$ . Then  $y \in B_y$  and  $B_y \subset \bigcup_{x \in U} B_x$ , so  $y$  is an element of the union.
    - ▶ Take any  $y \in \bigcup_{x \in U} B_x$ . Then there exists a  $z \in U$  so that  $y \in B_z$ , but by our choices of the basis elements, we have that  $B_z \subset U$ , so  $y \in B_z \subset U$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.
- ▶ The topology  $\mathcal{T}_{\mathcal{C}}$  generated by  $\mathcal{C}$  is actually  $\mathcal{T}$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.
    - ▶ (B1): Let  $x \in X$ . Since  $X \in \mathcal{T}$ , by definition of  $\mathcal{C}$  we get a  $C \in \mathcal{C}$  so that  $x \in C \subset X$ .
- ▶ The topology  $\mathcal{T}_{\mathcal{C}}$  generated by  $\mathcal{C}$  is actually  $\mathcal{T}$ .



## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.
    - ▶ (B1): Let  $x \in X$ . Since  $X \in \mathcal{T}$ , by definition of  $\mathcal{C}$  we get a  $C \in \mathcal{C}$  so that  $x \in C \subset X$ .
    - ▶ (B2): Let  $x \in C_1 \cap C_2$  for  $C_1, C_2 \in \mathcal{C}$ . Since  $C_1$  and  $C_2$  are open sets, so is  $C_1 \cap C_2$ . Therefore we get a  $C \in \mathcal{C}$  so that  $x \in C \subset C_1 \cap C_2$ .
  - ▶ The topology  $\mathcal{T}_{\mathcal{C}}$  generated by  $\mathcal{C}$  is actually  $\mathcal{T}$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.
    - ▶ (B1): Let  $x \in X$ . Since  $X \in \mathcal{T}$ , by definition of  $\mathcal{C}$  we get a  $C \in \mathcal{C}$  so that  $x \in C \subset X$ .
    - ▶ (B2): Let  $x \in C_1 \cap C_2$  for  $C_1, C_2 \in \mathcal{C}$ . Since  $C_1$  and  $C_2$  are open sets, so is  $C_1 \cap C_2$ . Therefore we get a  $C \in \mathcal{C}$  so that  $x \in C \subset C_1 \cap C_2$ .
  - ▶ The topology  $\mathcal{T}_{\mathcal{C}}$  generated by  $\mathcal{C}$  is actually  $\mathcal{T}$ .
    - ▶ Take any  $U \in \mathcal{T}_{\mathcal{C}}$ . By Lemma 2.3,  $U$  is a union of elements of  $\mathcal{C}$ . Since  $\mathcal{C} \subset \mathcal{T}$ , it follows from (T2) that  $U \in \mathcal{T}$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.4.** Let  $(X, \mathcal{T})$  be a topological space. Let  $\mathcal{C} \subset \mathcal{T}$  be a collection of open sets on  $X$  with the following property: for each set  $U \in \mathcal{T}$  and each  $x \in U$  there is a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Then  $\mathcal{C}$  is a basis for  $\mathcal{T}$ .
- ▶ *Proof.* There are two things to show.
  - ▶  $\mathcal{C}$  is a basis for a topology.
    - ▶ (B1): Let  $x \in X$ . Since  $X \in \mathcal{T}$ , by definition of  $\mathcal{C}$  we get a  $C \in \mathcal{C}$  so that  $x \in C \subset X$ .
    - ▶ (B2): Let  $x \in C_1 \cap C_2$  for  $C_1, C_2 \in \mathcal{C}$ . Since  $C_1$  and  $C_2$  are open sets, so is  $C_1 \cap C_2$ . Therefore we get a  $C \in \mathcal{C}$  so that  $x \in C \subset C_1 \cap C_2$ .
  - ▶ The topology  $\mathcal{T}_{\mathcal{C}}$  generated by  $\mathcal{C}$  is actually  $\mathcal{T}$ .
    - ▶ Take any  $U \in \mathcal{T}_{\mathcal{C}}$ . By Lemma 2.3,  $U$  is a union of elements of  $\mathcal{C}$ . Since  $\mathcal{C} \subset \mathcal{T}$ , it follows from (T2) that  $U \in \mathcal{T}$ .
    - ▶ Take any  $U \in \mathcal{T}$ . By definition of  $\mathcal{C}$ , for any  $x \in U$  we can find a  $C \in \mathcal{C}$  so that  $x \in C \subset U$ . Hence  $U \in \mathcal{T}_{\mathcal{C}}$ .

## 2.2. Basis for a topology

- For  $x \in \mathbb{R}^n$  and  $r > 0$ , let

$$B(x, r) = \{y \in \mathbb{R}^n : \|x - y\| < r\}$$

be the open ball centered in  $x$  with radius  $r$ .

## 2.2. Basis for a topology

- ▶ For  $x \in \mathbb{R}^n$  and  $r > 0$ , let

$$B(x, r) = \{y \in \mathbb{R}^n : \|x - y\| < r\}$$

be the open ball centered in  $x$  with radius  $r$ .

- ▶ **Proposition 2.5.** The collection

$$\mathcal{B} = \{B(x, r) \mid x \in \mathbb{R}^n, r > 0\}$$

is the basis for a topology on  $\mathbb{R}^n$ . The resulting topology  $\mathcal{T}_{\mathcal{B}}$  is called the standard topology and its open sets are exactly the open sets in analysis/calculus.

## 2.2. Basis for a topology

- ▶ **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:

## 2.2. Basis for a topology

- ▶ **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:
  - ▶ (1) The topology  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .

## 2.2. Basis for a topology

- **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:
- (1) The topology  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .
  - (2) For every  $x \in X$  and each basis element  $B \in \mathcal{B}$  satisfying  $x \in B$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .



## 2.2. Basis for a topology

- ▶ **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:
  - ▶ (1) The topology  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .
  - ▶ (2) For every  $x \in X$  and each basis element  $B \in \mathcal{B}$  satisfying  $x \in B$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .
- ▶ *Proof.* There are two things to prove.

## 2.2. Basis for a topology

- ▶ **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:
  - ▶ (1) The topology  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .
  - ▶ (2) For every  $x \in X$  and each basis element  $B \in \mathcal{B}$  satisfying  $x \in B$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .
- ▶ *Proof.* There are two things to prove.
  - ▶ Suppose  $\mathcal{T} \subset \mathcal{T}'$ . Let  $x$  in  $X$ , and let  $B \in \mathcal{B}$  satisfy  $x \in B$ . Then since  $B \in \mathcal{T}$ , we have  $B \in \mathcal{T}'$ . Since  $\mathcal{T}' = \mathcal{T}_{\mathcal{B}'}$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.6.** Let  $X$  be a set, and let  $\mathcal{B}$  and  $\mathcal{B}'$  be bases for topologies  $\mathcal{T}$  and  $\mathcal{T}'$  respectively; both on  $X$ . TFAE:
  - ▶ (1) The topology  $\mathcal{T}'$  is finer than  $\mathcal{T}$ .
  - ▶ (2) For every  $x \in X$  and each basis element  $B \in \mathcal{B}$  satisfying  $x \in B$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .
- ▶ *Proof.* There are two things to prove.
  - ▶ Suppose  $\mathcal{T} \subset \mathcal{T}'$ . Let  $x$  in  $X$ , and let  $B \in \mathcal{B}$  satisfy  $x \in B$ . Then since  $B \in \mathcal{T}$ , we have  $B \in \mathcal{T}'$ . Since  $\mathcal{T}' = \mathcal{T}_{\mathcal{B}'}$ , there is a basis element  $B' \in \mathcal{B}'$  so that  $x \in B' \subset B$ .
  - ▶ Suppose (2) holds. Let  $U \in \mathcal{T}$ , and let  $x \in U$  be any element. Then there is a  $B \in \mathcal{B}$  with  $x \in B \subset U$ , and by (2) we get  $B' \in \mathcal{B}'$  with  $x \in B' \subset B \subset U$ . This implies that  $U \in \mathcal{T}'$ .

## 2.2. Basis for a topology

- **Example.** We can define a basis for a topology on  $\mathbb{R}$  by letting  $\mathcal{B}_\ell$  consist of all sets of the form

$$\{x \in \mathbb{R} \mid a \leq x < b\},$$

where  $a, b \in \mathbb{R}$  vary. The topology  $\mathcal{T}_\ell$  generated by  $\mathcal{B}_\ell$  is called the *lower limit topology* on  $\mathbb{R}$ , and we write  $\mathbb{R}_\ell = (\mathbb{R}, \mathcal{T}_\ell)$ .

## 2.2. Basis for a topology

- **Example.** We can define a basis for a topology on  $\mathbb{R}$  by letting  $\mathcal{B}_\ell$  consist of all sets of the form

$$\{x \in \mathbb{R} \mid a \leq x < b\},$$

where  $a, b \in \mathbb{R}$  vary. The topology  $\mathcal{T}_\ell$  generated by  $\mathcal{B}_\ell$  is called the *lower limit topology* on  $\mathbb{R}$ , and we write  $\mathbb{R}_\ell = (\mathbb{R}, \mathcal{T}_\ell)$ .

- **Example.** Let  $K = \{1/n \mid n \in \mathbb{N}\} \subset \mathbb{R}$  and let  $\mathcal{B}_K$  consist of all open intervals as well as all sets of the form  $(a, b) \setminus K$ . Then  $\mathcal{B}_K$  is a basis and the topology  $\mathcal{T}_K$  that it generates is called the *K-topology* on  $\mathbb{R}$ . We write  $\mathbb{R}_K = (\mathbb{R}, \mathcal{T}_K)$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.7.** The topologies  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are both strictly finer than the std topology but are not comparable with each other.

## 2.2. Basis for a topology

- ▶ **Lemma 2.7.** The topologies  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are both strictly finer than the std topology but are not comparable with each other.
- ▶ *Proof.* There are three things to prove.

## 2.2. Basis for a topology

- ▶ **Lemma 2.7.** The topologies  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are both strictly finer than the std topology but are not comparable with each other.
- ▶ *Proof.* There are three things to prove.
  - ▶  $\mathbb{R}_\ell \supsetneq \mathbb{R}$ : Let  $x \in \mathbb{R}$  and let  $(a, b)$  contain  $x$ . Since  $[x, b) \subset (a, b)$ , Lemma 2.6 implies that  $\mathbb{R}_\ell \supset \mathbb{R}$ . It is strictly finer because  $[x, b)$  is open in  $\mathbb{R}_\ell$  but not in the std topology: There is no open interval  $B$  so that  $x \in B \subset [x, b)$ .



## 2.2. Basis for a topology

- ▶ **Lemma 2.7.** The topologies  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are both strictly finer than the std topology but are not comparable with each other.
- ▶ *Proof.* There are three things to prove.
  - ▶  $\mathbb{R}_\ell \supsetneq \mathbb{R}$ : Let  $x \in \mathbb{R}$  and let  $(a, b)$  contain  $x$ . Since  $[x, b) \subset (a, b)$ , Lemma 2.6 implies that  $\mathbb{R}_\ell \supset \mathbb{R}$ . It is strictly finer because  $[x, b)$  is open in  $\mathbb{R}_\ell$  but not in the std topology: There is no open interval  $B$  so that  $x \in B \subset [x, b)$ .
  - ▶  $\mathbb{R}_K \supsetneq \mathbb{R}$ : Let  $x \in \mathbb{R}$  and let  $(a, b)$  contain  $x$ . This interval belongs to  $\mathcal{B}_K$  so by Lemma 2.6 we have  $\mathbb{R}_K \supset \mathbb{R}$ . To see that it is strictly finer, consider the set  $U = (-1, 1) \setminus K \in \mathcal{T}_K$ . Then  $0 \in U$  but there is no open interval  $B$  so that  $0 \in B \subset U$ .

## 2.2. Basis for a topology

- ▶ **Lemma 2.7.** The topologies  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are both strictly finer than the std topology but are not comparable with each other.
- ▶ *Proof.* There are three things to prove.
  - ▶  $\mathbb{R}_\ell \supsetneq \mathbb{R}$ : Let  $x \in \mathbb{R}$  and let  $(a, b)$  contain  $x$ . Since  $[x, b) \subset (a, b)$ , Lemma 2.6 implies that  $\mathbb{R}_\ell \supset \mathbb{R}$ . It is strictly finer because  $[x, b)$  is open in  $\mathbb{R}_\ell$  but not in the std topology: There is no open interval  $B$  so that  $x \in B \subset [x, b)$ .
  - ▶  $\mathbb{R}_K \supsetneq \mathbb{R}$ : Let  $x \in \mathbb{R}$  and let  $(a, b)$  contain  $x$ . This interval belongs to  $\mathcal{B}_K$  so by Lemma 2.6 we have  $\mathbb{R}_K \supset \mathbb{R}$ . To see that it is strictly finer, consider the set  $U = (-1, 1) \setminus K \in \mathcal{T}_K$ . Then  $0 \in U$  but there is no open interval  $B$  so that  $0 \in B \subset U$ .
  - ▶  $\mathbb{R}_\ell$  and  $\mathbb{R}_K$  are not comparable with each other: Note that  $U \in \mathcal{T}_K$  but  $U \notin \mathcal{T}_\ell$ , and that  $[1, 2) \in \mathcal{T}_\ell$  but  $[1, 2) \notin \mathcal{T}_K$ .

- **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying

- ▶ **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying
  - ▶ (M1):  $d(x, y) = 0$  if and only if  $x = y$ ,

# Metric spaces

- ▶ **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying
  - ▶ (M1):  $d(x, y) = 0$  if and only if  $x = y$ ,
  - ▶ (M2):  $d(x, y) = d(y, x)$ ,

# Metric spaces

- **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying
- (M1):  $d(x, y) = 0$  if and only if  $x = y$ ,
  - (M2):  $d(x, y) = d(y, x)$ ,
  - (M3): the triangle inequality  $d(x, z) \leq d(x, y) + d(y, z)$ .

# Metric spaces

- ▶ **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying
  - ▶ (M1):  $d(x, y) = 0$  if and only if  $x = y$ ,
  - ▶ (M2):  $d(x, y) = d(y, x)$ ,
  - ▶ (M3): the triangle inequality  $d(x, z) \leq d(x, y) + d(y, z)$ .
- ▶ The function  $d$  is called a *metric*, and  $d(x, y)$  is called the *distance* from  $x$  to  $y$ .

- ▶ **Definition.** A *metric space*  $(X, d)$  is a set  $X$  together with a non-negative function  $d : X \times X \rightarrow \mathbb{R}_{\geq 0}$  satisfying
  - ▶ (M1):  $d(x, y) = 0$  if and only if  $x = y$ ,
  - ▶ (M2):  $d(x, y) = d(y, x)$ ,
  - ▶ (M3): the triangle inequality  $d(x, z) \leq d(x, y) + d(y, z)$ .
- ▶ The function  $d$  is called a *metric*, and  $d(x, y)$  is called the *distance* from  $x$  to  $y$ .
- ▶ For a metric space  $(X, d)$  the open *ball*  $B_d(x, r)$  centered at  $x$ , with radius  $r > 0$ , with respect to the metric  $d$  is defined as

$$B_d(x, r) = \{y \in X \mid d(x, y) < r\}.$$

We will use the open balls to define a topology, called *the metric topology* on any metric space.



- **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ *Proof.* We need to show that  $\mathcal{B}$  satisfies (B1) and (B2).

- ▶ **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ *Proof.* We need to show that  $\mathcal{B}$  satisfies (B1) and (B2).
  - ▶ (B1) follows since  $x \in B_d(x, r)$  for any  $r > 0$ .

- ▶ **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ *Proof.* We need to show that  $\mathcal{B}$  satisfies (B1) and (B2).
  - ▶ (B1) follows since  $x \in B_d(x, r)$  for any  $r > 0$ .
  - ▶ To see (B2), let  $x \in B_d(y_1, r_1) \cap B_d(y_2, r_2)$ .

- ▶ **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ *Proof.* We need to show that  $\mathcal{B}$  satisfies (B1) and (B2).
  - ▶ (B1) follows since  $x \in B_d(x, r)$  for any  $r > 0$ .
  - ▶ To see (B2), let  $x \in B_d(y_1, r_1) \cap B_d(y_2, r_2)$ .
    - ▶ We need to show that there is a  $r > 0$  so that

$$B_d(x, r) \subset B_d(y_1, r_1) \cap B_d(y_2, r_2).$$

- ▶ **Proposition 2.8.** If  $(X, d)$  is a metric space, then the collection

$$\mathcal{B} = \{B_d(x, r) \mid x \in X, r > 0\}$$

is a basis for a topology.

- ▶ *Proof.* We need to show that  $\mathcal{B}$  satisfies (B1) and (B2).
  - ▶ (B1) follows since  $x \in B_d(x, r)$  for any  $r > 0$ .
  - ▶ To see (B2), let  $x \in B_d(y_1, r_1) \cap B_d(y_2, r_2)$ .
    - ▶ We need to show that there is a  $r > 0$  so that

$$B_d(x, r) \subset B_d(y_1, r_1) \cap B_d(y_2, r_2).$$

- ▶ Choose  $r = \min(r_1 - d(x, y_1), r_2 - d(x, y_2))$ . For any  $z \in B_d(x, r)$ , by the triangle inequality we have

$$d(z, y_i) \leq d(z, x) + d(x, y_i) < r + d(x, y_i) \leq r_i$$

for  $i = 1, 2$ . This implies that  $z \in B_d(y_1, r_1) \cap B_d(y_2, r_2)$ .

- **Proposition 2.9.** A set  $U$  is open in the metric topology iff for every point  $x \in U$  there is an  $r > 0$  so that  $B_d(x, r) \subset U$ .

- ▶ **Proposition 2.9.** A set  $U$  is open in the metric topology iff for every point  $x \in U$  there is an  $r > 0$  so that  $B_d(x, r) \subset U$ .
- ▶ *Proof.* The direction  $(\Leftarrow)$  follows from the definition of the topology as induced by the basis of open balls. Suppose now  $U$  is open in the metric topology. Let  $x \in U$ .



- ▶ **Proposition 2.9.** A set  $U$  is open in the metric topology iff for every point  $x \in U$  there is an  $r > 0$  so that  $B_d(x, r) \subset U$ .
- ▶ *Proof.* The direction  $(\Leftarrow)$  follows from the definition of the topology as induced by the basis of open balls. Suppose now  $U$  is open in the metric topology. Let  $x \in U$ .
  - ▶ Since the topology is generated by the basis of open balls, there exists an open ball  $B_d(y, \epsilon)$  so that  $x \in B_d(y, \epsilon) \subset U$ .

- ▶ **Proposition 2.9.** A set  $U$  is open in the metric topology iff for every point  $x \in U$  there is an  $r > 0$  so that  $B_d(x, r) \subset U$ .
- ▶ *Proof.* The direction ( $\Leftarrow$ ) follows from the definition of the topology as induced by the basis of open balls. Suppose now  $U$  is open in the metric topology. Let  $x \in U$ .
  - ▶ Since the topology is generated by the basis of open balls, there exists an open ball  $B_d(y, \epsilon)$  so that  $x \in B_d(y, \epsilon) \subset U$ .
  - ▶ By setting  $r = \epsilon - d(x, y) > 0$  we see that

$$x \in B_d(x, r) \subset B_d(y, \epsilon) \subset U.$$

- ▶ **Proposition 2.9.** A set  $U$  is open in the metric topology iff for every point  $x \in U$  there is an  $r > 0$  so that  $B_d(x, r) \subset U$ .
- ▶ *Proof.* The direction  $(\Leftarrow)$  follows from the definition of the topology as induced by the basis of open balls. Suppose now  $U$  is open in the metric topology. Let  $x \in U$ .
  - ▶ Since the topology is generated by the basis of open balls, there exists an open ball  $B_d(y, \epsilon)$  so that  $x \in B_d(y, \epsilon) \subset U$ .
  - ▶ By setting  $r = \epsilon - d(x, y) > 0$  we see that

$$x \in B_d(x, r) \subset B_d(y, \epsilon) \subset U.$$

- ▶ **Remark.** For the case of  $\mathbb{R}^n$ , we recover the usual condition for a set to be open.

# Metric spaces

- ▶ **Example.** The Euclidean space  $\mathbb{R}^n$  is a metric space with metric  $d(x, y) = \|x - y\|$ .

# Metric spaces

- ▶ **Example.** The Euclidean space  $\mathbb{R}^n$  is a metric space with metric  $d(x, y) = \|x - y\|$ .
- ▶ **Example.** Let  $X$  be any set. We can define a metric on  $X$  by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y, \\ 1 & \text{if } x \neq y. \end{cases}$$

# Metric spaces

- ▶ **Example.** The Euclidean space  $\mathbb{R}^n$  is a metric space with metric  $d(x, y) = \|x - y\|$ .
- ▶ **Example.** Let  $X$  be any set. We can define a metric on  $X$  by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y, \\ 1 & \text{if } x \neq y. \end{cases}$$

- ▶ The metric  $d$  is often called the *discrete metric*. Let's describe the collection of open balls.

# Metric spaces

- ▶ **Example.** The Euclidean space  $\mathbb{R}^n$  is a metric space with metric  $d(x, y) = \|x - y\|$ .
- ▶ **Example.** Let  $X$  be any set. We can define a metric on  $X$  by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y, \\ 1 & \text{if } x \neq y. \end{cases}$$

- ▶ The metric  $d$  is often called the *discrete metric*. Let's describe the collection of open balls.
- ▶ Let  $x \in X$  be arbitrary. If  $r \leq 1$ , then  $B_d(x, r) = \{x\}$  while if  $r > 1$  then  $B_d(x, r) = X$ .

# Metric spaces

- ▶ **Example.** The Euclidean space  $\mathbb{R}^n$  is a metric space with metric  $d(x, y) = \|x - y\|$ .
- ▶ **Example.** Let  $X$  be any set. We can define a metric on  $X$  by

$$d(x, y) = \begin{cases} 0 & \text{if } x = y, \\ 1 & \text{if } x \neq y. \end{cases}$$

- ▶ The metric  $d$  is often called the *discrete metric*. Let's describe the collection of open balls.
- ▶ Let  $x \in X$  be arbitrary. If  $r \leq 1$ , then  $B_d(x, r) = \{x\}$  while if  $r > 1$  then  $B_d(x, r) = X$ .
- ▶ Thus the basis of open balls is

$$\mathcal{B} = \{\{x\} \mid x \in X\} \cup \{X\}.$$

Hence the topology induced by  $d$  is the discrete topology.



# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.

# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.
- ▶ If  $f : X \rightarrow Y$  is a function between two sets, and  $A \subset Y$  is a subset, then we define the *preimage* of  $A$  to be

$$f^{-1}(A) = \{x \in X \mid f(x) \in A\}.$$

# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.
- ▶ If  $f : X \rightarrow Y$  is a function between two sets, and  $A \subset Y$  is a subset, then we define the *preimage* of  $A$  to be

$$f^{-1}(A) = \{x \in X \mid f(x) \in A\}.$$

- ▶ **Proposition 2.10.** The preimage behaves nicely with respect to various operations of sets.

# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.
- ▶ If  $f : X \rightarrow Y$  is a function between two sets, and  $A \subset Y$  is a subset, then we define the *preimage* of  $A$  to be

$$f^{-1}(A) = \{x \in X \mid f(x) \in A\}.$$

- ▶ **Proposition 2.10.** The preimage behaves nicely with respect to various operations of sets.
  - ▶ If  $f : X \rightarrow Y$  and  $\{B_i\}_{i \in I}$  is a family of subsets of  $Y$ , then

$$f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i), \quad f^{-1}\left(\bigcap_{i \in I} B_i\right) = \bigcap_{i \in I} f^{-1}(B_i).$$

# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.
- ▶ If  $f : X \rightarrow Y$  is a function between two sets, and  $A \subset Y$  is a subset, then we define the *preimage* of  $A$  to be

$$f^{-1}(A) = \{x \in X \mid f(x) \in A\}.$$

- ▶ **Proposition 2.10.** The preimage behaves nicely with respect to various operations of sets.

- ▶ If  $f : X \rightarrow Y$  and  $\{B_i\}_{i \in I}$  is a family of subsets of  $Y$ , then

$$f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i), \quad f^{-1}\left(\bigcap_{i \in I} B_i\right) = \bigcap_{i \in I} f^{-1}(B_i).$$

- ▶ If  $B \subset Y$ , then  $f^{-1}(B^c) = f^{-1}(B)^c$ .

# Continuous functions

- ▶ Having the data of open sets turns out to be sufficient to define continuous functions.
- ▶ If  $f : X \rightarrow Y$  is a function between two sets, and  $A \subset Y$  is a subset, then we define the *preimage* of  $A$  to be

$$f^{-1}(A) = \{x \in X \mid f(x) \in A\}.$$

- ▶ **Proposition 2.10.** The preimage behaves nicely with respect to various operations of sets.

- ▶ If  $f : X \rightarrow Y$  and  $\{B_i\}_{i \in I}$  is a family of subsets of  $Y$ , then

$$f^{-1}\left(\bigcup_{i \in I} B_i\right) = \bigcup_{i \in I} f^{-1}(B_i), \quad f^{-1}\left(\bigcap_{i \in I} B_i\right) = \bigcap_{i \in I} f^{-1}(B_i).$$

- ▶ If  $B \subset Y$ , then  $f^{-1}(B^c) = f^{-1}(B)^c$ .
- ▶ If  $g : Y \rightarrow Z$  is another map and  $C \subset Z$ , then

$$(g \circ f)^{-1}(C) = f^{-1}(g^{-1}(C)).$$

- **Definition.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces. A function  $f : X \rightarrow Y$  is called *continuous* if  $f^{-1}(U) \in \mathcal{T}_X$  for all  $U \in \mathcal{T}_Y$ , or in words, if the preimages of open sets are open.

- ▶ **Definition.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces. A function  $f : X \rightarrow Y$  is called *continuous* if  $f^{-1}(U) \in \mathcal{T}_X$  for all  $U \in \mathcal{T}_Y$ , or in words, if the preimages of open sets are open.
- ▶ A function  $f : X \rightarrow Y$  is called *continuous at a point*  $x \in X$  if for every  $U \in \mathcal{T}_Y$  with  $f(x) \in U$  there is a  $V \in \mathcal{T}_X$  so that  $x \in V$  and  $f(V) \subset U$ .



# Continuous functions

- ▶ **Definition.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces. A function  $f : X \rightarrow Y$  is called *continuous* if  $f^{-1}(U) \in \mathcal{T}_X$  for all  $U \in \mathcal{T}_Y$ , or in words, if the preimages of open sets are open.
- ▶ A function  $f : X \rightarrow Y$  is called *continuous at a point*  $x \in X$  if for every  $U \in \mathcal{T}_Y$  with  $f(x) \in U$  there is a  $V \in \mathcal{T}_X$  so that  $x \in V$  and  $f(V) \subset U$ .
- ▶ **Remark.** The notion of “continuity” depends heavily on the topologies on the spaces under consideration.

# Continuous functions

- ▶ **Example.** Let  $X$  be a topological space. Then the identity map  $id : X \rightarrow X$  is continuous since  $id^{-1}(U) = U$  for every subset  $U \subset X$ .

# Continuous functions

- ▶ **Example.** Let  $X$  be a topological space. Then the identity map  $id : X \rightarrow X$  is continuous since  $id^{-1}(U) = U$  for every subset  $U \subset X$ .
- ▶ **Example.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces, and let  $y_0 \in Y$ . Then the constant map  $f : X \rightarrow Y$ ,  $f(x) = y_0$  for all  $x$ , is continuous.

# Continuous functions

- ▶ **Example.** Let  $X$  be a topological space. Then the identity map  $id : X \rightarrow X$  is continuous since  $id^{-1}(U) = U$  for every subset  $U \subset X$ .
- ▶ **Example.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces, and let  $y_0 \in Y$ . Then the constant map  $f : X \rightarrow Y$ ,  $f(x) = y_0$  for all  $x$ , is continuous.
- ▶ **Example.** Let  $X$  have the discrete topology, and let  $Y$  be any topological space. Then any map  $f : X \rightarrow Y$  is continuous, since  $f^{-1}(U) \in \mathcal{P}(X)$  for all  $U \subset Y$ .

# Continuous functions

- ▶ **Example.** Let  $X$  be a topological space. Then the identity map  $id : X \rightarrow X$  is continuous since  $id^{-1}(U) = U$  for every subset  $U \subset X$ .
- ▶ **Example.** Let  $(X, \mathcal{T}_X)$  and  $(Y, \mathcal{T}_Y)$  be topological spaces, and let  $y_0 \in Y$ . Then the constant map  $f : X \rightarrow Y$ ,  $f(x) = y_0$  for all  $x$ , is continuous.
- ▶ **Example.** Let  $X$  have the discrete topology, and let  $Y$  be any topological space. Then any map  $f : X \rightarrow Y$  is continuous, since  $f^{-1}(U) \in \mathcal{P}(X)$  for all  $U \subset Y$ .
- ▶ **Example.** Let  $X$  be any topological space, and let  $Y$  have the trivial topology. Then any map  $f : X \rightarrow Y$  is continuous, since  $f^{-1}(\emptyset) = \emptyset$  and  $f^{-1}(Y) = X$  which are both open.

- ▶ **Theorem 2.11.** The following properties hold:

# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .

# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .
  - ▶ (ii) A function  $f : X \rightarrow Y$  is continuous if and only if the preimage of any closed set is closed.



# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .
  - ▶ (ii) A function  $f : X \rightarrow Y$  is continuous if and only if the preimage of any closed set is closed.
  - ▶ (iii) A function  $f : X \rightarrow Y$  is continuous if and only if it is continuous at  $x$  for all  $x \in X$ .

# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .
  - ▶ (ii) A function  $f : X \rightarrow Y$  is continuous if and only if the preimage of any closed set is closed.
  - ▶ (iii) A function  $f : X \rightarrow Y$  is continuous if and only if it is continuous at  $x$  for all  $x \in X$ .
- ▶ *Proof.* (i) follows from  $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$ .  
(ii) follows from  $f^{-1}(C)^c = f^{-1}(C^c)$ . We now show (iii).

# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .
  - ▶ (ii) A function  $f : X \rightarrow Y$  is continuous if and only if the preimage of any closed set is closed.
  - ▶ (iii) A function  $f : X \rightarrow Y$  is continuous if and only if it is continuous at  $x$  for all  $x \in X$ .
- ▶ *Proof.* (i) follows from  $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$ .  
(ii) follows from  $f^{-1}(C)^c = f^{-1}(C^c)$ . We now show (iii).
  - ▶ Suppose  $f$  is continuous. Let  $x \in X$  and let  $U \in \mathcal{T}_Y$  with  $f(x) \in U$ . Then  $V = f^{-1}(U)$  is open in  $X$  and  $f(V) = U$ , so  $f$  is continuous at  $x$ .

# Continuous functions

- ▶ **Theorem 2.11.** The following properties hold:
  - ▶ (i) If  $f : X \rightarrow Y$  and  $g : Y \rightarrow Z$  are continuous, then so is  $g \circ f : X \rightarrow Z$ .
  - ▶ (ii) A function  $f : X \rightarrow Y$  is continuous if and only if the preimage of any closed set is closed.
  - ▶ (iii) A function  $f : X \rightarrow Y$  is continuous if and only if it is continuous at  $x$  for all  $x \in X$ .
- ▶ *Proof.* (i) follows from  $(g \circ f)^{-1}(U) = f^{-1}(g^{-1}(U))$ .  
(ii) follows from  $f^{-1}(C)^c = f^{-1}(C^c)$ . We now show (iii).
  - ▶ Suppose  $f$  is continuous. Let  $x \in X$  and let  $U \in \mathcal{T}_Y$  with  $f(x) \in U$ . Then  $V = f^{-1}(U)$  is open in  $X$  and  $f(V) = U$ , so  $f$  is continuous at  $x$ .
  - ▶ Suppose  $f$  is continuous at  $x$  for all  $x \in X$ . Let  $U \in \mathcal{T}_Y$ . We need to show that  $f^{-1}(U) \in \mathcal{T}_X$ . For each  $x \in f^{-1}(U)$ , there exists  $V_x \in \mathcal{T}_X$  so that  $x \in V_x$  and  $f(V_x) \subset U$ . Now  $f^{-1}(U) = \bigcup_{x \in f^{-1}(U)} V_x$  and thus open since each  $V_x$  is.

# Continuous functions

- ▶ Recall from analysis that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called *continuous at a point*  $x \in \mathbb{R}^n$  if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that  $\|x - y\| < \delta \Rightarrow \|f(x) - f(y)\| < \epsilon$ .  
A function is called *continuous* if it is cont. at every point.

# Continuous functions

- ▶ Recall from analysis that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called *continuous at a point*  $x \in \mathbb{R}^n$  if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that  $\|x - y\| < \delta \Rightarrow \|f(x) - f(y)\| < \epsilon$ .

A function is called *continuous* if it is cont. at every point.

- ▶ **Theorem 2.12.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces with their induced metric topologies. Then a function  $f : X \rightarrow Y$  is continuous if and only if

$$\forall x \in X, \forall \epsilon > 0, \exists \delta > 0 : d_X(x, y) < \delta \Rightarrow d_Y(f(x), f(y)) < \epsilon.$$

# Continuous functions

- ▶ Recall from analysis that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called *continuous at a point*  $x \in \mathbb{R}^n$  if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that  $\|x - y\| < \delta \Rightarrow \|f(x) - f(y)\| < \epsilon$ .

A function is called *continuous* if it is cont. at every point.

- ▶ **Theorem 2.12.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces with their induced metric topologies. Then a function  $f : X \rightarrow Y$  is continuous if and only if

$$\forall x \in X, \forall \epsilon > 0, \exists \delta > 0 : d_X(x, y) < \delta \Rightarrow d_Y(f(x), f(y)) < \epsilon.$$

- ▶ *Proof.* Theorem 2.12 follows from Theorem 2.11(iii) and Lemma 2.13.

# Continuous functions

- ▶ Recall from analysis that a function  $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$  is called *continuous at a point*  $x \in \mathbb{R}^n$  if for every  $\epsilon > 0$  there exists  $\delta > 0$  such that  $\|x - y\| < \delta \Rightarrow \|f(x) - f(y)\| < \epsilon$ .

A function is called *continuous* if it is cont. at every point.

- ▶ **Theorem 2.12.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces with their induced metric topologies. Then a function  $f : X \rightarrow Y$  is continuous if and only if

$$\forall x \in X, \forall \epsilon > 0, \exists \delta > 0 : d_X(x, y) < \delta \Rightarrow d_Y(f(x), f(y)) < \epsilon.$$

- ▶ *Proof.* Theorem 2.12 follows from Theorem 2.11(iii) and Lemma 2.13.
- ▶ **Lemma 2.13.** Let  $(X, d_X)$  and  $(Y, d_Y)$  be metric spaces with the metric topologies. Then a function  $f : X \rightarrow Y$  is continuous at a point  $x \in X$  if and only if

$$\forall \epsilon > 0, \exists \delta > 0 : f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon). \quad (1)$$



# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset f(V) \subset B_{d_Y}(f(x), \epsilon)$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
- ▶ Suppose (1) holds for  $f$ . Let  $U$  be an open set in  $Y$  containing  $f(x)$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
- ▶ Suppose (1) holds for  $f$ . Let  $U$  be an open set in  $Y$  containing  $f(x)$ .
  - ▶ By Proposition 2.9, the openness of  $U$  implies that there exists  $\epsilon > 0$  so that  $B_{d_Y}(f(x), \epsilon) \subset U$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
- ▶ Suppose (1) holds for  $f$ . Let  $U$  be an open set in  $Y$  containing  $f(x)$ .
  - ▶ By Proposition 2.9, the openness of  $U$  implies that there exists  $\epsilon > 0$  so that  $B_{d_Y}(f(x), \epsilon) \subset U$ .
  - ▶ By (1) we then get a  $\delta > 0$  with  $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$ .

# Continuous functions: Proof of Lemma 2.13

- ▶ Suppose  $f$  is continuous at  $x \in X$ . Let  $\epsilon > 0$ .
  - ▶ There is an open set  $V$  in  $X$  such that  $x \in V$  and  $f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ By Proposition 2.9, the openness of  $V$  implies that there is a  $\delta > 0$  so that  $B_{d_X}(x, \delta) \subset V$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset f(V) \subset B_{d_Y}(f(x), \epsilon)$ .
- ▶ Suppose (1) holds for  $f$ . Let  $U$  be an open set in  $Y$  containing  $f(x)$ .
  - ▶ By Proposition 2.9, the openness of  $U$  implies that there exists  $\epsilon > 0$  so that  $B_{d_Y}(f(x), \epsilon) \subset U$ .
  - ▶ By (1) we then get a  $\delta > 0$  with  $f(B_{d_X}(x, \delta)) \subset B_{d_Y}(f(x), \epsilon)$ .
  - ▶ Hence  $f(B_{d_X}(x, \delta)) \subset U$ . Since  $B_{d_X}(x, \delta)$  is open in  $X$  and contains  $x$  we are done.