

1. Use Gauss's lemma to calculate $(\frac{5}{23})$.

Consider the multiples of 5 up to $\frac{23-1}{2} = 11$ times 5. These are the integers 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55. The least residues mod 23 of these integers are 5, 10, 15, 20, 2, 7, 12, 17, 22, 4, 9. Of these least residues, exactly 5 exceed $\frac{23}{2} = 11.5$. So $(\frac{5}{23}) = (-1)^5 = -1$.

2. (a) Prove that $4^n \equiv 4 \pmod{12}$ for all integers $n \geq 1$.

- (b) If p is a prime of the form $4^n + 1$, then is 3 a quadratic residue or nonresidue mod p ?

(a) $4^2 \equiv 16 \equiv 4 \pmod{12}$. $4^3 \equiv 4 \cdot 4^2 \equiv 4 \cdot 4 \equiv 4 \pmod{12}$, $4^4 \equiv 4 \cdot 4^3 \equiv 4 \cdot 4 \equiv 4 \pmod{12}$, and so on. The best way to write this up is induction, but the above is acceptable too.

(b) Let p be an odd prime. We know (from class) that 3 is a quadratic residue mod p if and only if p is of the form $12k+1$ or $12k+11$. If p is of the form 4^n+1 , then by part (a), $p \equiv 4+1 \equiv 5 \pmod{12}$. In other words, it is of the form $12k+5$. Thus 3 is a quadratic non-residue mod p .

3. Let p be an odd prime. Use Quadratic Reciprocity to calculate $(\frac{127}{311})$. Note that 127 and 311 are both primes.

$$\begin{aligned} \left(\frac{127}{311}\right) &= \left(\frac{311}{127}\right)(-1)^{\frac{311-1}{2} \frac{127-1}{2}} = -\left(\frac{311}{127}\right) = -\left(\frac{57}{127}\right) = -\left(\frac{3}{127}\right)\left(\frac{19}{127}\right) = -\left(\frac{127}{3}\right)\left(\frac{127}{19}\right)(-1)^{\frac{127-1}{2} \frac{3-1}{2}}(-1)^{\frac{127-1}{2} \frac{19-1}{2}} \\ &= -\left(\frac{127}{3}\right)\left(\frac{127}{19}\right) = -\left(\frac{1}{3}\right)\left(\frac{13}{19}\right) = -\left(\frac{13}{19}\right) = -\left(\frac{19}{13}\right)(-1)^{\frac{13-1}{2} \frac{19-1}{2}} = -\left(\frac{6}{13}\right) = -\left(\frac{3}{13}\right)\left(\frac{2}{13}\right) = \left(\frac{3}{13}\right), \text{ where} \\ &\text{we used } \left(\frac{2}{13}\right) = -1 \text{ since } 13 = 8k+5. \text{ Continuing with the calculation, } \left(\frac{3}{13}\right) = \left(\frac{13}{3}\right)(-1)^{\frac{13-1}{2} \frac{3-1}{2}} = \\ &\left(\frac{13}{3}\right) = \left(\frac{1}{3}\right) = 1. \end{aligned}$$

Final answer, $(\frac{127}{311}) = 1$.

4. Use Quadratic Reciprocity to determine when 5 is a quadratic residue mod p and when it is not.

$(\frac{5}{p}) = (\frac{p}{5})(-1)^{\frac{5-1}{2} \frac{p-1}{2}} = (\frac{p}{5})$. The quadratic residues of 5 are 1 and 4. Thus $(\frac{p}{5}) = 1$ if and only if $p \equiv 1 \pmod{5}$ or $p \equiv 4 \pmod{5}$. In summary,

$$\left(\frac{5}{p}\right) = \begin{cases} 1 & \text{if and only if } p = 5k+1 \text{ or } p = 5k+4 \\ -1 & \text{if and only if } p = 5k+2 \text{ or } p = 5k+3. \end{cases}$$