

## 5. Connectedness

Math 4341 (Topology)

# Homeomorphisms

# Homeomorphisms

- ▶ **Definition.** A bijection  $f : X \rightarrow Y$  between two topological spaces is called a *homeomorphism* if both  $f$  and its inverse  $f^{-1}$  are continuous. In this case, we say that  $X$  and  $Y$  are *homeomorphic* and we write  $X \simeq Y$ .

# Homeomorphisms

- ▶ **Definition.** A bijection  $f : X \rightarrow Y$  between two topological spaces is called a *homeomorphism* if both  $f$  and its inverse  $f^{-1}$  are continuous. In this case, we say that  $X$  and  $Y$  are *homeomorphic* and we write  $X \simeq Y$ .
- ▶ Note that  $\simeq$  satisfies the property of an equivalence relation.

# Homeomorphisms

- ▶ **Definition.** A bijection  $f : X \rightarrow Y$  between two topological spaces is called a *homeomorphism* if both  $f$  and its inverse  $f^{-1}$  are continuous. In this case, we say that  $X$  and  $Y$  are *homeomorphic* and we write  $X \simeq Y$ .
- ▶ Note that  $\simeq$  satisfies the property of an equivalence relation.
- ▶ **Example.** Let  $f : (-1, 1) \rightarrow \mathbb{R}$  be the bijective map

$$f(x) = \tan\left(\frac{\pi x}{2}\right)$$

whose inverse is  $f^{-1}(x) = \frac{2}{\pi} \arctan x$ . Then both  $f$  and  $f^{-1}$  are continuous so  $(-1, 1)$  and  $\mathbb{R}$  are homeomorphic.

- **Example.** Let  $B^n := B(0, 1)$  be the unit ball in  $\mathbb{R}^n$ . Then  $B^n \simeq \mathbb{R}^n$ . This is because the map  $f : B^n \rightarrow \mathbb{R}^n$  given by

$$f(x) = \frac{x}{1 - \|x\|}$$

is a continuous bijection with inverse  $f^{-1}(x) = \frac{x}{1 + \|x\|}$ .

# Homeomorphisms

- **Example.** Let  $B^n := B(0, 1)$  be the unit ball in  $\mathbb{R}^n$ . Then  $B^n \simeq \mathbb{R}^n$ . This is because the map  $f : B^n \rightarrow \mathbb{R}^n$  given by

$$f(x) = \frac{x}{1 - \|x\|}$$

is a continuous bijection with inverse  $f^{-1}(x) = \frac{x}{1 + \|x\|}$ .

- **Definition.** Let  $X$  and  $Y$  be topological spaces. A function  $f : X \rightarrow Y$  is called an *embedding* if  $f : X \rightarrow f(X)$  is a homeomorphism; where  $f(X) \subset Y$  has the subspace topology.

- ▶ **Example.** Let  $B^n := B(0, 1)$  be the unit ball in  $\mathbb{R}^n$ . Then  $B^n \simeq \mathbb{R}^n$ . This is because the map  $f : B^n \rightarrow \mathbb{R}^n$  given by

$$f(x) = \frac{x}{1 - \|x\|}$$

is a continuous bijection with inverse  $f^{-1}(x) = \frac{x}{1 + \|x\|}$ .

- ▶ **Definition.** Let  $X$  and  $Y$  be topological spaces. A function  $f : X \rightarrow Y$  is called an *embedding* if  $f : X \rightarrow f(X)$  is a homeomorphism; where  $f(X) \subset Y$  has the subspace topology.
- ▶ **Example.** If  $X$  is a top. space and  $Y \subset X$  a subspace, then the inclusion  $\iota : Y \rightarrow X$  given by  $\iota(x) = x$  is an embedding.



# The $n$ -dimensional sphere

- **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .
- ▶ *Proof.* Let  $x = (x_1, \dots, x_{n+1}) \in S^n \setminus \{p\}$  so that  $x_{n+1} \neq 1$ .

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .
- ▶ *Proof.* Let  $x = (x_1, \dots, x_{n+1}) \in S^n \setminus \{p\}$  so that  $x_{n+1} \neq 1$ .
  - ▶ Define the *stereographic projection* of  $x$  by

$$f(x) = f(x_1, \dots, x_n, x_{n+1}) = \frac{1}{1 - x_{n+1}}(x_1, \dots, x_n) \in \mathbb{R}^n.$$

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .
- ▶ *Proof.* Let  $x = (x_1, \dots, x_{n+1}) \in S^n \setminus \{p\}$  so that  $x_{n+1} \neq 1$ .
  - ▶ Define the *stereographic projection* of  $x$  by

$$f(x) = f(x_1, \dots, x_n, x_{n+1}) = \frac{1}{1 - x_{n+1}}(x_1, \dots, x_n) \in \mathbb{R}^n.$$

- ▶ Geometrically, if one draws a straight line through  $x$  and  $p$ , then its intersection with  $\mathbb{R}^n \times \{0\}$  is the point  $(f(x), 0)$ .

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .
- ▶ *Proof.* Let  $x = (x_1, \dots, x_{n+1}) \in S^n \setminus \{p\}$  so that  $x_{n+1} \neq 1$ .
  - ▶ Define the *stereographic projection* of  $x$  by

$$f(x) = f(x_1, \dots, x_n, x_{n+1}) = \frac{1}{1 - x_{n+1}}(x_1, \dots, x_n) \in \mathbb{R}^n.$$

- ▶ Geometrically, if one draws a straight line through  $x$  and  $p$ , then its intersection with  $\mathbb{R}^n \times \{0\}$  is the point  $(f(x), 0)$ .
- ▶  $f$  is continuous because each of its components are.

# The $n$ -dimensional sphere

- ▶ **Definition.** The  $n$ -sphere is the set

$$S^n = \{x \in \mathbb{R}^{n+1} : \|x\| = 1\} \subset \mathbb{R}^{n+1}$$

with the subspace topology from  $\mathbb{R}^{n+1}$ .

- ▶ **Proposition 5.1.** Let  $p = (0, 0, \dots, 0, 1) \in S^n$  be the “north pole”. Then  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ .
- ▶ *Proof.* Let  $x = (x_1, \dots, x_{n+1}) \in S^n \setminus \{p\}$  so that  $x_{n+1} \neq 1$ .
  - ▶ Define the *stereographic projection* of  $x$  by

$$f(x) = f(x_1, \dots, x_n, x_{n+1}) = \frac{1}{1 - x_{n+1}}(x_1, \dots, x_n) \in \mathbb{R}^n.$$

- ▶ Geometrically, if one draws a straight line through  $x$  and  $p$ , then its intersection with  $\mathbb{R}^n \times \{0\}$  is the point  $(f(x), 0)$ .
- ▶  $f$  is continuous because each of its components are.
- ▶  $f$  has a continuous inverse  $g : \mathbb{R}^n \rightarrow S^n \setminus \{p\}$  given by

$$g(y_1, \dots, y_n) = (t(y)y_1, \dots, t(y)y_n, 1 - t(y)),$$

where  $t(y) = 2/(1 + \|y\|^2)$ .

- **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.



- ▶ **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.
- ▶ **Example.** The subspace  $(0, 1) \cup (2, 3) \subset \mathbb{R}$  has a separation.

- ▶ **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.
- ▶ **Example.** The subspace  $(0, 1) \cup (2, 3) \subset \mathbb{R}$  has a separation.
- ▶ **Remark.** If  $X = U \cup V$  is a separation, then  $U = X \setminus V$  and  $V = X \setminus U$  are both open and closed.

- ▶ **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.
- ▶ **Example.** The subspace  $(0, 1) \cup (2, 3) \subset \mathbb{R}$  has a separation.
- ▶ **Remark.** If  $X = U \cup V$  is a separation, then  $U = X \setminus V$  and  $V = X \setminus U$  are both open and closed.
- ▶ **Lemma 5.2.** A topological space  $X$  is connected iff  $\emptyset$  and  $X$  are the only subsets of  $X$  that are both open and closed.

- ▶ **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.
- ▶ **Example.** The subspace  $(0, 1) \cup (2, 3) \subset \mathbb{R}$  has a separation.
- ▶ **Remark.** If  $X = U \cup V$  is a separation, then  $U = X \setminus V$  and  $V = X \setminus U$  are both open and closed.
- ▶ **Lemma 5.2.** A topological space  $X$  is connected iff  $\emptyset$  and  $X$  are the only subsets of  $X$  that are both open and closed.
- ▶ **Example.** The rational numbers  $\mathbb{Q} \subset \mathbb{R}$  are not connected.

- ▶ **Definition.** Let  $X$  be a top. space. A *separation* of  $X$  is a pair  $U, V$  of disjoint non-empty open subsets of  $X$  such that  $X = U \cup V$ . We say  $X$  is *connected* if it has no separation.
- ▶ **Example.** The subspace  $(0, 1) \cup (2, 3) \subset \mathbb{R}$  has a separation.
- ▶ **Remark.** If  $X = U \cup V$  is a separation, then  $U = X \setminus V$  and  $V = X \setminus U$  are both open and closed.
- ▶ **Lemma 5.2.** A topological space  $X$  is connected iff  $\emptyset$  and  $X$  are the only subsets of  $X$  that are both open and closed.
- ▶ **Example.** The rational numbers  $\mathbb{Q} \subset \mathbb{R}$  are not connected.
  - ▶ Choose any irrational number  $a \in \mathbb{R}$ . Then

$$\mathbb{Q} = ((-\infty, a) \cap \mathbb{Q}) \cup (\mathbb{Q} \cap (a, \infty)),$$

which is a separation.

- ▶ **Example.** If  $X$  has the discrete topology and consists of at least two points, then  $X = \{x\} \cup (X \setminus \{x\})$  is a separation of  $X$ , so  $X$  is not connected.

- ▶ **Example.** If  $X$  has the discrete topology and consists of at least two points, then  $X = \{x\} \cup (X \setminus \{x\})$  is a separation of  $X$ , so  $X$  is not connected.
- ▶ **Lemma 5.3.** Let  $X = U \cup V$  for disjoint open sets  $U$  and  $V$ , and let  $Y \subset X$ . If  $Y$  is connected, then  $Y \subset U$  or  $Y \subset V$ .

- ▶ **Example.** If  $X$  has the discrete topology and consists of at least two points, then  $X = \{x\} \cup (X \setminus \{x\})$  is a separation of  $X$ , so  $X$  is not connected.
- ▶ **Lemma 5.3.** Let  $X = U \cup V$  for disjoint open sets  $U$  and  $V$ , and let  $Y \subset X$ . If  $Y$  is connected, then  $Y \subset U$  or  $Y \subset V$ .
- ▶ **Proof.** We will show the contrapositive of the statement, so assume that  $Y \cap U \neq \emptyset$  and  $Y \cap V \neq \emptyset$ . Then

$$Y = Y \cap X = Y \cap (U \cup V) = (Y \cap U) \cup (Y \cap V)$$

is a separation of  $Y$  (since  $Y \cap U$  and  $Y \cap V$  are disjoint, non-empty and open in the subspace topology).



- ▶ **Theorem 5.4.** Let  $\{A_i\}_{i \in I}$  be a collection of connected subspaces of a topological space  $X$  with a common point  $x \in X$ ; i.e.  $x \in A_i$  for all  $i \in I$ . Then  $\bigcup_{i \in I} A_i$  is connected.

- ▶ **Theorem 5.4.** Let  $\{A_i\}_{i \in I}$  be a collection of connected subspaces of a topological space  $X$  with a common point  $x \in X$ ; i.e.  $x \in A_i$  for all  $i \in I$ . Then  $\bigcup_{i \in I} A_i$  is connected.
- ▶ **Proof.** Suppose  $\bigcup_{i \in I} A_i = U \cup V$  is a separation.

- ▶ **Theorem 5.4.** Let  $\{A_i\}_{i \in I}$  be a collection of connected subspaces of a topological space  $X$  with a common point  $x \in X$ ; i.e.  $x \in A_i$  for all  $i \in I$ . Then  $\bigcup_{i \in I} A_i$  is connected.
- ▶ **Proof.** Suppose  $\bigcup_{i \in I} A_i = U \cup V$  is a separation.
  - ▶ Assume without loss of generality that  $x \in U$ . By Lemma 5.3 we have for each  $i$  that either  $A_i \subset U$  or  $A_i \subset V$ .

- ▶ **Theorem 5.4.** Let  $\{A_i\}_{i \in I}$  be a collection of connected subspaces of a topological space  $X$  with a common point  $x \in X$ ; i.e.  $x \in A_i$  for all  $i \in I$ . Then  $\bigcup_{i \in I} A_i$  is connected.
- ▶ **Proof.** Suppose  $\bigcup_{i \in I} A_i = U \cup V$  is a separation.
  - ▶ Assume without loss of generality that  $x \in U$ . By Lemma 5.3 we have for each  $i$  that either  $A_i \subset U$  or  $A_i \subset V$ .
  - ▶ Since  $x \in A_i$  we must have  $A_i \subset U$  for all  $i \in I$ . This implies that  $\bigcup_{i \in I} A_i \subset U$ , so  $V$  must be empty.

- ▶ **Theorem 5.5.** Let  $A \subset X$  be connected. If a subset  $B \subset X$  satisfies  $A \subset B \subset \bar{A}$ , then  $B$  is also connected. In particular,  $\bar{A}$  is connected whenever  $A$  is.

- ▶ **Theorem 5.5.** Let  $A \subset X$  be connected. If a subset  $B \subset X$  satisfies  $A \subset B \subset \bar{A}$ , then  $B$  is also connected. In particular,  $\bar{A}$  is connected whenever  $A$  is.
- ▶ *Proof.* Suppose  $B = U \cup V$  is a separation.

- ▶ **Theorem 5.5.** Let  $A \subset X$  be connected. If a subset  $B \subset X$  satisfies  $A \subset B \subset \bar{A}$ , then  $B$  is also connected. In particular,  $\bar{A}$  is connected whenever  $A$  is.
- ▶ *Proof.* Suppose  $B = U \cup V$  is a separation.
  - ▶ By Lemma 5.3 we must have that  $A \subset U$  or  $A \subset V$ , so assume without loss of generality that  $A \subset U$ . Then  $B \subset \bar{A} \subset \bar{U}$  (where all closures are in  $X$ ).

- ▶ **Theorem 5.5.** Let  $A \subset X$  be connected. If a subset  $B \subset X$  satisfies  $A \subset B \subset \bar{A}$ , then  $B$  is also connected. In particular,  $\bar{A}$  is connected whenever  $A$  is.
- ▶ *Proof.* Suppose  $B = U \cup V$  is a separation.
  - ▶ By Lemma 5.3 we must have that  $A \subset U$  or  $A \subset V$ , so assume without loss of generality that  $A \subset U$ . Then  $B \subset \bar{A} \subset \bar{U}$  (where all closures are in  $X$ ).
  - ▶ By definition of the subspace topology, there are open sets  $U'$  and  $V'$  in  $X$  so that  $U = B \cap U'$ ,  $V = B \cap V'$ , and

$$U = B \cap U' = B \setminus V' \subset X \setminus V'.$$

The latter space is closed so  $\bar{U} \subset X \setminus V' \subset X \setminus V$ .



- ▶ **Theorem 5.5.** Let  $A \subset X$  be connected. If a subset  $B \subset X$  satisfies  $A \subset B \subset \bar{A}$ , then  $B$  is also connected. In particular,  $\bar{A}$  is connected whenever  $A$  is.
- ▶ *Proof.* Suppose  $B = U \cup V$  is a separation.
  - ▶ By Lemma 5.3 we must have that  $A \subset U$  or  $A \subset V$ , so assume without loss of generality that  $A \subset U$ . Then  $B \subset \bar{A} \subset \bar{U}$  (where all closures are in  $X$ ).
  - ▶ By definition of the subspace topology, there are open sets  $U'$  and  $V'$  in  $X$  so that  $U = B \cap U'$ ,  $V = B \cap V'$ , and

$$U = B \setminus V = B \setminus V' \subset X \setminus V'.$$

The latter space is closed so  $\bar{U} \subset X \setminus V' \subset X \setminus V$ .

- ▶ Hence  $B \subset \bar{U} \subset X \setminus V$  which means  $B \cap V = \emptyset$ , so  $V = \emptyset$ .

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.
- ▶ *Proof.* Suppose that  $f(X) = U \cup V$  is a separation.

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.
- ▶ *Proof.* Suppose that  $f(X) = U \cup V$  is a separation.
  - ▶ Then  $f^{-1}(U)$  and  $f^{-1}(V)$  are disjoint open subsets of  $X$  with  $X = f^{-1}(U) \cup f^{-1}(V)$ . Since  $X$  is connected, either  $f^{-1}(U)$  or  $f^{-1}(V)$  is empty.

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.
- ▶ *Proof.* Suppose that  $f(X) = U \cup V$  is a separation.
  - ▶ Then  $f^{-1}(U)$  and  $f^{-1}(V)$  are disjoint open subsets of  $X$  with  $X = f^{-1}(U) \cup f^{-1}(V)$ . Since  $X$  is connected, either  $f^{-1}(U)$  or  $f^{-1}(V)$  is empty.
  - ▶ Assume  $f^{-1}(U) = \emptyset$ . Since  $U \subset f(X)$  we must have  $U = \emptyset$ .

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.
- ▶ *Proof.* Suppose that  $f(X) = U \cup V$  is a separation.
  - ▶ Then  $f^{-1}(U)$  and  $f^{-1}(V)$  are disjoint open subsets of  $X$  with  $X = f^{-1}(U) \cup f^{-1}(V)$ . Since  $X$  is connected, either  $f^{-1}(U)$  or  $f^{-1}(V)$  is empty.
  - ▶ Assume  $f^{-1}(U) = \emptyset$ . Since  $U \subset f(X)$  we must have  $U = \emptyset$ .
- ▶ **Corollary 5.7.** [Intermediate value theorem] Let  $f : X \rightarrow \mathbb{R}$  be continuous and assume that  $X$  is connected. If there is an  $r \in \mathbb{R}$  and  $x, y \in X$  so that  $f(x) < r < f(y)$ , then there is a  $z \in X$  with  $f(z) = r$ .

- ▶ **Theorem 5.6.** Let  $f : X \rightarrow Y$  be a continuous map between top. spaces. If  $X$  is connected, then  $f(X)$  is also connected.
- ▶ *Proof.* Suppose that  $f(X) = U \cup V$  is a separation.
  - ▶ Then  $f^{-1}(U)$  and  $f^{-1}(V)$  are disjoint open subsets of  $X$  with  $X = f^{-1}(U) \cup f^{-1}(V)$ . Since  $X$  is connected, either  $f^{-1}(U)$  or  $f^{-1}(V)$  is empty.
  - ▶ Assume  $f^{-1}(U) = \emptyset$ . Since  $U \subset f(X)$  we must have  $U = \emptyset$ .
- ▶ **Corollary 5.7.** [Intermediate value theorem] Let  $f : X \rightarrow \mathbb{R}$  be continuous and assume that  $X$  is connected. If there is an  $r \in \mathbb{R}$  and  $x, y \in X$  so that  $f(x) < r < f(y)$ , then there is a  $z \in X$  with  $f(z) = r$ .
- ▶ *Proof.* By Theorem 5.6  $f(X) \subset \mathbb{R}$  is connected. This implies that  $r \in [f(x), f(y)] \subset f(X)$  (why ???).

- **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.



- ▶ **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.
- ▶ *Proof.* ( $\Rightarrow$ ) Use Theorem 5.6 and the fact that the projection  $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$  is continuous.

- ▶ **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.
- ▶ *Proof.* ( $\Rightarrow$ ) Use Theorem 5.6 and the fact that the projection  $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$  is continuous.
- ▶ ( $\Leftarrow$ ) We show that  $X \times Y$  is connected when  $X$  and  $Y$  are.

- ▶ **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.
- ▶ *Proof.* ( $\Rightarrow$ ) Use Theorem 5.6 and the fact that the projection  $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$  is continuous.
- ▶ ( $\Leftarrow$ ) We show that  $X \times Y$  is connected when  $X$  and  $Y$  are.
  - ▶ Fix  $x_0 \in X$  and let  $A_{x_0} = \{x_0\} \times Y$ . Then  $A_{x_0}$  is the image of the continuous map  $Y \rightarrow X \times Y$  given by  $y \mapsto (x_0, y)$ , so  $A_{x_0}$  is connected by Theorem 5.6.

- ▶ **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.
- ▶ *Proof.* ( $\Rightarrow$ ) Use Theorem 5.6 and the fact that the projection  $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$  is continuous.
- ▶ ( $\Leftarrow$ ) We show that  $X \times Y$  is connected when  $X$  and  $Y$  are.
  - ▶ Fix  $x_0 \in X$  and let  $A_{x_0} = \{x_0\} \times Y$ . Then  $A_{x_0}$  is the image of the continuous map  $Y \rightarrow X \times Y$  given by  $y \mapsto (x_0, y)$ , so  $A_{x_0}$  is connected by Theorem 5.6.
  - ▶ Similarly  $B_y = X \times \{y\}$  is connected for all  $y \in Y$ . By Theorem 5.4,  $A_{x_0} \cup B_y$  is connected for all  $y \in Y$  since  $(x_0, y)$  is contained in both  $A_{x_0}$  and  $B_y$ .

- ▶ **Theorem 5.8.** Let  $X_1, \dots, X_n$  be topological spaces. Then  $X_1 \times \dots \times X_n$  is connected iff every  $X_i$  is.
- ▶ *Proof.* ( $\Rightarrow$ ) Use Theorem 5.6 and the fact that the projection  $\pi_i : X_1 \times \dots \times X_n \rightarrow X_i$  is continuous.
- ▶ ( $\Leftarrow$ ) We show that  $X \times Y$  is connected when  $X$  and  $Y$  are.
  - ▶ Fix  $x_0 \in X$  and let  $A_{x_0} = \{x_0\} \times Y$ . Then  $A_{x_0}$  is the image of the continuous map  $Y \rightarrow X \times Y$  given by  $y \mapsto (x_0, y)$ , so  $A_{x_0}$  is connected by Theorem 5.6.
  - ▶ Similarly  $B_y = X \times \{y\}$  is connected for all  $y \in Y$ . By Theorem 5.4,  $A_{x_0} \cup B_y$  is connected for all  $y \in Y$  since  $(x_0, y)$  is contained in both  $A_{x_0}$  and  $B_y$ .
  - ▶ Now clearly,

$$X \times Y = \bigcup_{y \in Y} A_{x_0} \cup B_y,$$

and all the sets on the RHS have the common point  $(x_0, *)$ , where  $*$   $\in Y$ . Therefore,  $X \times Y$  is connected by Theorem 5.4.

- ▶ **Corollary 5.9.**  $\mathbb{R}^n$  is connected for all  $n \in \mathbb{N}$ .

- ▶ **Corollary 5.9.**  $\mathbb{R}^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ **Prop. 5.10.** The  $n$ -sphere  $S^n$  is connected for all  $n \in \mathbb{N}$ .

- ▶ **Corollary 5.9.**  $\mathbb{R}^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ **Prop. 5.10.** The  $n$ -sphere  $S^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ *Proof.* Recall from Proposition 5.1 that  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ , where  $p$  is the north pole. It follows that  $S^n \setminus \{p\}$  is connected. Since  $\overline{S^n \setminus \{p\}} = S^n$ , the result follows from Theorem 5.5.



- ▶ **Corollary 5.9.**  $\mathbb{R}^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ **Prop. 5.10.** The  $n$ -sphere  $S^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ *Proof.* Recall from Proposition 5.1 that  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ , where  $p$  is the north pole. It follows that  $S^n \setminus \{p\}$  is connected. Since  $\overline{S^n \setminus \{p\}} = S^n$ , the result follows from Theorem 5.5.
- ▶ **Prop. 5.11.** For any  $n \in \mathbb{N}$ , we have  $S^n \not\simeq \mathbb{R}$ .

- ▶ **Corollary 5.9.**  $\mathbb{R}^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ **Prop. 5.10.** The  $n$ -sphere  $S^n$  is connected for all  $n \in \mathbb{N}$ .
- ▶ *Proof.* Recall from Proposition 5.1 that  $S^n \setminus \{p\} \simeq \mathbb{R}^n$ , where  $p$  is the north pole. It follows that  $S^n \setminus \{p\}$  is connected. Since  $\overline{S^n \setminus \{p\}} = S^n$ , the result follows from Theorem 5.5.
- ▶ **Prop. 5.11.** For any  $n \in \mathbb{N}$ , we have  $S^n \not\simeq \mathbb{R}$ .
- ▶ *Proof.* Suppose  $f : S^n \rightarrow \mathbb{R}$  is a homeomorphism. Then  $S^n \setminus \{p\} \simeq \mathbb{R} \setminus \{f(p)\}$ . We obtain a contradiction, since  $S^n \setminus \{p\} \simeq \mathbb{R}^n$  is connected while  $\mathbb{R} \setminus \{f(p)\}$  is not connected.

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .
  - ▶ If for any pair  $x, y$  in  $X$  there is a path from  $x$  to  $y$ , we say that  $X$  is *path-connected*.

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .
  - ▶ If for any pair  $x, y$  in  $X$  there is a path from  $x$  to  $y$ , we say that  $X$  is *path-connected*.
- ▶ **Prop. 5.12.** A path-connected space is connected.

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .
  - ▶ If for any pair  $x, y$  in  $X$  there is a path from  $x$  to  $y$ , we say that  $X$  is *path-connected*.
- ▶ **Prop. 5.12.** A path-connected space is connected.
- ▶ *Proof.* Suppose  $X = U \cup V$  be a separation.

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .
  - ▶ If for any pair  $x, y$  in  $X$  there is a path from  $x$  to  $y$ , we say that  $X$  is *path-connected*.
- ▶ **Prop. 5.12.** A path-connected space is connected.
- ▶ *Proof.* Suppose  $X = U \cup V$  be a separation.
  - ▶ Let  $\gamma : [0, 1] \rightarrow X$  be any path. By Theorem 5.6 we see that  $\gamma([0, 1])$  is connected and by Lemma 5.3,  $\gamma([0, 1])$  is contained entirely in either  $U$  or in  $V$ .

# Paths and path-connectedness

- ▶ **Definition.** Given two points  $x$  and  $y$  in a topological space  $X$ , a *path* from  $x$  to  $y$  is a continuous map  $\gamma : [0, 1] \rightarrow X$  such that  $\gamma(0) = x$  and  $\gamma(1) = y$ .
  - ▶ If for any pair  $x, y$  in  $X$  there is a path from  $x$  to  $y$ , we say that  $X$  is *path-connected*.
- ▶ **Prop. 5.12.** A path-connected space is connected.
- ▶ *Proof.* Suppose  $X = U \cup V$  be a separation.
  - ▶ Let  $\gamma : [0, 1] \rightarrow X$  be any path. By Theorem 5.6 we see that  $\gamma([0, 1])$  is connected and by Lemma 5.3,  $\gamma([0, 1])$  is contained entirely in either  $U$  or in  $V$ .
  - ▶ This means that it is not possible to find paths from points in  $U$  to points in  $V$ . Hence  $X$  is not path-connected.



- **Example.** For  $x, y \in \mathbb{R}^n$ , the path  $\gamma : [0, 1] \rightarrow \mathbb{R}^n$  defined by

$$\gamma(t) = (1 - t)x + ty$$

is a path from  $x$  to  $y$ , so  $\mathbb{R}^n$  is path-connected.

# Paths and path-connectedness

- ▶ **Example.** For  $x, y \in \mathbb{R}^n$ , the path  $\gamma : [0, 1] \rightarrow \mathbb{R}^n$  defined by

$$\gamma(t) = (1 - t)x + ty$$

is a path from  $x$  to  $y$ , so  $\mathbb{R}^n$  is path-connected.

- ▶ **Example.** A subset  $A$  of  $\mathbb{R}^n$  is called *convex* if for any  $x, y \in A$ , the image of the straight line  $\gamma(t) = (1 - t)x + ty$  belongs to  $A$ . As in the previous example, it follows that convex subsets are path-connected and thus also connected.

# Paths and path-connectedness

- ▶ **Example.** For  $x, y \in \mathbb{R}^n$ , the path  $\gamma : [0, 1] \rightarrow \mathbb{R}^n$  defined by

$$\gamma(t) = (1 - t)x + ty$$

is a path from  $x$  to  $y$ , so  $\mathbb{R}^n$  is path-connected.

- ▶ **Example.** A subset  $A$  of  $\mathbb{R}^n$  is called *convex* if for any  $x, y \in A$ , the image of the straight line  $\gamma(t) = (1 - t)x + ty$  belongs to  $A$ . As in the previous example, it follows that convex subsets are path-connected and thus also connected.
  - ▶ Some examples of convex subsets are the upper half-plane

$$\{(x_1, \dots, x_n) \in \mathbb{R}^n \mid x_n > 0\}$$


and any ball  $B(x, r)$ ,  $x \in \mathbb{R}^n$ ,  $r > 0$ .

# Paths and path-connectedness

- ▶ A connected space does not need to be path-connected. A counter-example is the so-called topologist's sine curve

$$S = \{(x, y) \in \mathbb{R}^2 \mid y = \sin(1/x), x > 0\} \cup \{(0, y) \mid -1 \leq y \leq 1\},$$

that is, the closure of the graph of  $x \mapsto \sin(1/x)$  for  $x > 0$ .  
For details, see Section 24 of Munkres' textbook.



# Connected components

- ▶ **Proposition 5.13.** Let  $X$  be a topological space. Define a relation  $\sim$  on  $X$  by declaring that  $x \sim y$  if and only if there is a connected set  $A \subset X$  such that  $x, y \in A$ . Then  $\sim$  is an equivalence relation. The equivalence classes of  $\sim$  are called the *connected components* of  $X$ .

# Connected components

- ▶ **Proposition 5.13.** Let  $X$  be a topological space. Define a relation  $\sim$  on  $X$  by declaring that  $x \sim y$  if and only if there is a connected set  $A \subset X$  such that  $x, y \in A$ . Then  $\sim$  is an equivalence relation. The equivalence classes of  $\sim$  are called the *connected components* of  $X$ .
- ▶ *Proof.* We see that  $x \sim x$  for every  $x$ , since  $\{x\}$  is connected.

- ▶ **Proposition 5.13.** Let  $X$  be a topological space. Define a relation  $\sim$  on  $X$  by declaring that  $x \sim y$  if and only if there is a connected set  $A \subset X$  such that  $x, y \in A$ . Then  $\sim$  is an equivalence relation. The equivalence classes of  $\sim$  are called the *connected components* of  $X$ .
- ▶ *Proof.* We see that  $x \sim x$  for every  $x$ , since  $\{x\}$  is connected.
  - ▶ If  $x \sim y$  there exists a connected set  $A$  with  $x, y \in A$ . Then clearly  $y, x \in A$  so  $y \sim x$ .

- ▶ **Proposition 5.13.** Let  $X$  be a topological space. Define a relation  $\sim$  on  $X$  by declaring that  $x \sim y$  if and only if there is a connected set  $A \subset X$  such that  $x, y \in A$ . Then  $\sim$  is an equivalence relation. The equivalence classes of  $\sim$  are called the *connected components* of  $X$ .
- ▶ *Proof.* We see that  $x \sim x$  for every  $x$ , since  $\{x\}$  is connected.
  - ▶ If  $x \sim y$  there exists a connected set  $A$  with  $x, y \in A$ . Then clearly  $y, x \in A$  so  $y \sim x$ .
  - ▶ If  $x \sim y$  and  $y \sim z$  we get connected sets  $A$  and  $B$  such that  $x, y \in A$  and  $y, z \in B$ . Let  $C = A \cup B$ . Then  $x, z \in C$ , and  $C$  is connected by Theorem 5.4, so  $x \sim z$ .



- **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then

# Connected components

- ▶ **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
  - ▶ (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,

# Connected components

- **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
- (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,

# Connected components

- **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
- (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,
  - (iii)  $C_i \subset X$  is connected for each  $i \in I$ , and

# Connected components

- **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
- (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,
  - (iii)  $C_i \subset X$  is connected for each  $i \in I$ , and
  - (iv)  $C_i$  is closed for all  $i \in I$ .

# Connected components

- ▶ **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
  - ▶ (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - ▶ (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,
  - ▶ (iii)  $C_i \subset X$  is connected for each  $i \in I$ , and
  - ▶ (iv)  $C_i$  is closed for all  $i \in I$ .
- ▶ *Proof.* (ii) Let  $Y \subset X$  be connected, and let  $x \in Y$ . Since  $y \sim x$  for all  $y \in Y$ , so  $Y \subset [x]$ .

# Connected components

- ▶ **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
  - ▶ (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - ▶ (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,
  - ▶ (iii)  $C_i \subset X$  is connected for each  $i \in I$ , and
  - ▶ (iv)  $C_i$  is closed for all  $i \in I$ .
- ▶ *Proof.* (ii) Let  $Y \subset X$  be connected, and let  $x \in Y$ . Since  $y \sim x$  for all  $y \in Y$ , so  $Y \subset [x]$ .
  - ▶ (iii) Fix  $x_0 \in C_i$ . For every  $y \in C_i$  there a connected subset  $A_y$  so that  $x_0, y \in A_y$ . Then  $A_y \subset C_i$  by (ii). Since  $C_i = \bigcup_{y \in C_i} A_y$  and all of the  $A_y$  contain  $x_0$ ,  $C_i$  is connected by Theorem 5.4.

# Connected components

- ▶ **Proposition 5.14.** Let  $\{C_i\}_{i \in I}$  be the set of connected components of a topological space  $X$ . Then
  - ▶ (i)  $X = \bigcup_{i \in I} C_i$  and the  $C_i$  are pairwise disjoint,
  - ▶ (ii) if  $Y \subset X$  is connected, then  $Y \subset C_i$  for some  $i \in I$ ,
  - ▶ (iii)  $C_i \subset X$  is connected for each  $i \in I$ , and
  - ▶ (iv)  $C_i$  is closed for all  $i \in I$ .
- ▶ *Proof.* (ii) Let  $Y \subset X$  be connected, and let  $x \in Y$ . Since  $y \sim x$  for all  $y \in Y$ , so  $Y \subset [x]$ .
  - ▶ (iii) Fix  $x_0 \in C_i$ . For every  $y \in C_i$  there a connected subset  $A_y$  so that  $x_0, y \in A_y$ . Then  $A_y \subset C_i$  by (ii). Since  $C_i = \bigcup_{y \in C_i} A_y$  and all of the  $A_y$  contain  $x_0$ ,  $C_i$  is connected by Theorem 5.4.
  - ▶ (iv) We will show that  $C_i = \overline{C_i}$ . Write  $C_i = [x]$  for any  $x \in C_i$ . Let  $y \in \overline{C_i}$ . Then  $\overline{C_i}$  is a subset containing both  $x$  and  $y$ , and  $\overline{C_i}$  is connected by Theorem 5.5, so  $y \in [x] = C_i$ .



# Connected components

- ▶ **Remark.** It follows from Proposition 5.14 that connected components are open if there are only finitely many of them.

# Connected components

- ▶ **Remark.** It follows from Proposition 5.14 that connected components are open if there are only finitely many of them.
- ▶ **Example.** We claim that the connected components of  $\mathbb{Q}$  are the singleton sets  $\{x\}$ . Indeed, let  $X$  be any subset of  $\mathbb{Q}$  containing at least two points, and suppose that  $x, y \in X$ ,  $x \neq y$ . There there is an irrational number  $r$ ,  $x < r < y$ , and

$$X = (X \cap (-\infty, r)) \cup (X \cap (r, \infty))$$

is a separation of  $X$ . Hence  $X$  is not connected.

# Connected components

- ▶ **Remark.** It follows from Proposition 5.14 that connected components are open if there are only finitely many of them.
- ▶ **Example.** We claim that the connected components of  $\mathbb{Q}$  are the singleton sets  $\{x\}$ . Indeed, let  $X$  be any subset of  $\mathbb{Q}$  containing at least two points, and suppose that  $x, y \in X$ ,  $x \neq y$ . There there is an irrational number  $r$ ,  $x < r < y$ , and

$$X = (X \cap (-\infty, r)) \cup (X \cap (r, \infty))$$

is a separation of  $X$ . Hence  $X$  is not connected.

- ▶ Since the topology on  $\mathbb{Q}$  is not the discrete one, the connected component  $\{x\}$  is not open for any  $x$ .

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.

# $\mathbb{R}$ is connected

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.
- ▶ *Proof:* Assume that  $U \cup V$  is a separation of  $\mathbb{R}$ . Let  $a, b \in \mathbb{R}$  with  $a \in U$  and  $b \in V$ , and without loss of generality we can assume that  $a < b$ .

# $\mathbb{R}$ is connected

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.
- ▶ *Proof:* Assume that  $U \cup V$  is a separation of  $\mathbb{R}$ . Let  $a, b \in \mathbb{R}$  with  $a \in U$  and  $b \in V$ , and without loss of generality we can assume that  $a < b$ .
- ▶ Let  $S = \{x \in \mathbb{R} \mid [a, x] \subset U\}$ . This is nonempty since  $a \in S$ . Also  $S$  is bounded above by  $b$  (otherwise  $b \in [a, x] \subset U$ ).

# $\mathbb{R}$ is connected

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.
- ▶ *Proof:* Assume that  $U \cup V$  is a separation of  $\mathbb{R}$ . Let  $a, b \in \mathbb{R}$  with  $a \in U$  and  $b \in V$ , and without loss of generality we can assume that  $a < b$ .
- ▶ Let  $S = \{x \in \mathbb{R} \mid [a, x] \subset U\}$ . This is nonempty since  $a \in S$ . Also  $S$  is bounded above by  $b$  (otherwise  $b \in [a, x] \subset U$ ).
- ▶ Let  $c = \sup S$ . If  $c \in V$ , then there exists an open interval  $(c - \varepsilon', c + \varepsilon') \subset V$ . Since  $c = \sup S$ , there exists  $y \in (c - \varepsilon', c) \cap S$ . Then  $y \in S$  implies that  $[a, y] \subset U$ . Both  $U$  and  $V$  contain  $y$ , a contradiction. Hence  $c \in U$ .

# $\mathbb{R}$ is connected

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.
- ▶ *Proof:* Assume that  $U \cup V$  is a separation of  $\mathbb{R}$ . Let  $a, b \in \mathbb{R}$  with  $a \in U$  and  $b \in V$ , and without loss of generality we can assume that  $a < b$ .
- ▶ Let  $S = \{x \in \mathbb{R} \mid [a, x] \subset U\}$ . This is nonempty since  $a \in S$ . Also  $S$  is bounded above by  $b$  (otherwise  $b \in [a, x] \subset U$ ).
- ▶ Let  $c = \sup S$ . If  $c \in V$ , then there exists an open interval  $(c - \varepsilon', c + \varepsilon') \subset V$ . Since  $c = \sup S$ , there exists  $y \in (c - \varepsilon', c) \cap S$ . Then  $y \in S$  implies that  $[a, y] \subset U$ . Both  $U$  and  $V$  contain  $y$ , a contradiction. Hence  $c \in U$ .
- ▶ Since  $U$  is open, there exists an open interval  $(c - \varepsilon, c + \varepsilon) \subset U$ . Since  $c = \sup S$ , there exists  $y \in (c - \varepsilon, c) \cap S$ . Then  $y \in S$  implies that  $[a, y] \subset U$ .



# $\mathbb{R}$ is connected

- ▶ **Theorem 5.15.**  $\mathbb{R}$  is connected.
- ▶ *Proof:* Assume that  $U \cup V$  is a separation of  $\mathbb{R}$ . Let  $a, b \in \mathbb{R}$  with  $a \in U$  and  $b \in V$ , and without loss of generality we can assume that  $a < b$ .
- ▶ Let  $S = \{x \in \mathbb{R} \mid [a, x] \subset U\}$ . This is nonempty since  $a \in S$ . Also  $S$  is bounded above by  $b$  (otherwise  $b \in [a, x] \subset U$ ).
- ▶ Let  $c = \sup S$ . If  $c \in V$ , then there exists an open interval  $(c - \varepsilon', c + \varepsilon') \subset V$ . Since  $c = \sup S$ , there exists  $y \in (c - \varepsilon', c) \cap S$ . Then  $y \in S$  implies that  $[a, y] \subset U$ . Both  $U$  and  $V$  contain  $y$ , a contradiction. Hence  $c \in U$ .
- ▶ Since  $U$  is open, there exists an open interval  $(c - \varepsilon, c + \varepsilon) \subset U$ . Since  $c = \sup S$ , there exists  $y \in (c - \varepsilon, c) \cap S$ . Then  $y \in S$  implies that  $[a, y] \subset U$ .
- ▶ Note that  $[y, c + \varepsilon/2] \subset (c - \varepsilon, c + \varepsilon) \subset U$ . Hence  $[a, c + \varepsilon/2] = [a, y] \cup [y, c + \varepsilon/2] \subset U$ . Then  $c + \varepsilon/2 \in S$ , which contradicts  $c = \sup S$ .