

§3. Constructing Topologies

Math 4341 (Topology)

Subspace topology

Subspace topology

- **Definition.** Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$ be any subset of X . Then the collection

$$\mathcal{T}_Y = \{Y \cap U \mid U \in \mathcal{T}\}$$

is called *the subspace topology*.

Subspace topology

- **Definition.** Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$ be any subset of X . Then the collection

$$\mathcal{T}_Y = \{Y \cap U \mid U \in \mathcal{T}\}$$

is called *the subspace topology*.

- **Lemma 3.1.** The collection \mathcal{T}_Y defines a topology on Y .

Subspace topology

- ▶ **Definition.** Let (X, \mathcal{T}) be a topological space, and let $Y \subset X$ be any subset of X . Then the collection

$$\mathcal{T}_Y = \{Y \cap U \mid U \in \mathcal{T}\}$$

is called *the subspace topology*.

- ▶ **Lemma 3.1.** The collection \mathcal{T}_Y defines a topology on Y .
- ▶ *Proof.* (T1) is obvious. (T2) and (T3) follow from De Morgan's Laws:

$$\bigcup_{i \in I} (Y \cap U_i) = Y \cap \bigcup_{i \in I} U_i, \quad \bigcap_{i \in I} (Y \cap U_i) = Y \cap \bigcap_{i \in I} U_i.$$

Subspace topology

- ▶ Equipping Y with the subspace topology, we will call Y a *subspace* of X . If a set U belongs to \mathcal{T}_Y we will often say that U is *open in Y* .

Subspace topology

- ▶ Equipping Y with the subspace topology, we will call Y a *subspace* of X . If a set U belongs to \mathcal{T}_Y we will often say that U is *open in Y* .
- ▶ **Remark.** A subspace might have open sets that are not open in the full topological space. For instance, let $X = \mathbb{R}$ and $Y = [0, \infty)$. Then the half-open interval $[0, 1)$ is open in Y since $[0, 1) = Y \cap (-1, 1)$, but $[0, 1)$ is not open in X .

Subspace topology

- ▶ Equipping Y with the subspace topology, we will call Y a *subspace* of X . If a set U belongs to \mathcal{T}_Y we will often say that U is *open in Y* .
- ▶ **Remark.** A subspace might have open sets that are not open in the full topological space. For instance, let $X = \mathbb{R}$ and $Y = [0, \infty)$. Then the half-open interval $[0, 1)$ is open in Y since $[0, 1) = Y \cap (-1, 1)$, but $[0, 1)$ is not open in X .
 - ▶ However, if $Y \subset X$ is open in X and $U \subset Y$ is open in Y , then U is open in X .

Subspace topology

- ▶ Equipping Y with the subspace topology, we will call Y a *subspace* of X . If a set U belongs to \mathcal{T}_Y we will often say that U is *open in Y* .
- ▶ **Remark.** A subspace might have open sets that are not open in the full topological space. For instance, let $X = \mathbb{R}$ and $Y = [0, \infty)$. Then the half-open interval $[0, 1)$ is open in Y since $[0, 1) = Y \cap (-1, 1)$, but $[0, 1)$ is not open in X .
 - ▶ However, if $Y \subset X$ is open in X and $U \subset Y$ is open in Y , then U is open in X .
- ▶ **Example.** The subspace topology on $\mathbb{Z} \subset \mathbb{R}$ is the discrete topology on \mathbb{Z} : the set $\{n\}$ is open in \mathbb{Z} for any integer n .

Subspace topology

- ▶ Equipping Y with the subspace topology, we will call Y a *subspace* of X . If a set U belongs to \mathcal{T}_Y we will often say that U is *open in Y* .
- ▶ **Remark.** A subspace might have open sets that are not open in the full topological space. For instance, let $X = \mathbb{R}$ and $Y = [0, \infty)$. Then the half-open interval $[0, 1)$ is open in Y since $[0, 1) = Y \cap (-1, 1)$, but $[0, 1)$ is not open in X .
 - ▶ However, if $Y \subset X$ is open in X and $U \subset Y$ is open in Y , then U is open in X .
- ▶ **Example.** The subspace topology on $\mathbb{Z} \subset \mathbb{R}$ is the discrete topology on \mathbb{Z} : the set $\{n\}$ is open in \mathbb{Z} for any integer n .
 - ▶ On the other hand, the subspace topology on $\mathbb{Q} \subset \mathbb{R}$ is *not* the discrete topology, essentially because any non-empty open interval in \mathbb{R} contains infinitely many rational numbers.

Subspace topology

- **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then

Subspace topology

- ▶ **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
 - ▶ (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,

Subspace topology

- **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
- (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,
 - (ii) if Z is a topological space and $f : X \rightarrow Z$ is a continuous map, then the restriction $f|_Y : Y \rightarrow Z$ is also continuous,

Subspace topology

- **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
- (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,
 - (ii) if Z is a topological space and $f : X \rightarrow Z$ is a continuous map, then the restriction $f|_Y : Y \rightarrow Z$ is also continuous,
 - (iii) a set $F \subset Y$ is closed in Y if and only if there is a set $G \subset X$ which is closed in X so that $F = Y \cap G$.

Subspace topology

- ▶ **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
 - ▶ (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,
 - ▶ (ii) if Z is a topological space and $f : X \rightarrow Z$ is a continuous map, then the restriction $f|_Y : Y \rightarrow Z$ is also continuous,
 - ▶ (iii) a set $F \subset Y$ is closed in Y if and only if there is a set $G \subset X$ which is closed in X so that $F = Y \cap G$.
- ▶ *Proof.* (i) follows from $\iota^{-1}(U) = U \cap Y$.

Subspace topology

- ▶ **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
 - ▶ (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,
 - ▶ (ii) if Z is a topological space and $f : X \rightarrow Z$ is a continuous map, then the restriction $f|_Y : Y \rightarrow Z$ is also continuous,
 - ▶ (iii) a set $F \subset Y$ is closed in Y if and only if there is a set $G \subset X$ which is closed in X so that $F = Y \cap G$.
- ▶ *Proof.* (i) follows from $\iota^{-1}(U) = U \cap Y$.
 - ▶ (ii) follows from $f|_Y^{-1}(U) = f^{-1}(U) \cap Y$.

Subspace topology

- **Proposition 3.2.** Let (X, \mathcal{T}) be a topological space, and let (Y, \mathcal{T}_Y) be a subspace. Then
- (i) the inclusion map $\iota : Y \rightarrow X$ given by $\iota(y) = y$ is cont.,
 - (ii) if Z is a topological space and $f : X \rightarrow Z$ is a continuous map, then the restriction $f|_Y : Y \rightarrow Z$ is also continuous,
 - (iii) a set $F \subset Y$ is closed in Y if and only if there is a set $G \subset X$ which is closed in X so that $F = Y \cap G$.
- *Proof.* (i) follows from $\iota^{-1}(U) = U \cap Y$.
- (ii) follows from $f|_Y^{-1}(U) = f^{-1}(U) \cap Y$.
 - (iii): Suppose $F \subset Y$ is closed in Y . Then there is an open set U in X so that $Y \setminus F = Y \cap U$. Note that

$$F = Y \setminus (Y \setminus F) = Y \setminus (Y \cap U) = Y \setminus U = Y \cap (X \setminus U).$$

Suppose $F = Y \cap G$ for some closed set $G \subset X$. Then

$$Y \setminus F = Y \setminus (Y \cap G) = Y \cap (X \setminus G),$$

which is open in Y .

Subspace topology: The pasting lemma

- **Lemma 3.5.** Let X be a topological space, and let $U, V \subset X$ be two open subsets such that $X = U \cup V$. Let $f : U \rightarrow Y$ and $g : V \rightarrow Y$ be two functions so that $f|_{U \cap V} = g|_{U \cap V}$. Then f and g are continuous w.r.t. the subspace topologies on U and V if and only if the function $h : X \rightarrow Y$ given by

$$h(x) = \begin{cases} f(x) & \text{if } x \in U, \\ g(x) & \text{if } x \in V, \end{cases}$$

is continuous.

Subspace topology: The pasting lemma

- **Lemma 3.5.** Let X be a topological space, and let $U, V \subset X$ be two open subsets such that $X = U \cup V$. Let $f : U \rightarrow Y$ and $g : V \rightarrow Y$ be two functions so that $f|_{U \cap V} = g|_{U \cap V}$. Then f and g are continuous w.r.t. the subspace topologies on U and V if and only if the function $h : X \rightarrow Y$ given by

$$h(x) = \begin{cases} f(x) & \text{if } x \in U, \\ g(x) & \text{if } x \in V, \end{cases}$$

is continuous.

- *Proof.* (\Rightarrow) follows from

$$\begin{aligned} h^{-1}(W) &= \{x \in X \mid h(x) \in W\} \\ &= \{x \in U \mid h(x) \in W\} \cup \{x \in V \mid h(x) \in W\} \\ &= \{x \in U \mid f(x) \in W\} \cup \{x \in V \mid g(x) \in W\} \\ &= f^{-1}(W) \cup g^{-1}(W). \end{aligned}$$

Subspace topology: The pasting lemma

- ▶ **Remark.** The exact same result would hold if we replaced “open” with “closed” everywhere in the statement of the pasting lemma.

Subspace topology: The pasting lemma

- ▶ **Remark.** The exact same result would hold if we replaced “open” with “closed” everywhere in the statement of the pasting lemma.
- ▶ The result would also be true if we replaced U and V with an infinite collection of open (or closed) sets $\{U_i\}_{i \in I}$ so that $X = \bigcup_{i \in I} U_i$.

Subspace topology: The pasting lemma

- ▶ **Remark.** The exact same result would hold if we replaced “open” with “closed” everywhere in the statement of the pasting lemma.
- ▶ The result would also be true if we replaced U and V with an infinite collection of open (or closed) sets $\{U_i\}_{i \in I}$ so that $X = \bigcup_{i \in I} U_i$.
- ▶ Note that if $Y \subset X$ is open (resp. closed) in X and $U \subset Y$ is open (resp. closed) in Y , then U is open (resp. closed) in X .

The product topology

- **Definition.** Let I be an index set. Given a set X , we define an I -tuple of elements of X to be a function $x : I \rightarrow X$.

The product topology

- ▶ **Definition.** Let I be an index set. Given a set X , we define an I -tuple of elements of X to be a function $x : I \rightarrow X$.
 - ▶ If i is an element of I , we often denote the value of x at i by x_i ; we call it the i th *coordinate* of x .

The product topology

- ▶ **Definition.** Let I be an index set. Given a set X , we define an I -tuple of elements of X to be a function $x : I \rightarrow X$.
 - ▶ If i is an element of I , we often denote the value of x at i by x_i ; we call it the i th *coordinate* of x .
 - ▶ And we often denote the function x itself by the symbol $(x_i)_{i \in I}$.

The product topology

- ▶ **Definition.** Let I be an index set. Given a set X , we define an I -tuple of elements of X to be a function $x : I \rightarrow X$.
 - ▶ If i is an element of I , we often denote the value of x at i by x_i ; we call it the i th *coordinate* of x .
 - ▶ And we often denote the function x itself by the symbol $(x_i)_{i \in I}$.
- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be an indexed family of sets; let $X = \bigcup_{i \in I} X_i$. The *Cartesian product* of this indexed family, denoted by

$$\prod_{i \in I} X_i,$$

is defined to be the set of all I -tuples $(x_i)_{i \in I}$ of elements of X such that $x_i \in X_i$ for each $i \in I$.

The product topology

- ▶ **Definition.** Let I be an index set. Given a set X , we define an I -tuple of elements of X to be a function $x : I \rightarrow X$.
 - ▶ If i is an element of I , we often denote the value of x at i by x_i ; we call it the i th *coordinate* of x .
 - ▶ And we often denote the function x itself by the symbol $(x_i)_{i \in I}$.
- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be an indexed family of sets; let $X = \bigcup_{i \in I} X_i$. The *Cartesian product* of this indexed family, denoted by

$$\prod_{i \in I} X_i,$$

is defined to be the set of all I -tuples $(x_i)_{i \in I}$ of elements of X such that $x_i \in X_i$ for each $i \in I$.

- ▶ That is, it is the set of all functions

$$x : I \rightarrow \bigcup_{i \in I} X_i$$

such that $x(i) \in X_i$ for each $i \in I$.

The product topology

- **Definition.** Let $\{X_i\}_{i \in I}$ be an indexed family of topological spaces. Let us take as a basis for a topology on the product space

$$\prod_{i \in I} X_i$$

the collection of all sets of the form

$$\prod_{i \in I} U_i,$$

where U_i is open in X_i , for each $i \in I$. (Exercise: check that this is indeed a basis.)

The product topology

- **Definition.** Let $\{X_i\}_{i \in I}$ be an indexed family of topological spaces. Let us take as a basis for a topology on the product space

$$\prod_{i \in I} X_i$$

the collection of all sets of the form

$$\prod_{i \in I} U_i,$$

where U_i is open in X_i , for each $i \in I$. (Exercise: check that this is indeed a basis.)

- The topology generated by this basis is called the *box topology*.

The product topology

► **Definition.** Let X be a set.

The product topology

- ▶ **Definition.** Let X be a set.
 - ▶ A *subbasis* \mathcal{C} for a topology on X is a collection of subsets that cover X , meaning that their union is all of X .

The product topology

- ▶ **Definition.** Let X be a set.
 - ▶ A *subbasis* \mathcal{C} for a topology on X is a collection of subsets that cover X , meaning that their union is all of X .
 - ▶ The topology $\mathcal{T}_{\mathcal{C}}$ generated by \mathcal{C} consists of all unions of all finite intersections of elements in \mathcal{C} .

The product topology

- ▶ **Definition.** Let X be a set.
 - ▶ A *subbasis* \mathcal{C} for a topology on X is a collection of subsets that cover X , meaning that their union is all of X .
 - ▶ The topology $\mathcal{T}_{\mathcal{C}}$ generated by \mathcal{C} consists of all unions of all finite intersections of elements in \mathcal{C} .
- ▶ **Remark.** $\mathcal{T}_{\mathcal{C}}$ is the coarsest topology containing \mathcal{C} , meaning that it has as few open sets as possible while still including the elements in \mathcal{C} as open sets.

The product topology

- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be a family of topological spaces, and let $X = \prod_{i \in I} X_i$.

The product topology

- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be a family of topological spaces, and let $X = \prod_{i \in I} X_i$.
 - ▶ For every $i \in I$, there is a natural map $\pi_i : X \rightarrow X_i$, called the projection onto X_i , which maps $\pi_i(x) = x(i)$, where we think of $x \in X$ as a map $I \rightarrow \bigcup_{i \in I} X_i$.

The product topology

- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be a family of topological spaces, and let $X = \prod_{i \in I} X_i$.
 - ▶ For every $i \in I$, there is a natural map $\pi_i : X \rightarrow X_i$, called the projection onto X_i , which maps $\pi_i(x) = x(i)$, where we think of $x \in X$ as a map $I \rightarrow \bigcup_{i \in I} X_i$.
 - ▶ We define a topology on X , called the *product topology*, to be the coarsest topology such that π_i is continuous for every i .

The product topology

- ▶ **Definition.** Let $\{X_i\}_{i \in I}$ be a family of topological spaces, and let $X = \prod_{i \in I} X_i$.
 - ▶ For every $i \in I$, there is a natural map $\pi_i : X \rightarrow X_i$, called the projection onto X_i , which maps $\pi_i(x) = x(i)$, where we think of $x \in X$ as a map $I \rightarrow \bigcup_{i \in I} X_i$.
 - ▶ We define a topology on X , called the *product topology*, to be the coarsest topology such that π_i is continuous for every i .
- ▶ **Remark.** The product topology on $X = \prod_{i \in I} X_i$ is generated by the subbasis \mathcal{C} which consists of all sets of the form $\pi_i^{-1}(U)$, where U is an open set in X_i .

The product topology

- ▶ Consider the case of a product of just two topological spaces (X_1, \mathcal{T}_{X_1}) and (X_2, \mathcal{T}_{X_2}) .

The product topology

- ▶ Consider the case of a product of just two topological spaces (X_1, \mathcal{T}_{X_1}) and (X_2, \mathcal{T}_{X_2}) .
 - ▶ Let U and V be open sets in X_1 and X_2 respectively. Then $\pi_1^{-1}(U) = U \times X_2$ and $\pi_2^{-1}(V) = X_1 \times V$ are examples of open sets in $X_1 \times X_2$. Their intersection is the set $\pi_1^{-1}(U) \cap \pi_2^{-1}(V) = U \times V$, and the topology on $X_1 \times X_2$ consists of all unions of this form.

The product topology

- ▶ Consider the case of a product of just two topological spaces (X_1, \mathcal{T}_{X_1}) and (X_2, \mathcal{T}_{X_2}) .
 - ▶ Let U and V be open sets in X_1 and X_2 respectively. Then $\pi_1^{-1}(U) = U \times X_2$ and $\pi_2^{-1}(V) = X_1 \times V$ are examples of open sets in $X_1 \times X_2$. Their intersection is the set $\pi_1^{-1}(U) \cap \pi_2^{-1}(V) = U \times V$, and the topology on $X_1 \times X_2$ consists of all unions of this form.
 - ▶ In symbols, if we let

$$\mathcal{B} = \{U \times V \mid U \in \mathcal{T}_{X_1}, V \in \mathcal{T}_{X_2}\},$$

then \mathcal{B} is a basis for the product topology on $X_1 \times X_2$.

The product topology

- ▶ Consider the case of a product of just two topological spaces (X_1, \mathcal{T}_{X_1}) and (X_2, \mathcal{T}_{X_2}) .
 - ▶ Let U and V be open sets in X_1 and X_2 respectively. Then $\pi_1^{-1}(U) = U \times X_2$ and $\pi_2^{-1}(V) = X_1 \times V$ are examples of open sets in $X_1 \times X_2$. Their intersection is the set $\pi_1^{-1}(U) \cap \pi_2^{-1}(V) = U \times V$, and the topology on $X_1 \times X_2$ consists of all unions of this form.
 - ▶ In symbols, if we let

$$\mathcal{B} = \{U \times V \mid U \in \mathcal{T}_{X_1}, V \in \mathcal{T}_{X_2}\},$$

then \mathcal{B} is a basis for the product topology on $X_1 \times X_2$.

- ▶ Similarly, open sets in $X = \prod_{i \in I} X_i$ are unions of sets of the form $\prod_{i \in I} U_i$, where U_i is open in X_i for each $i \in I$, and $U_i = X_i$ for all but finitely many i .

The product topology

- ▶ **Theorem 3.8.** The box topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i . The product topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i and U_i equals X_i except for finitely many values of i .

The product topology

- ▶ **Theorem 3.8.** The box topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i . The product topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i and U_i equals X_i except for finitely many values of i .
- ▶ **Remark.** Two things are clear. First, for finite products $\prod_{i=1}^n X_i$ the two topologies are precisely the same. Second, the box topology is in general finer than the product topology.

The product topology

- ▶ **Theorem 3.8.** The box topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i . The product topology on $\prod X_i$ has as basis all sets of the form $\prod U_i$, where U_i is open in X_i for each i and U_i equals X_i except for finitely many values of i .
- ▶ **Remark.** Two things are clear. First, for finite products $\prod_{i=1}^n X_i$ the two topologies are precisely the same. Second, the box topology is in general finer than the product topology.
- ▶ What is not so clear is why we prefer the product topology to the box topology. We will find that a number of important theorems about finite products will also hold for arbitrary products if we use the product topology, but not if we use the box topology.

The product topology

- **Theorem 3.9.** Let X be a topological space, and let $\{Y_i\}_{i \in I}$ be a family of topological spaces. A function $f : X \rightarrow \prod_{i \in I} Y_i$ consists of a family of functions $\{f_i\}_{i \in I}$ where $f_i : X \rightarrow Y_i$ for all $i \in I$. Then f is continuous iff f_i is continuous for every i .

The product topology

- ▶ **Theorem 3.9.** Let X be a topological space, and let $\{Y_i\}_{i \in I}$ be a family of topological spaces. A function $f : X \rightarrow \prod_{i \in I} Y_i$ consists of a family of functions $\{f_i\}_{i \in I}$ where $f_i : X \rightarrow Y_i$ for all $i \in I$. Then f is continuous iff f_i is continuous for every i .
- ▶ *Proof.* There are two things to prove.

The product topology

- ▶ **Theorem 3.9.** Let X be a topological space, and let $\{Y_i\}_{i \in I}$ be a family of topological spaces. A function $f : X \rightarrow \prod_{i \in I} Y_i$ consists of a family of functions $\{f_i\}_{i \in I}$ where $f_i : X \rightarrow Y_i$ for all $i \in I$. Then f is continuous iff f_i is continuous for every i .
- ▶ *Proof.* There are two things to prove.
 - ▶ Suppose f is continuous. Since each π_i is continuous, so is every $f_i = \pi_i \circ f$.

The product topology

- ▶ **Theorem 3.9.** Let X be a topological space, and let $\{Y_i\}_{i \in I}$ be a family of topological spaces. A function $f : X \rightarrow \prod_{i \in I} Y_i$ consists of a family of functions $\{f_i\}_{i \in I}$ where $f_i : X \rightarrow Y_i$ for all $i \in I$. Then f is continuous iff f_i is continuous for every i .
- ▶ *Proof.* There are two things to prove.
 - ▶ Suppose f is continuous. Since each π_i is continuous, so is every $f_i = \pi_i \circ f$.
 - ▶ Suppose all the f_i are continuous. We will show that the preimages of elements of the subbasis are open. That is, let U be an open set in $\prod_{i \in I} Y_i$ of the form $U = \pi_j^{-1}(V)$ where V is open in Y_j . Then $f^{-1}(U) = f^{-1}(\pi_j^{-1}(V)) = f_j^{-1}(V)$, which is open by assumption.

The product topology

- **Example.** Consider \mathbb{R}^ω , the countably infinite product of \mathbb{R} with itself. That is

$$\mathbb{R}^\omega = \prod_{i=1}^{\infty} X_i$$

where $X_i = \mathbb{R}$ for each i . Let us define a function $f : \mathbb{R} \rightarrow \mathbb{R}^\omega$ by the equation

$$f(t) = (t, t, t, \dots);$$

the i th coordinate function of f is the function $f_i(t) = t$.

The product topology

- **Example.** Consider \mathbb{R}^ω , the countably infinite product of \mathbb{R} with itself. That is

$$\mathbb{R}^\omega = \prod_{i=1}^{\infty} X_i$$

where $X_i = \mathbb{R}$ for each i . Let us define a function $f : \mathbb{R} \rightarrow \mathbb{R}^\omega$ by the equation

$$f(t) = (t, t, t, \dots);$$

the i th coordinate function of f is the function $f_i(t) = t$.

- Each of the coordinate functions $f_i : \mathbb{R} \rightarrow \mathbb{R}$ is continuous; therefore, the function f is continuous if \mathbb{R}^ω is given the product topology.

The product topology

- ▶ But f is not continuous if \mathbb{R}^ω is given the box topology. Consider, for example, the basis element

$$B = (-1, 1) \times (-1/2, 1/2) \times (-1/3, 1/3) \times \cdots$$

for the box topology. We assert that $f^{-1}(B)$ is not open in \mathbb{R} . If $f^{-1}(B)$ were open in \mathbb{R} , it would contain some interval $(-\delta, \delta)$ about the point 0. This would mean that $f((-\delta, \delta)) \subset B$. Applying π_i to both sides of the inclusion we obtain

$$f_i((-\delta, \delta)) = (-\delta, \delta) \subset (-1/i, 1/i)$$

for all i , a contradiction.