Mathematical Analysis II

conf.univ., dr. Elena Cojuhari

elena.cojuhari@mate.utm.md
Technical University of Moldova



2022

Stewart, p.621 ex. 23

Stewart, p.600 ex. 22

Adv. Eng. Math., p.17

Reduction to Separable Form (homogeneous ODE)

ex. 7,8,9,10,16, 17, p.18

Refresh from the previous seminar Ex.23. p.621 Stewart

$$\frac{dy}{dx} + P(x)y = Q(x)y'' \qquad \frac{\text{nonlinear DE, I}}{\text{Bernoulli DE}}$$

$$y(x) \sim 1 u(x)$$
 $u = y^{1-n}$
 $u' = (1-h) y^{-n} \cdot y'$

$$(1-n)y^{-n}y' + (1-n)P(x)y^{-1-h} = (1-n)Q(x)$$

 $u' + (1-n)P(x)u = (1-n)Q(x)$ linear DE, I

Refresh from the previous seminar

Ex. 25, p. 621 Stewart

$$y' + \frac{2}{x}y = \frac{y^3}{x^2}$$
 $u = y^{-2}$
 $u' = -\frac{2}{y^3}y'$
 $2y^3y' - \frac{4}{x}y^{-2} = -2 \cdot \frac{1}{x^2}$

Linear DE

Adv. Eng. Math. Ex.4, p.18

$$y' \sin 2\pi x = \pi y \cos 2\pi x$$
Separable ODE, I

$$\frac{dy}{dx} \sin 2\pi x = \pi y \cos 2\pi x$$

$$\frac{1}{y} dy = \pi \frac{\cos 2\pi x}{\sin 2\pi x} dx$$

Adv. Eng. Math. Ex.6, p.18

$$y' = e^{2x-1} y^2$$
Separable DE, I

$$y^{2}dy = e^{2x-1}dx$$

$$-y^{2} = \frac{1}{2}e^{2x-1} + ($$

Adv. Eng. Math. Ex.7, p.18

$$xy' = y + 2x^3 \sin^2 \frac{y}{x}$$
 homogeneous ODE, I

$$t = \frac{y}{x}$$

$$y = tx$$

$$y(x) \sim t(x)$$

$$x(t/x+t) = tx + 2x^3 \sin^2 t$$

$$2x^2t' + tx = tx + 2x^3 \sin^2 t$$

CONT.

 $x^2 t' = 2 x^3 sin^2 t$

Separable DE. I

$$\frac{dt}{\sin^2 t} = 2x dx$$

$$-ctgt = x^2 + C$$

$$ctg\frac{y}{x} = -x^2 + C$$

$$ctg = -x + c$$

$$y = x \operatorname{arcctg}(-x^2 + c)$$

Adv. Eng. Math. Ex.17, p.18

$$xy'=y+3x^{4}cos^{2}(\frac{t}{x})$$
, $y(1)=0$
 $y=tx$
 $y'=t'x+t$
 $x(t'x+t)=tx+3x^{4}cos^{2}t$
 $x^{2}t'+tx=tx+3x^{4}cos^{2}t$

$$x^{2}t' = 3x^{4} \cos^{2}t$$

$$\frac{dt}{\cos^{2}t} = 3x^{2} dx$$

$$t_{gt} = \alpha^{3} + c$$

the general solution:

$$y = x \operatorname{arcta}(x^{2}+c)$$
 $y(1)=0$

$$y = x \operatorname{arcta}(x^{2}+c)$$

$$y = 0 = 1^{3}+c$$

$$y = x \operatorname{arctg}(x^{3} + 1)$$

Homogeneous DE, I

$$\frac{dy}{dx} = \frac{x+y}{x-y}$$

$$\frac{dy}{dx} = \frac{x^2 + xy + y^3}{x^2}$$

$$y' = \frac{y}{x} - e$$

$$y = \frac{x + 2y - 4}{2x - y - 3}$$

$$(x^2+xy)dx-(xy+y^2)dy=0$$

$$M(x,y) dx + N(x,y) dy = 0$$

$$M(dx_1dy) = (\alpha x)^2 + dx \cdot dy = d^2(x^2 + xy) = 0$$

$$= d^2M(x,y)$$

The given equation is a homogeneous DE

y=tx

Exercise. CONT.

$$(x^2+xy)dx-(xy+y^2)dy=0$$

The given equation is a homogeneous DE

The substitution:
$$y=tx$$
 or $y(x)=xt(x)$

$$(x^{2}+xtx)dx - (xtx+(tx)^{2})(xtt+tdx)=0$$

$$(x^{2}(1+t)-tx^{2}t(1+t)]dx-x\cdot x^{2}t(1+t)dt=0$$

$$x^{2}(1+t)(1-t^{2})dx=x^{3}t(1+t)dt$$

Exercise. CONT.

Exercise. CONT.

$$x^{2}(1+t)(1-t^{2}) dx = x^{3}t(1+t)dt$$

$$\frac{dx}{x} = \frac{tdt}{1-t^{2}}$$

$$h/1-t^2/=2h/12t+C \Rightarrow h/1-t^2=h/2t^2e$$

the general solution:

$$x^2 - y^2 = c$$

$$y' = (y + 4x)^{2} \text{ ODE, } I$$

$$y(x) \sim V(x)$$

$$V = y + 4x$$

$$V' = y' + 4$$

Separable ODE, I
$$\mathcal{U}' - 4 = \mathcal{U}^2$$

$$\mathcal{U}' = \mathcal{U}^2 + 4$$

$$\frac{dr}{dx} = v^2 + 4$$

$$\frac{dy}{dx} = \frac{x+y-3}{x-y-1}$$

ODE, I , it is reducible to a homogeneous DE

Substitution:

#= f(=) dy = f(3) da

$$\begin{cases} x = 2 + u \Rightarrow dx = du \\ y = 1 + v \Rightarrow dy = dv \\ y(x) \sim v V(u) \end{cases}$$

$$\frac{dv}{du} = \frac{u+v}{u-v}$$
 is a homogeneous DE, I $v(u) \sim 1 + (u)$

$$\frac{udt+tdu}{du} = \frac{u+tu}{u-tu}$$

$$dt + u+tu$$

$$u\frac{dt}{du} + t = \frac{u + tu}{u - tu}$$

$$u\frac{dt}{du} = \frac{1+t}{1-t} - t$$

$$U \frac{dt}{du} = \frac{1+t^2}{1-t}$$
 Separable ODE, I

$$\frac{1-t}{1+t^{2}} dt = \frac{du}{u}$$

$$\int \frac{dt}{1+t^{2}} - \int \frac{t}{1+t^{2}} dt = \int \frac{du}{u}$$

$$arctgt - \frac{1}{2} ln(1+t^{2}) = ln|u| + C$$

$$arctgt = ln|cu\sqrt{1+t^{2}}|$$

$$cu\sqrt{1+t^2} = e^{arctgt}$$

$$cu\sqrt{1+\frac{v^2}{u^2}} = e^{arctgt}$$

CONT

$$C\sqrt{u^2+v^2} = e$$
the general solution:
$$C\sqrt{(\chi-z)^2+(\chi-1)^2} = e$$

$$C\sqrt{(\chi-z)^2+(\chi-1)^2} = e$$

$$(x^2+2y^2)$$
 dy = xy dx

$$(x^{2}+2t^{2}x^{2})(x dt+t dx) = x^{2}t dx$$

$$x^{3}(1+2t^{2}) dt = (-x^{2}t-2t^{2}x^{2}+x^{2}t)dx$$

$$x^{3}(1+2t^{2}) dt = -2t^{3}x^{2}dx$$

CONT

$$\frac{1+2t^2}{t^3}dt = -2\frac{x^2}{x^3}dx$$

$$\int t^{-3}dt + 2\int \frac{dt}{t} = -2\int \frac{dz}{x}$$

$$-\frac{1}{2t^2} + 2\ln|t| = -2\ln|x| + C$$

$$\ln|cxt| = \frac{1}{4} \cdot \frac{1}{t^2}$$
the general solution:
$$\frac{x^2}{x^2} = \frac{1}{2} \cdot \frac{1}{2}$$

Exercise.

$$(3z^2y+y^3)dy - (x^3+3xy^2)dx = 0$$

 $y = tx$; $dy = xdt + tdx$

 $(3x^{2}+x+(tx)^{3})(x+t+tdx)-(x^{2}+3x(t^{2})^{2})dx=0$

 $x(3x^3t+x^3t^3)dt+(3x^3t^2+x^3t^4-x^3-3x^2t^2)dx$

 $x^4t(3+t^2)dt = x^3(1-t^4)dx$ $\frac{t(3+t^2)}{4-t^4}dt = \frac{dx}{x}$

