

# Taylor approximations

conf.dr. Bostan Viorel

Spring 2014

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$**  is

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$**  is

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$$

For example, Taylor series of  $f(x) = e^x$  at  $a = 2.7$  is

$$f(2.7) + f'(2.7)(x - 2.7) + \frac{f''(2.7)}{2!}(x - 2.7)^2 + \frac{f'''(2.7)}{3!}(x - 2.7)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$$

For example, Taylor series of  $f(x) = e^x$  at  $a = 2.7$  is

$$f(2.7) + f'(2.7)(x - 2.7) + \frac{f''(2.7)}{2!}(x - 2.7)^2 + \frac{f'''(2.7)}{3!}(x - 2.7)^3 + \dots$$
$$e^{2.7} + e^{2.7}(x - 2.7) + \frac{e^{2.7}}{2!}(x - 2.7)^2 + \frac{e^{2.7}}{3!}(x - 2.7)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$$

For example, Taylor series of  $f(x) = e^x$  at  $a = 5$  is

$$f(5) + f'(5)(x - 5) + \frac{f''(5)}{2!}(x - 5)^2 + \frac{f'''(5)}{3!}(x - 5)^3 + \dots$$
$$e^5 + e^5(x - 5) + \frac{e^5}{2!}(x - 5)^2 + \frac{e^5}{3!}(x - 5)^3 + \dots$$



# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

For example, Taylor series of  $f(x) = \cos x$  at  $a = \frac{\pi}{6}$  is

$$f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x - \frac{\pi}{6}\right) + \frac{f''\left(\frac{\pi}{6}\right)}{2!}\left(x - \frac{\pi}{6}\right)^2 + \frac{f'''\left(\frac{\pi}{6}\right)}{3!}\left(x - \frac{\pi}{6}\right)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

For example, Taylor series of  $f(x) = \cos x$  at  $a = \frac{\pi}{6}$  is

$$f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x - \frac{\pi}{6}\right) + \frac{f''\left(\frac{\pi}{6}\right)}{2!}\left(x - \frac{\pi}{6}\right)^2 + \frac{f'''\left(\frac{\pi}{6}\right)}{3!}\left(x - \frac{\pi}{6}\right)^3 + \dots$$
$$\cos \frac{\pi}{6} - \sin \frac{\pi}{6} \left(x - \frac{\pi}{6}\right) - \frac{\cos \frac{\pi}{6}}{2!} \left(x - \frac{\pi}{6}\right)^2 + \frac{\sin \frac{\pi}{6}}{3!} \left(x - \frac{\pi}{6}\right)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$

**Taylor series of function  $f$  at  $a$  is**

$$f(a) + f'(a)(x-a) + \frac{f''(a)}{2!}(x-a)^2 + \frac{f'''(a)}{3!}(x-a)^3 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x-a)^n$$

For example, Taylor series of  $f(x) = \cos x$  at  $a = \frac{\pi}{6}$  is

$$f\left(\frac{\pi}{6}\right) + f'\left(\frac{\pi}{6}\right)\left(x - \frac{\pi}{6}\right) + \frac{f''\left(\frac{\pi}{6}\right)}{2!}\left(x - \frac{\pi}{6}\right)^2 + \frac{f'''\left(\frac{\pi}{6}\right)}{3!}\left(x - \frac{\pi}{6}\right)^3 + \dots$$
$$\cos \frac{\pi}{6} - \sin \frac{\pi}{6} \left(x - \frac{\pi}{6}\right) - \frac{\cos \frac{\pi}{6}}{2!} \left(x - \frac{\pi}{6}\right)^2 + \frac{\sin \frac{\pi}{6}}{3!} \left(x - \frac{\pi}{6}\right)^3 + \dots$$
$$\frac{\sqrt{3}}{2} - \frac{1}{2} \left(x - \frac{\pi}{6}\right) - \frac{\sqrt{3}}{4} \left(x - \frac{\pi}{6}\right)^2 + \frac{1}{12} \left(x - \frac{\pi}{6}\right)^3 + \dots$$

# Taylor and Maclaurin Series

If  $a = 0$  then Taylor series is called **Maclaurin series of function  $f$**  and is defined by

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

# Taylor and Maclaurin Series

If  $a = 0$  then Taylor series is called **Maclaurin series of function  $f$**  and is defined by

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n$$

For example, Maclaurin series of  $f(x) = \cos x$  is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$

# Taylor and Maclaurin Series

If  $a = 0$  then Taylor series is called **Maclaurin series of function  $f$**  and is defined by

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

For example, Maclaurin series of  $f(x) = \cos x$  is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\cos 0 - \sin 0 \cdot x - \frac{\cos 0}{2!}x^2 + \frac{\sin 0}{3!}x^3 + \frac{\cos 0}{4!}x^4 + \dots$$

# Taylor and Maclaurin Series

If  $a = 0$  then Taylor series is called **Maclaurin series of function  $f$**  and is defined by

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!}x^n$$

For example, Maclaurin series of  $f(x) = \cos x$  is

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$
$$\cos 0 - \sin 0 \cdot x - \frac{\cos 0}{2!}x^2 + \frac{\sin 0}{3!}x^3 + \frac{\cos 0}{4!}x^4 + \dots$$
$$1 - 0 \cdot x - \frac{1}{2!}x^2 + \frac{0}{3!}x^3 + \frac{1}{4!}x^4 - \frac{0}{5!}x^5 - \frac{1}{6!}x^6 + \dots$$



# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$  we can construct a **Taylor series of function  $f$  at  $a$**  is

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$  we can construct a **Taylor series of function  $f$  at  $a$**  is

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

or a **Maclaurin series of function  $f$**

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$  we can construct a **Taylor series of function  $f$  at  $a$**  is

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

or a **Maclaurin series of function  $f$**

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$

- 1 Under what conditions will these series converge (absolutely)?

# Taylor and Maclaurin Series

For a given function  $f : \mathbb{R} \rightarrow \mathbb{R}$  and a given number  $a \in \mathbb{R}$  we can construct a **Taylor series of function  $f$  at  $a$**  is

$$f(a) + f'(a)(x - a) + \frac{f''(a)}{2!}(x - a)^2 + \frac{f'''(a)}{3!}(x - a)^3 + \dots$$

or a **Maclaurin series of function  $f$**

$$f(0) + f'(0)x + \frac{f''(0)}{2!}x^2 + \frac{f'''(0)}{3!}x^3 + \frac{f^{(4)}(0)}{4!}x^4 + \dots$$

- 1 Under what conditions will these series converge (absolutely)?
- 2 If the series converge, will its sum be  $f(x)$ ?

# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots \quad \forall x \in \mathbb{R}$$

# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots \quad \forall x \in \mathbb{R}$$

$$\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \quad \forall x \in \mathbb{R}$$

# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots \quad \forall x \in \mathbb{R}$$

$$\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \quad \forall x \in \mathbb{R}$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots, \quad |x| < 1$$



# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots \quad \forall x \in \mathbb{R}$$

$$\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \quad \forall x \in \mathbb{R}$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots, \quad |x| < 1$$

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + x^4 - x^5 + x^6 + \dots, \quad |x| < 1$$

# Taylor and Maclaurin Series

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots \quad \forall x \in \mathbb{R}$$

$$\sin x = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots \quad \forall x \in \mathbb{R}$$

$$\cos x = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots \quad \forall x \in \mathbb{R}$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots, \quad |x| < 1$$

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + x^4 - x^5 + x^6 + \dots, \quad |x| < 1$$

$$\ln(1+x) = x - \frac{1}{3}x^3 + \frac{1}{5}x^5 - \frac{1}{7}x^7 + \dots \quad |x| < 1$$

# Taylor Polynomials

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n + \frac{1}{(n+1)!}x^{n+1} + \dots$$

# Taylor Polynomials

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n + \frac{1}{(n+1)!}x^{n+1} + \dots$$

# Taylor Polynomials


$$e^x = \underbrace{1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n}_{=T_n(x)} + \underbrace{\frac{1}{(n+1)!}x^{n+1} + \dots}_{=R_n(x)}$$

# Taylor Polynomials


$$e^x = \underbrace{1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n}_{=T_n(x)} + \underbrace{\frac{1}{(n+1)!}x^{n+1} + \dots}_{=R_n(x)}$$

$$e^x = T_n(x) + R_n(x)$$

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots, \quad |x| < 1$$




$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$







$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$




$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$




$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$







$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$



$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$







$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$



$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$



# Function whose Taylor series doesn't converge to the function itself

$$f(x) = \begin{cases} 0, & \text{if } x = 0 \\ e^{-\frac{1}{x^2}}, & \text{if } x \neq 0 \end{cases}$$





# Srīnivāsa Aiyangār Rāmānujan

better known as Srinivasa Iyengar Ramanujan  
(22 December 1887 – 26 April 1920)



# Srīnivāsa Aiyangār Rāmānujan

$$1 - 5 \left( \frac{1}{2} \right)^3 + 9 \left( \frac{1 \cdot 3}{2 \cdot 4} \right)^3 - 13 \left( \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^3 + \dots = \frac{2}{\pi}$$

$$1 - 5 \left( \frac{1}{2} \right)^3 + 9 \left( \frac{1 \cdot 3}{2 \cdot 4} \right)^3 - 13 \left( \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^3 + \dots = \frac{2}{\pi}$$

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}} = 3$$

$$1 - 5 \left( \frac{1}{2} \right)^3 + 9 \left( \frac{1 \cdot 3}{2 \cdot 4} \right)^3 - 13 \left( \frac{1 \cdot 3 \cdot 5}{2 \cdot 4 \cdot 6} \right)^3 + \dots = \frac{2}{\pi}$$

$$\sqrt{1 + 2\sqrt{1 + 3\sqrt{1 + 4\sqrt{1 + \dots}}}} = 3$$

$$1 + \frac{1}{1 \cdot 3} + \frac{1}{1 \cdot 3 \cdot 5} + \frac{1}{1 \cdot 3 \cdot 5 \cdot 7} + \dots + \frac{1}{1 + \frac{1}{1 + \frac{2}{1 + \frac{3}{1 + \frac{4}{1 + \dots}}}}} = \sqrt{\frac{\pi e}{2}}$$

One day Hardy came to Ramanujan who was at that time lying in a London;s hospital.

One day Hardy came to Ramanujan who was at that time lying in a London;s hospital.

Hardy said to Ramanujan that the cab he was taking to arrive at the hospital had a quite **unitersting** number: **1729**.

One day Hardy came to Ramanujan who was at that time lying in a London;s hospital.

Hardy said to Ramanujan that the cab he was taking to arrive at the hospital had a quite **unitersting** number: **1729**.

Immediately, Ramanujan replied that it was actually a very interesting number mathematically...



One day Hardy came to Ramanujan who was at that time lying in a London;s hospital.

Hardy said to Ramanujan that the cab he was taking to arrive at the hospital had a quite **unitersting** number: **1729**.

Immediately, Ramanujan replied that it was actually a very interesting number mathematically...

being the smallest number representable in two different ways as a sum of two cubes:

$$1729 = 1^3 + 12^3 = 9^3 + 10^3.$$

One day Hardy came to Ramanujan who was at that time lying in a London;s hospital.

Hardy said to Ramanujan that the cab he was taking to arrive at the hospital had a quite **unitersting** number: **1729**.

Immediately, Ramanujan replied that it was actually a very interesting number mathematically...

being the smallest number representable in two different ways as a sum of two cubes:

$$1729 = 1^3 + 12^3 = 9^3 + 10^3.$$

Generalizations of this idea have spawned the notion of "taxicab numbers".