Mathematics for Computer Science


Prof.dr.hab. Viorel Bostan

Technical University of Moldova viorel.bostan@adm.utm.md

Lecture 10














FIGURE 12. The U_8 skeleton (a), its classical realization (b), and its low-dimensional spectral realizations. (The $(1+\sqrt{2})$ -realization is only pseudo-classical: the hexagonal faces are not regular.)









Graph Theory History





Leonard Euler, 1707-1783

Graph Theory started in 1736 when Leonard Euler while walking in Konigsberg posed a problem known today as "Bridges of Konigsberg problem".

Graph Theory History





Leonard Euler, 1707-1783

Graph Theory started in 1736 when Leonard Euler while walking in Konigsberg posed a problem known today as "Bridges of Konigsberg problem".

Later, Vandermonde studied the knot problem using methods developed by Leibniz.

Graph Theory History





Leonard Euler, 1707-1783

Graph Theory started in 1736 when Leonard Euler while walking in Konigsberg posed a problem known today as "Bridges of Konigsberg problem".


Later, Vandermonde studied the knot problem using methods developed by Leibniz.

Euler also introduced his famous formula relating the number of edges, vertices, and faces of a convex polyhedron, generalized later by Cauchy.

Seven Bridges of Konigsberg




Gedenkblatt jur sechshundert jährigen Dubelfeier der Koniglichen Baupt und Residen; Stadt Romigsberg in Breufen.




Seven Bridges of Konigsberg





Seven Bridges of Konigsberg





Euler's Question

Is it possible to find a walk through the city that would cross each bridge once and only once?













The 4 Color Conjecture was first proposed on October 23, 1852 by Francis Guthrie, while trying to color the map of counties of England, noticed that only four different colors were needed.

10 / 67



Theorem (4 Color Theorem, Intuitive Formulation)

Given any separation of a plane into contiguous regions, called a map, the regions can be colored using at most four colors so that no two adjacent regions have the same color.



Theorem (4 Color Theorem, Intuitive Formulation)

Given any separation of a plane into contiguous regions, called a map, the regions can be colored using at most four colors so that no two adjacent regions have the same color.

There were several early failed attempts at proving the theorem one of which was given by A. Kempe in 1879.



Theorem (4 Color Theorem, Intuitive Formulation)

Given any separation of a plane into contiguous regions, called a map, the regions can be colored using at most four colors so that no two adjacent regions have the same color.

There were several early failed attempts at proving the theorem one of which was given by A. Kempe in 1879.

Another "proof" was given by P.G. Tait in 1880.



Theorem (4 Color Theorem, Intuitive Formulation)

Given any separation of a plane into contiguous regions, called a map, the regions can be colored using at most four colors so that no two adjacent regions have the same color.

There were several early failed attempts at proving the theorem one of which was given by A. Kempe in 1879.

Another "proof" was given by P.G. Tait in 1880.

It was not until 1890 that Kempe's proof was shown incorrect by P. Heawood, who also proved the 5 color theorem and generalized the 4 color conjecture to surfaces of arbitrary genus.



Theorem (4 Color Theorem, Intuitive Formulation)

Given any separation of a plane into contiguous regions, called a map, the regions can be colored using at most four colors so that no two adjacent regions have the same color.

There were several early failed attempts at proving the theorem one of which was given by A. Kempe in 1879.

Another "proof" was given by P.G. Tait in 1880.

It was not until 1890 that Kempe's proof was shown incorrect by P. Heawood, who also proved the 5 color theorem and generalized the 4 color conjecture to surfaces of arbitrary genus.

In 1891 Tait's proof was shown incorrect by J. Petersen.



The 4 color theorem was proven in 1976 by K. Appel and W .Haken.



The 4 color theorem was proven in 1976 by K. Appel and W .Haken.

It was the first major theorem to be proved using a computer.



The 4 color theorem was proven in 1976 by K. Appel and W .Haken.

It was the first major theorem to be proved using a computer.

Appel and Haken's approach started by showing that there is a particular set of 1,936 maps, each of which cannot be part of a smallest-sized counterexample to the 4 color theorem. Appel and Haken used a special-purpose computer program to confirm that each of these maps had this property.



The 4 color theorem was proven in 1976 by K. Appel and W .Haken.


It was the first major theorem to be proved using a computer.

Appel and Haken's approach started by showing that there is a particular set of 1,936 maps, each of which cannot be part of a smallest-sized counterexample to the 4 color theorem. Appel and Haken used a special-purpose computer program to confirm that each of these maps had this property.

A simpler proof using the same ideas and still relying on computers was published in 1997 by Robertson, Sanders, Seymour, and Thomas.


Graph Theory: Computer Networks






Graphs in Computer Science





Graphs in Computer Science







How many handshakes (at most) are between,



How many handshakes (at most) are between, say, ...



How many handshakes (at most) are between, say, ...

Donald Trump and



How many handshakes (at most) are between, say, ...

Donald Trump and Toader de la Stâna din Vale?



How many handshakes (at most) are between, say, ...

Donald Trump and Toader de la Stâna din Vale?







How many handshakes (at most) are between, say, ...

Donald Trump and Toader de la Stâna din Vale?















How many handshakes are between Queen Elizabeth and a zulu warrior?















How many handshakes are between Queen Elizabeth and a zulu warrior?













Six degrees of separation is the theory that everyone and everything is six or fewer steps away, by way of introduction, from any other person in the world.

6 Degree of separation conjecture



A chain of "a friend of a friend" statements can be made to connect any two people in a maximum of six steps.

6 Degree of separation conjecture



A chain of "a friend of a friend" statements can be made to connect any two people in a maximum of six steps.

It was originally set out by F.Karinthy in 1929 and popularized by a play written by J.Guare.

6 Degree of separation conjecture




A chain of "a friend of a friend" statements can be made to connect any two people in a maximum of six steps.

It was originally set out by F.Karinthy in 1929 and popularized by a play written by J.Guare. Karinthy wrote:

A fascinating game grew out of this discussion. One of us suggested performing the following experiment to prove that the population of the Earth is closer together now than they have ever been before. We should select any person from the 1.5 billion inhabitants of the Earth – anyone, anywhere at all. He bet us that, using no more than five individuals, one of whom is a personal acquaintance, he could contact the selected individual using nothing except the network of personal acquaintances.







A game, called 6 degrees of Kevin Bacon, was created by four students in US.



A game, called **6 degrees of Kevin Bacon**, was created by four students in US. The goal is to link any actor to Kevin Bacon through no more than 6 connections, where 2 actors are connected if they have appeared in a movie or commercial together.



A game, called **6 degrees of Kevin Bacon**, was created by four students in US. The goal is to link any actor to Kevin Bacon through no more than 6 connections, where 2 actors are connected if they have appeared in a movie or commercial together.





A game, called **6 degrees of Kevin Bacon**, was created by four students in US. The goal is to link any actor to Kevin Bacon through no more than 6 connections, where 2 actors are connected if they have appeared in a movie or commercial together.



Google made it possible to search for any given actor's Bacon Number through their search engine.



Facebook and scientists have lanched a project called "Small World" in order to check the "6 Degree of Separation Theory".



Facebook and scientists have lanched a project called "Small World" in order to check the "6 Degree of Separation Theory".

The new study assessed in 2011 the relationships between 721 million active users (more than 10% of the global population) of the social network.



Facebook and scientists have lanched a project called "Small World" in order to check the "6 Degree of Separation Theory".

The new study assessed in 2011 the relationships between 721 million active users (more than 10% of the global population) of the social network.

Using state-of-the-art algorithms. . . we were able to approximate the number of 'hops' (degrees of separation) between all pairs of individuals on Facebook."



Facebook and scientists have lanched a project called "Small World" in order to check the "6 Degree of Separation Theory".

The new study assessed in 2011 the relationships between 721 million active users (more than 10% of the global population) of the social network.

Using state-of-the-art algorithms. . . we were able to approximate the number of 'hops' (degrees of separation) between all pairs of individuals on Facebook."

Researchers found that the average number of connections between people has dropped since 2008 to four.



Facebook and scientists have lanched a project called "Small World" in order to check the "6 Degree of Separation Theory".

The new study assessed in 2011 the relationships between 721 million active users (more than 10% of the global population) of the social network.

Using state-of-the-art algorithms. . . we were able to approximate the number of 'hops' (degrees of separation) between all pairs of individuals on Facebook."

Researchers found that the average number of connections between people has dropped since 2008 to four.

We found that six degrees actually overstates the number of links between typical pairs of users: While 99.6% of all pairs of users are connected by paths with 5 degrees (6 hops), 92% are connected by only four degrees (5 hops)





Each person in the world (at least among the 1.59 billion people active on Facebook) is connected to every other person by an average of (3.57) three and a half other people.



Paul Erdős, who died in 1996 at the age of 83, was the most prolific mathematician in history.



Paul Erdős, who died in 1996 at the age of 83, was the most prolific mathematician in history.

He boasts 1525 published papers (and counting, with posthumous publications) and 511 collaborations;



Paul Erdős, who died in 1996 at the age of 83, was the most prolific mathematician in history.

He boasts 1525 published papers (and counting, with posthumous publications) and 511 collaborations;

Erdős number is computed as follows: Paul Erdős itself receives the Erdős number 0. An author's Erdős number is 1 if he has co-authored a paper with Erdős, 2 if he has co-authored a paper with someone who has co-authored a paper with Erdős, etc.



Paul Erdős, who died in 1996 at the age of 83, was the most prolific mathematician in history.

He boasts 1525 published papers (and counting, with posthumous publications) and 511 collaborations;

Erdős number is computed as follows: Paul Erdős itself receives the Erdős number 0. An author's Erdős number is 1 if he has co-authored a paper with Erdős, 2 if he has co-authored a paper with someone who has co-authored a paper with Erdős, etc.

Thus, 9779 people can lay claim to having an Erdős number of at most 2.



Paul Erdős, who died in 1996 at the age of 83, was the most prolific mathematician in history.

He boasts 1525 published papers (and counting, with posthumous publications) and 511 collaborations;

Erdős number is computed as follows: Paul Erdős itself receives the Erdős number 0. An author's Erdős number is 1 if he has co-authored a paper with Erdős, 2 if he has co-authored a paper with someone who has co-authored a paper with Erdős, etc.

Thus, 9779 people can lay claim to having an Erdős number of at most 2.

Some famous names have low Erdős numbers— Bill Gates has an Erdős number of 4, John von Neumann is 3, and Albert Einstein is 2.


Erdős number (2017)



Erdős number	
1	511
2	9267
3	33605
4	83642
5	87760
6	40014
7	11591
8	3146
9	819
10	244
11	68
12	23
13	5

Thus the median Erdős number is 5; the mean is 4.65, and the standard deviation is 1.21.







My Erdos number is:



My Erdos number is: 4



My Erdos number is: 4

Here is the proof:

- 4. Viorel Bostan coauthored with Weimin Han
- 3. Weimin Han coauthored with Michael J. McAsey
- 2. Michael J. McAsey coauthored with Lee A. Rubel
- 1. Lee A. Rubel coauthored with Paul Erdős

You can check your Erdos number on the website: http://www.ams.org/mathscinet/collaborationDistance.html




































Shortest path in graphs








Google Maps Fastest Roundtrip Solver



Airline routes in Europe







Transportation Map







Definition

Let V be an arbitrary set. A **graph** is a pair of sets G = (V, E) satisfying $E \subset V^2 \equiv V \times V$.



Definition

Let V be an arbitrary set. A **graph** is a pair of sets G = (V, E) satisfying $E \subset V^2 \equiv V \times V$.

Note that the elements of E are pairs of elements from V. To avoid ambiguities in notation, always assume that $V \cap E = \emptyset$.



Definition

Let V be an arbitrary set. A **graph** is a pair of sets G = (V, E) satisfying $E \subset V^2 \equiv V \times V$.

Note that the elements of E are pairs of elements from V. To avoid ambiguities in notation, always assume that $V \cap E = \emptyset$.

The elements of V are called **vertices** (or nodes, or points) of the graph G, the elements of E are called its **edges** (or lines, or arcs).



Definition

Let V be an arbitrary set. A **graph** is a pair of sets G = (V, E) satisfying $E \subset V^2 \equiv V \times V$.

Note that the elements of E are pairs of elements from V. To avoid ambiguities in notation, always assume that $V \cap E = \emptyset$.

The elements of V are called **vertices** (or nodes, or points) of the graph G, the elements of E are called its **edges** (or lines, or arcs).

The usual way to picture a graph is by drawing a dot for each vertex and joining two of these dots by a line, if the corresponding two vertices form an edge.

Graph definition. Example




$$G = (V, E), V = \{0, 1, ..., 6\}, E = \{(0, 2), (0, 4), (2, 4), (1, 3), (4, 5)\}.$$

Graph definition. Example




$$G = (V, E), V = \{0, 1, ..., 6\}, E = \{(0, 2), (0, 4), (2, 4), (1, 3), (4, 5)\}.$$



Graph definition. Example



$$G = (V, E), V = \{0, 1, ..., 6\}, E = \{(0, 2), (0, 4), (2, 4), (1, 3), (4, 5)\}.$$



Important!

Just how these dots and lines are drawn is considered irrelevant:

All that matters is the information which pairs of vertices form edges!

Order of a graph




Definition

A graph with vertex set V is said to be a graph on V. The **vertex set** of a graph G is referred to as V(G), its **edge set** as E(G).

Definition

The number of vertices of a graph G is called its **order**, written as |G|; its number of edges is denoted by ||G||.



Finite and trivial graphs. Vertex incident with an edge



Definition

Graphs are **finite** or **infinite** according to their order; unless otherwise stated, the graphs we consider are all finite.

Definition

For the empty graph (\emptyset, \emptyset) we simply write \emptyset .

Definition

A graph of order 0 or 1 is called **trivial**.

Finite and trivial graphs. Vertex incident with an edge



Definition

Graphs are **finite** or **infinite** according to their order; unless otherwise stated, the graphs we consider are all finite.

Definition

For the empty graph (\emptyset, \emptyset) we simply write \emptyset .

Definition

A graph of order 0 or 1 is called **trivial**.

Definition

A vertex v is **incident** with an edge e if $v \in e$; then e is an edge at v. The two vertices incident with an edge are its **endvertices** or ends, and an edge joins its ends. An edge $\{x, y\}$ is usually written as xy (or yx).

Adjacent vertices and edges. Complete graphs



Definition

The set of all the edges in E at a vertex v is denoted by E(v).

Definition

Two vertices x and y of G are **adjacent**, or neighbours, if xy is an edge of G. Two edges e and f are **adjacent** if they have an endvertex in common.

Adjacent vertices and edges. Complete graphs



Definition

The set of all the edges in E at a vertex v is denoted by E(v).

Definition

Two vertices x and y of G are **adjacent**, or neighbours, if xy is an edge of G. Two edges e and f are **adjacent** if they have an endvertex in common.

Definition

If all the vertices of G are pairwise adjacent, then G is called **complete**. A complete graph on n vertices is denoted K^n ; a K^3 is called

Adjacent vertices and edges. Complete graphs



Definition

The set of all the edges in E at a vertex v is denoted by E(v).

Definition


Two vertices x and y of G are **adjacent**, or neighbours, if xy is an edge of G. Two edges e and f are **adjacent** if they have an endvertex in common.

Definition

If all the vertices of G are pairwise adjacent, then G is called **complete**. A complete graph on n vertices is denoted K^n ; a K^3 is called a **triangle**.

Complete graphs






Complete graphs K^3 (triangle), K^4 and K^5

Complete graphs






Complete graphs K^3 (triangle), K^4 and K^5

Problem

Count the number of edges in complete graph K^n .

Complete graphs





Graph K^{16}




Definition

Let G=(V,E) and G'=(V',E') be two graphs. We call G and G' isomorphic, and write $G\simeq G'$, if there exists a bijection $\varphi:V\to V'$ with $xy\in E\Longleftrightarrow \varphi(x)\varphi(y)\in E'\forall x,y\in V$. Such a map φ is called an **isomorphism**; if G=G', it is called an **automorphism**.



Definition


Let G=(V,E) and G'=(V',E') be two graphs. We call G and G' isomorphic, and write $G\simeq G'$, if there exists a bijection $\varphi:V\to V'$ with $xy\in E\Longleftrightarrow \varphi(x)\varphi(y)\in E'\forall x,y\in V$. Such a map φ is called an **isomorphism**; if G=G', it is called an **automorphism**.





Definition


Let G=(V,E) and G'=(V',E') be two graphs. We call G and G' isomorphic, and write $G\simeq G'$, if there exists a bijection $\varphi:V\to V'$ with $xy\in E\Longleftrightarrow \varphi(x)\varphi(y)\in E'\forall x,y\in V$. Such a map φ is called an **isomorphism**; if G=G', it is called an **automorphism**.



$$\varphi(1) = B$$
, $\varphi(2) = A$, $\varphi(3) = C$, $\varphi(4) = D$

Isomorphic graphs. Graph invariant






We do not distinguish between isomorphic graphs, and write G = G' rather than $G \simeq G'$.

Isomorphic graphs. Graph invariant






We do not distinguish between isomorphic graphs, and write G = G' rather than $G \simeq G'$.

Definition

A map taking graphs as arguments is called a **graph invariant** if it assigns equal values to isomorphic graphs.

Isomorphic graphs. Graph invariant






We do not distinguish between isomorphic graphs, and write G = G' rather than $G \simeq G'$.


Definition

A map taking graphs as arguments is called a **graph invariant** if it assigns equal values to isomorphic graphs.


The number of vertices and number of edges of a graph are 2 simple graph invariant.
















$$\varphi(1) = 1$$
, $\varphi(2) = 3$, $\varphi(3) = 5$, $\varphi(4) = 2$, $\varphi(5) = 4$








$$\varphi(1) = 1$$
, $\varphi(2) = 3$, $\varphi(3) = 5$, $\varphi(4) = 2$, $\varphi(5) = 4$





$$\varphi(1)=1,\quad \varphi(2)=3,\quad \varphi(3)=5,\quad \varphi(4)=2,\quad \varphi(5)=4$$

Remark

No one has yet found a general procedure for determining whether two graphs are isomorphic that is guaranteed to run in polynomial time in |V|.



Definition

If $V' \subset V$ and $E' \subset E$, then G' is called a **subgraph** of G, written as $G' \subset G$.



Definition

If $V' \subset V$ and $E' \subset E$, then G' is called a **subgraph** of G, written as $G' \subset G$.

Definition

If $G' \subset G$ and G' contains all the edges $xy \in E$ with $x, y \in V'$, then G' is an **induced** subgraph of G and write G' = G[V'].



Definition

If $V' \subset V$ and $E' \subset E$, then G' is called a **subgraph** of G, written as $G' \subset G$.

Definition

If $G' \subset G$ and G' contains all the edges $xy \in E$ with $x, y \in V'$, then G' is an **induced** subgraph of G and write G' = G[V'].

Thus if $U \subset V$ is any set of vertices, then G[U] denotes the graph on U whose edges are precisely the edges of G with both ends in U.



Definition

If $V' \subset V$ and $E' \subset E$, then G' is called a **subgraph** of G, written as $G' \subset G$.

Definition

If $G' \subset G$ and G' contains all the edges $xy \in E$ with $x, y \in V'$, then G' is an **induced** subgraph of G and write G' = G[V'].

Thus if $U \subset V$ is any set of vertices, then G[U] denotes the graph on U whose edges are precisely the edges of G with both ends in U.

If H is a subgraph of G, not necessarily induced, we abbreviate G[V(H)] to G[H].



Definition

If $V' \subset V$ and $E' \subset E$, then G' is called a **subgraph** of G, written as $G' \subset G$.

Definition

If $G' \subset G$ and G' contains all the edges $xy \in E$ with $x, y \in V'$, then G' is an **induced** subgraph of G and write G' = G[V'].

Thus if $U \subset V$ is any set of vertices, then G[U] denotes the graph on U whose edges are precisely the edges of G with both ends in U.


If H is a subgraph of G, not necessarily induced, we abbreviate G[V(H)] to G[H].

Definition

Let $G' \subset G$. Then G' is a **spanning subgraph** of G if V' spans all of G, i.e. if V' = V. In other words subgraph G' contains all the vertices of G.

Induced subgraph. Spanning subgraph.






■ G_1 and G_2 are subgraphs of G.

Induced subgraph. Spanning subgraph.






- G_1 and G_2 are subgraphs of G.
- G_1 is an induced subgraph of G, while G_2 is not.

Induced subgraph. Spanning subgraph.





- G_1 and G_2 are subgraphs of G.
- G_1 is an induced subgraph of G, while G_2 is not.
- G_2 is a spanning subgraph of G, while G_1 is not.

Union, intersection and difference of two graphs



Definition (Graph union and intersection)

We set $G \cup G' = (V \cup V', E \cup E')$ and $G \cap G' = (V \cap V', E \cap E')$. If $G \cap G' = \emptyset$, then G and G' are called **disjoint**.

Definition (Graph difference)

If U is any set of vertices (usually of G), we write G - U for G[V - U]. In other words, G - U is obtained from G by deleting all the vertices in U - V and their incident edges.


Definition

If $U = \{v\}$ is a singleton, we write G - v rather than $G - \{v\}$. Instead of G - V(G') we simply write G - G'.

For a subset F of V^2 we write G - F = (V; E - F) and $G + F = (V; E \cup F)$; as above, $G - \{e\}$ and $G + \{e\}$ are abbreviated to G - e and G + e.


Union and intersection of two graphs





Difference of two graphs






Star operation on graphs



Definition (Star operation on graphs)

If G_1 and G_2 are disjoint, we denote by $G_1 * G_2$ the graph obtained from $G_1 \cup G_2$ by joining all the vertices of G_1 to all the vertices of G_2 .




$$G_1$$
 (blue), G_2 (red), $G_1 * G_2$ (green)

Star operation on graphs



For example, $K^2 * K^3 = K^5$.




$$K^2$$
 (blue), K^3 (red), $K^2 * K^3 = K^5$ (green).

Star operation on graphs



For example, $K^2 * K^3 = K^5$.



$$K^2 * K^3 = K^5$$
 (green).

Complement of a graph



Definition (Graph complement)


The **complement** \overline{G} of G is the graph on V with edge set $V^2 - E$. In other words, eliminate all the edges from G and add all the edges not present in G.

Complement of a graph



Definition (Graph complement)

The **complement** \overline{G} of G is the graph on V with edge set $V^2 - E$. In other words, eliminate all the edges from G and add all the edges not present in G.



A graph isomorphic to its complement.

Neighbors of a vertex. Vertex degree.



Definition

The set of **neighbors** of a vertex v in G is denoted by $N_G(v)$, or N(v). For $U \subset V$, the neighbors in V - U of vertices in U are called **neighbors** of U, N(U).

Neighbors of a vertex. Vertex degree.



Definition

The set of **neighbors** of a vertex v in G is denoted by $N_G(v)$, or N(v). For $U \subset V$, the neighbors in V - U of vertices in U are called **neighbors** of U, N(U).

Definition

Degree (or valency) $d_G(v) = d(v)$ of v is the number of edges at v, that is the number of neighbors of v.

A vertex of degree 0 is called **isolated** vertex.

Neighbors of a vertex. Vertex degree.



Definition

The set of **neighbors** of a vertex v in G is denoted by $N_G(v)$, or N(v). For $U \subset V$, the neighbors in V - U of vertices in U are called **neighbors** of U, N(U).

Definition

Degree (or valency) $d_G(v) = d(v)$ of v is the number of edges at v, that is the number of neighbors of v.

A vertex of degree 0 is called **isolated** vertex.

Definition


Minimum degree of G and **maximum degree** of G are numbers:


$$\delta(G) = \min_{v \in V} \{d_G(v)\}, \quad \Delta(G) = \max_{v \in V} \{d_G(v)\}$$

Vertex degree. Minimum and maximum degrees.



Consider the graph G = (V, E):






$$d_G(E) = d(E) = 5$$
, $d(C) = 0$, $\delta(G) = 0$, $\Delta(G) = 5$

k-regular graphs. Cubic graph



Definition (Regular graph)


If all the vertices of a graph G have the same degree k, then G is called k— **regular**, or simply **regular**. A 3—regular graph is called **cubic**.



A 2-regular graph (blue) and a 3-regular (cubic) graph (red).

Adjacency Matrix






$$a_{ij} = \left\{ egin{array}{ll} 1, & ext{if } (v_i, v_j) \in E \ 0, & ext{otherwise} \end{array}
ight.$$

Adjacency Matrix





$$a_{ij} = \begin{cases} 1, & \text{if } (v_i, v_j) \in E \\ 0, & \text{otherwise} \end{cases}$$

```
    1
    2
    3
    4
    5
    6
    8

    1
    0
    1
    1
    1
    1
    1

    2
    1
    0
    0
    1
    0
    0
    0

    3
    1
    0
    0
    1
    1
    0
    1


    4
    0
    1
    1
    0
    1
    1
    0

    5
    1
    0
    1
    1
    0
    1
    1
    0
    1

    6
    1
    0
    1
    0
    1
    1
    0
    1
    0
```

Incidence Matrix






$$a_{ij} = \begin{cases} 1, & \text{if } v_i \text{ and } e_j \text{ are incident} \\ 0, & \text{otherwise} \end{cases}$$

Incidence Matrix






$$a_{ij} = \begin{cases} 1, & \text{if } v_i \text{ and } e_j \text{ are incident} \\ 0, & \text{otherwise} \end{cases}$$

	1	2	3	4	5	6	7	8	9	10	11
1	1	1	0	1	0	0	0	0	0	0	0
2	1	0	1	0	0	0	0	0	0	0	0
3	0	1	0	0	0	0	1	0	1	0	0
4	0	0	1	0	1	1	0	0	0	0	0
5	0	0	0	1	1	0	1	0	0	1	0
6	0	0	0	0	0	1	0	0	0	1	1
7	0	0	0	0	0	0	0	1	1	0	1


Adjacency List (Simple Graph)






Adjacency List (Simple Graph)






Adjacency List (Directed Graph)





Adjacency List (Directed Graph)





Average Degree of a Graph



Definition

The number

$$d(G) = \frac{1}{|G|} \sum_{v \in V} d_G(v)$$

is called the **average degree** of G.

Average Degree of a Graph



Definition

The number

$$d(G) = \frac{1}{|G|} \sum_{v \in V} d_G(v)$$

is called the **average degree** of *G*.

Clearly, average degree

$$\delta(G) \leq d(G) \leq \Delta(G)$$
.

The average degree quantifies globally what is measured locally by the vertex degrees: the number of edges of G per vertex.

Average Degree of a Graph



Definition

The number

$$d(G) = \frac{1}{|G|} \sum_{v \in V} d_G(v)$$

is called the **average degree** of G.

Clearly, average degree

$$\delta(G) \leq d(G) \leq \Delta(G)$$
.

The average degree quantifies globally what is measured locally by the vertex degrees: the number of edges of G per vertex.

Sometimes it will be convenient to express this ratio directly, as

$$\varepsilon(G) = \frac{\|G\|}{|G|}.$$



The quantities d and ε are related. Indeed, if we sum up all the vertex degrees in G, we count every edge exactly twice: once from each of its ends.



The quantities d and ε are related. Indeed, if we sum up all the vertex degrees in G, we count every edge exactly twice: once from each of its ends.

Thus,

$$||G|| = \frac{1}{2} \sum_{v \in V} d_G(v) = \frac{1}{2} d(G) \cdot |G|$$



The quantities d and ε are related. Indeed, if we sum up all the vertex degrees in G, we count every edge exactly twice: once from each of its ends.

Thus,

$$||G|| = \frac{1}{2} \sum_{v \in V} d_G(v) = \frac{1}{2} d(G) \cdot |G|$$

and therefore

$$\varepsilon(G)=\frac{1}{2}d(G).$$

Corollary

The number of vertices of odd degree in a graph is always even.



The quantities d and ε are related. Indeed, if we sum up all the vertex degrees in G, we count every edge exactly twice: once from each of its ends.

Thus,

$$||G|| = \frac{1}{2} \sum_{v \in V} d_G(v) = \frac{1}{2} d(G) \cdot |G|$$

and therefore

$$\varepsilon(G)=\frac{1}{2}d(G).$$

Corollary


The number of vertices of odd degree in a graph is always even.

Lemma (Handshaking lemma)

For any graph the sum of the degrees of vertices equals twice the number of edges.

Average Degree





$$G_1$$
 (left), G_2 (right). $d(G_1)=4$, $\varepsilon(G_1)=2$, $d(G_2)=\frac{22}{7}=3.14$, $\varepsilon(G_2)=\frac{11}{7}=1.57$

Lecture summary



- Graph theory history;
- Six degrees of separation;
- Graph applications;
- Graph definition, Graph representations;
- Complete graph, Isomorphic graphs;
- Subgraphs, Induced and spanning graph;
- Operations on graphs;
- Adjacency and Incident matrices;
- Vertex degree, Average degree, Regular graph;
- Handshaking lemma.