

# Numerical Analysis

Prof.dr.hab. Bostan Viorel

# Calculation of functions

Using hand calculations, a hand calculator, or a computer, what are the basic operations of which we are capable?

# Calculation of functions

Using hand calculations, a hand calculator, or a computer, what are the basic operations of which we are capable?

Addition, subtraction, multiplication, and division

# Calculation of functions

Using hand calculations, a hand calculator, or a computer, what are the basic operations of which we are capable?

Addition, subtraction, multiplication, and division

(and even this will usually require a truncation of the quotient at some point).

# Calculation of functions

Using hand calculations, a hand calculator, or a computer, what are the basic operations of which we are capable?

Addition, subtraction, multiplication, and division

(and even this will usually require a truncation of the quotient at some point).

In addition, we can make logical decisions, such as deciding:

$$a > b, \quad a = b, \quad a < b$$

# Calculation of functions

Using hand calculations, a hand calculator, or a computer, what are the basic operations of which we are capable?

Addition, subtraction, multiplication, and division

(and even this will usually require a truncation of the quotient at some point).

In addition, we can make logical decisions, such as deciding:

$$a > b, \quad a = b, \quad a < b$$

Furthermore, we can carry out only a finite number of such operations.

# Calculation of functions

In evaluating functions  $f(x)$  we are limited to the evaluation of polynomials:

# Calculation of functions

In evaluating functions  $f(x)$  we are limited to the evaluation of polynomials:

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$



# Calculation of functions

In evaluating functions  $f(x)$  we are limited to the evaluation of polynomials:

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

We will discuss later how to evaluate **efficiently** the polynomials;

# Calculation of functions

In evaluating functions  $f(x)$  we are limited to the evaluation of polynomials:

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

We will discuss later how to evaluate **efficiently** the polynomials;

How to evaluate other functions such as

$$e^x, \quad \cos x, \quad \log x, \quad \tan x, \quad \text{etc?}$$

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

Consider evaluating it for  $x$  near to 0.

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

Consider evaluating it for  $x$  near to 0.

Look for a polynomial  $p(x)$  whose values will be the same as those of  $e^x$  within acceptable accuracy.

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

Consider evaluating it for  $x$  near to 0.

Look for a polynomial  $p(x)$  whose values will be the same as those of  $e^x$  to within acceptable accuracy.

Begin with a linear polynomial  $P_1(x) = a_0 + a_1x$ .

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

Consider evaluating it for  $x$  near to 0.

Look for a polynomial  $p(x)$  whose values will be the same as those of  $e^x$  to within acceptable accuracy.

Begin with a linear polynomial  $P_1(x) = a_0 + a_1x$ .

Then to make its graph look like that of  $e^x$ , we ask that the graph of  $y = P_1(x)$  be tangent to that of  $y = e^x$  at  $x = 0$ .

# Taylor polynomial approximations

Consider function  $f(x) = e^x$

Consider evaluating it for  $x$  near to 0.

Look for a polynomial  $p(x)$  whose values will be the same as those of  $e^x$  to within acceptable accuracy.


Begin with a linear polynomial  $P_1(x) = a_0 + a_1x$ .

Then to make its graph look like that of  $e^x$ , we ask that the graph of  $y = P_1(x)$  be tangent to that of  $y = e^x$  at  $x = 0$ .

and we get  $P_1(x) = 1 + x$ .



# Taylor polynomial approximations



# Taylor polynomial approximations

Look for a quadratic polynomial  $P_2(x) = a_0 + a_1x + a_2x^2$

# Taylor polynomial approximations

Look for a quadratic polynomial  $P_2(x) = a_0 + a_1x + a_2x^2$

Make it tangent; and to determine  $a_2$ , ask that  $P_2(x)$  and  $e^x$  have the same “curvature” at the origin.

# Taylor polynomial approximations

Look for a quadratic polynomial  $P_2(x) = a_0 + a_1x + a_2x^2$

Make it tangent; and to determine  $a_2$ , ask that  $P_2(x)$  and  $e^x$  have the same “curvature” at the origin.

Combining these requirements, we have for  $f(x) = e^x$  that

$$P_2(0) = f(0), \quad P_2'(0) = f'(0), \quad P_2''(0) = f''(0)$$

# Taylor polynomial approximations

Look for a quadratic polynomial  $P_2(x) = a_0 + a_1x + a_2x^2$

Make it tangent; and to determine  $a_2$ , ask that  $P_2(x)$  and  $e^x$  have the same “curvature” at the origin.

Combining these requirements, we have for  $f(x) = e^x$  that

$$P_2(0) = f(0), \quad P_2'(0) = f'(0), \quad P_2''(0) = f''(0)$$

This yields the approximation

$$P_2(x) = 1 + x + \frac{1}{2}x^2.$$

# Taylor polynomial approximations

We continue this pattern, looking for a polynomial

$$P_n(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

# Taylor polynomial approximations

We continue this pattern, looking for a polynomial

$$P_n(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

Such that

$$P_n(0) = f(0), \quad P'_n(0) = f'(0), \quad P''_n(0) = f''(0), \quad P_n^{(n)}(0) = f^{(n)}(0)$$

# Taylor polynomial approximations

We continue this pattern, looking for a polynomial

$$P_n(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

Such that

$$P_n(0) = f(0), \quad P'_n(0) = f'(0), \quad P''_n(0) = f''(0), \quad P_n^{(n)}(0) = f^{(n)}(0)$$

This leads to the formula

$$P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$



# Taylor polynomial approximations

We continue this pattern, looking for a polynomial

$$P_n(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

Such that

$$P_n(0) = f(0), \quad P'_n(0) = f'(0), \quad P''_n(0) = f''(0), \quad P_n^{(n)}(0) = f^{(n)}(0)$$

This leads to the formula

$$P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

In the vicinity of 0 we will have an approximation of  $f(x) = e^x$  by polynomials  $P_n(x)$  :

$$e^x \approx P_n(x)$$

# Taylor polynomial approximations

We continue this pattern, looking for a polynomial

$$P_n(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

Such that

$$P_n(0) = f(0), \quad P'_n(0) = f'(0), \quad P''_n(0) = f''(0), \quad P_n^{(n)}(0) = f^{(n)}(0)$$

This leads to the formula

$$P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

In the vicinity of 0 we will have an approximation of  $f(x) = e^x$  by polynomials  $P_n(x)$  :

$$e^x \approx P_n(x)$$


Moreover, as  $n \rightarrow \infty$  it can be shown that  $P_n(x) \rightarrow e^x$

# Taylor polynomial approximations

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$

# Taylor polynomial approximations

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \dots, \quad x \in \mathbb{R}$$




# Taylor polynomial approximations

What are the problems when evaluating points  $x$  that are far from 0?

# Taylor polynomial approximations

What are the problems when evaluating points  $x$  that are far from 0?



# Taylor polynomial error formula

Let  $f(x)$  be a given function, and assume it has derivatives around some point  $x = a$  (with as many derivatives as we find necessary).

# Taylor polynomial error formula

Let  $f(x)$  be a given function, and assume it has derivatives around some point  $x = a$  (with as many derivatives as we find necessary).

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas



# Taylor polynomial error formula

Let  $f(x)$  be a given function, and assume it has derivatives around some point  $x = a$  (with as many derivatives as we find necessary).

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas

$$f(x) - P_n(x) = \frac{1}{(n+1)!} (x-a)^{n+1} f^{(n+1)}(\xi)$$

# Taylor polynomial error formula

Let  $f(x)$  be a given function, and assume it has derivatives around some point  $x = a$  (with as many derivatives as we find necessary).

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas

$$f(x) - P_n(x) = \frac{1}{(n+1)!} (x-a)^{n+1} f^{(n+1)}(\xi)$$

or

$$f(x) - P_n(x) = \frac{1}{n!} \int_a^x (x-t)^n f^{n+1}(t) dt$$

# Taylor polynomial error formula

Let  $f(x)$  be a given function, and assume it has derivatives around some point  $x = a$  (with as many derivatives as we find necessary).

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas

$$f(x) - P_n(x) = \frac{1}{(n+1)!} (x-a)^{n+1} f^{(n+1)}(\xi)$$

or

$$f(x) - P_n(x) = \frac{1}{n!} \int_a^x (x-t)^n f^{(n+1)}(t) dt$$

where point  $\xi$  is restricted to the interval bounded by  $x$  and  $a$ , and otherwise  $\xi$  is unknown.

# Taylor polynomial error formula

For the case  $a = 0$  we have

# Taylor polynomial error formula

For the case  $a = 0$  we have

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas

$$f(x) - P_n(x) = \frac{1}{(n+1)!} x^{n+1} f^{(n+1)}(\xi)$$

or

$$f(x) - P_n(x) = \frac{1}{n!} \int_0^x (x-t)^n f^{n+1}(t) dt$$

# Taylor polynomial error formula

For the case  $a = 0$  we have

For the error in the Taylor polynomial  $P_n(x)$ , we have the formulas

$$f(x) - P_n(x) = \frac{1}{(n+1)!} x^{n+1} f^{(n+1)}(\xi)$$

or

$$f(x) - P_n(x) = \frac{1}{n!} \int_0^x (x-t)^n f^{(n+1)}(t) dt$$

where point  $\xi$  is restricted to the interval bounded by  $x$  and  $0$ , and otherwise  $\xi$  is unknown.

# Taylor polynomial approximations

For the case of considered function  $f(x) = e^x$  we get

# Taylor polynomial approximations

For the case of considered function  $f(x) = e^x$  we get

$$e^x - P_n(x) = \frac{1}{(n+1)!} x^n e^{\xi}$$

with  $\xi$  located between 0 and  $x$ .



# Taylor polynomial approximations

For the case of considered function  $f(x) = e^x$  we get

$$e^x - P_n(x) = \frac{1}{(n+1)!} x^n e^{\xi}$$

with  $\xi$  located between 0 and  $x$ .

Note that for  $x \approx 0$ , we must have  $\xi \approx 0$  and

$$e^x - P_n(x) \approx \frac{1}{(n+1)!} x^n$$

# Taylor polynomial approximations

For the case of considered function  $f(x) = e^x$  we get

$$e^x - P_n(x) = \frac{1}{(n+1)!} x^n e^{\xi}$$

with  $\xi$  located between 0 and  $x$ .


Note that for  $x \approx 0$ , we must have  $\xi \approx 0$  and

$$e^x - P_n(x) \approx \frac{1}{(n+1)!} x^{n+1}$$


This last term is also the final term in  $P_{n+1}(x)$ , and thus

$$e^x - P_n(x) \approx P_{n+1}(x) - P_n(x)$$


$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$



$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$







$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$



$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \frac{x^8}{8!} - \frac{x^{10}}{10!} + \frac{x^{12}}{12!} - \frac{x^{14}}{14!} + \frac{x^{16}}{16!} - \frac{x^{18}}{18!} + \dots$$




# Taylor polynomial approximations

$$\frac{1}{1-x} = 1 + x + x^2 + x^3 + x^4 + x^5 + x^6 + \dots, \quad |x| < 1$$



# Function whose Taylor series doesn't converge to the function itself

$$f(x) = \begin{cases} 0, & \text{if } x = 0 \\ e^{-\frac{1}{x^2}}, & \text{if } x \neq 0 \end{cases}$$



# Computing approximation to e

Coming back to earlier approximation

$$e^x \approx P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$



# Computing approximation to e

Coming back to earlier approximation

$$e^x \approx P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

Then let  $x = 1$  to get

$$e \approx P_n(1) = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!}, \quad x \in \mathbb{R}$$

# Computing approximation to e

Coming back to earlier approximation

$$e^x \approx P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

Then let  $x = 1$  to get

$$e \approx P_n(1) = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!}, \quad x \in \mathbb{R}$$

For the error,

$$e - P_n(1) = \frac{1}{(n+1)!}e^{\xi}, \quad 0 \leq \xi \leq 1$$

# Computing approximation to e

Coming back to earlier approximation

$$e^x \approx P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

Then let  $x = 1$  to get

$$e \approx P_n(1) = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!}, \quad x \in \mathbb{R}$$

For the error,

$$e - P_n(1) = \frac{1}{(n+1)!}e^{\xi}, \quad 0 \leq \xi \leq 1$$

To bound the error, we have

$$e^0 \leq e^{\xi} \leq e^1$$

# Computing approximation to e

Coming back to earlier approximation

$$e^x \approx P_n(x) = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \dots + \frac{1}{n!}x^n, \quad x \in \mathbb{R}$$

Then let  $x = 1$  to get

$$e \approx P_n(1) = 1 + 1 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{n!}, \quad x \in \mathbb{R}$$

For the error,

$$e - P_n(1) = \frac{1}{(n+1)!}e^{\xi}, \quad 0 \leq \xi \leq 1$$

To bound the error, we have

$$e^0 \leq e^{\xi} \leq e^1$$

$$\frac{1}{(n+1)!} \leq e - P_n(1) \leq \frac{1}{(n+1)!}e$$

# Taylor polynomial approximations

Suppose we would like to have an approximation accurate for  $e$  within  $10^{-5}$ .

# Taylor polynomial approximations

Suppose we would like to have an approximation accurate for  $e$  within  $10^{-5}$ .

How large we should choose  $n$ ?

# Taylor polynomial approximations

Suppose we would like to have an approximation accurate for  $e$  within  $10^{-5}$ .

How large we should choose  $n$ ?

From the formula

$$\frac{1}{(n+1)!} \leq e - P_n(1) \leq \frac{1}{(n+1)!} e$$

# Taylor polynomial approximations

Suppose we would like to have an approximation accurate for  $e$  within  $10^{-5}$ .

How large we should choose  $n$ ?

From the formula

$$\frac{1}{(n+1)!} \leq e - P_n(1) \leq \frac{1}{(n+1)!} e$$

we choose  $n$  large enough to have

$$\frac{e}{(n+1)!} \leq 10^{-5}$$



# Taylor polynomial approximations

Suppose we would like to have an approximation accurate for  $e$  within  $10^{-5}$ .

How large we should choose  $n$ ?

From the formula

$$\frac{1}{(n+1)!} \leq e - P_n(1) \leq \frac{1}{(n+1)!} e$$

we choose  $n$  large enough to have

$$\frac{e}{(n+1)!} \leq 10^{-5}$$

which is true if  $n \geq 8$ .

# Taylor polynomial approximations

In fact,

$$e - P_8(1) \leq \frac{e}{9!} \approx 7.5 \cdot 10^{-6}$$

# Taylor polynomial approximations

In fact,

$$e - P_8(1) \leq \frac{e}{9!} \approx 7.5 \cdot 10^{-6}$$

Then calculate  $P_8(1)$  :

$$P_8(1) = 2 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{8!} = 2.71827877,$$

# Taylor polynomial approximations

In fact,

$$e - P_8(1) \leq \frac{e}{9!} \approx 7.5 \cdot 10^{-6}$$

Then calculate  $P_8(1)$  :

$$P_8(1) = 2 + \frac{1}{2!} + \frac{1}{3!} + \frac{1}{4!} + \dots + \frac{1}{8!} = 2.71827877,$$

and true error is

$$e - P_8(1) \approx 3.06 \cdot 10^{-6}$$

# Evaluation of polynomials

Consider having a polynomial

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

# Evaluation of polynomials

Consider having a polynomial

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

which you need to evaluate for many values of  $x$ .

# Evaluation of polynomials

Consider having a polynomial

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

which you need to evaluate for many values of  $x$ .

How do you evaluate it?

# Evaluation of polynomials

Consider having a polynomial

$$p(x) = a_0 + a_1x + a_2x^2 + \dots a_{n-1}x^{n-1} + a_nx^n$$

which you need to evaluate for many values of  $x$ .

How do you evaluate it?

This may seem a strange question, but the answer is not as obvious as you might think.



# Evaluation of polynomials

The standard way, written in a loose algorithmic format is:

# Evaluation of polynomials

The standard way, written in a loose algorithmic format is:

```
poly = a(0)
for j = 1 : n
    poly = poly + a(j)*x^j
end
```

# Evaluation of polynomials

The standard way, written in a loose algorithmic format is:

```
poly = a(0)
for j = 1 : n
    poly = poly + a(j)*x^j
end
```

To compare the costs of different numerical methods, we do an operations count, and then we compare these for the competing methods.

# Evaluation of polynomials

The standard way, written in a loose algorithmic format is:

```
poly = a(0)
for j = 1 : n
    poly = poly + a(j)*x^j
end
```

To compare the costs of different numerical methods, we do an operations count, and then we compare these for the competing methods.

Above, the counts are as follows:

$$\begin{aligned} \text{additions} &: n \\ \text{multiplications} &: 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2} \end{aligned}$$

# Evaluation of polynomials

Next, do the terms  $x^j$  recursively:

$$x^j = x \cdot x^{j-1}$$

# Evaluation of polynomials

Next, do the terms  $x^j$  recursively:

$$x^j = x \cdot x^{j-1}$$

Then to compute  $\{x^2, x^3, \dots, x^n\}$  will cost  $n - 1$  multiplications.

# Evaluation of polynomials

Next, do the terms  $x^j$  recursively:

$$x^j = x \cdot x^{j-1}$$

Then to compute  $\{x^2, x^3, \dots, x^n\}$  will cost  $n - 1$  multiplications.

The evaluation algorithm becomes

```
poly = a(0) + a(1)*x
power = x
for j = 2 : n
    power = x*power
    poly = poly + a(j)*power
end
```

# Evaluation of polynomials

Next, do the terms  $x^j$  recursively:

$$x^j = x \cdot x^{j-1}$$

Then to compute  $\{x^2, x^3, \dots, x^n\}$  will cost  $n - 1$  multiplications.

The evaluation algorithm becomes

```
poly = a(0) + a(1)*x
power = x
for j = 2 : n
    power = x*power
    poly = poly + a(j)*power
end
```

The total operations cost is

*additions* :  $n$

*multiplications* :  $n + n - 1 = 2n - 1$



# Evaluation of polynomials

When  $n$  is even moderately large, this is much less than for the first method of evaluating  $p(x)$ .

# Evaluation of polynomials

When  $n$  is even moderately large, this is much less than for the first method of evaluating  $p(x)$ .

For example, with  $n = 20$ ,

# Evaluation of polynomials

When  $n$  is even moderately large, this is much less than for the first method of evaluating  $p(x)$ .

For example, with  $n = 20$ ,

the first method has 210 multiplications,

# Evaluation of polynomials

When  $n$  is even moderately large, this is much less than for the first method of evaluating  $p(x)$ .

For example, with  $n = 20$ ,

the first method has 210 multiplications,

whereas the second has 39 multiplications.

# Evaluation of polynomials

# Evaluation of polynomials

Consider **nested multiplication**.

# Evaluation of polynomials

Consider **nested multiplication**.

As examples of particular degrees, write

$$n = 2 : p(x) = a_0 + x(a_1 + a_2x)$$

$$n = 3 : p(x) = a_0 + x(a_1 + x(a_2 + a_3x))$$

$$n = 4 : p(x) = a_0 + x(a_1 + x(a_2 + x(a_3 + a_4x)))$$

# Evaluation of polynomials

Consider **nested multiplication**.

As examples of particular degrees, write

$$n = 2 : p(x) = a_0 + x(a_1 + a_2x)$$

$$n = 3 : p(x) = a_0 + x(a_1 + x(a_2 + a_3x))$$

$$n = 4 : p(x) = a_0 + x(a_1 + x(a_2 + x(a_3 + a_4x)))$$

These contain, respectively, 2, 3, and 4 multiplications.



# Evaluation of polynomials

Consider **nested multiplication**.

As examples of particular degrees, write

$$n = 2 : p(x) = a_0 + x(a_1 + a_2x)$$

$$n = 3 : p(x) = a_0 + x(a_1 + x(a_2 + a_3x))$$

$$n = 4 : p(x) = a_0 + x(a_1 + x(a_2 + x(a_3 + a_4x)))$$

These contain, respectively, 2, 3, and 4 multiplications.

This is less than the second method, which would have need 3, 5, and 7 multiplications,

# Evaluation of polynomials

Consider **nested multiplication**.

As examples of particular degrees, write

$$n = 2 : p(x) = a_0 + x(a_1 + a_2x)$$

$$n = 3 : p(x) = a_0 + x(a_1 + x(a_2 + a_3x))$$

$$n = 4 : p(x) = a_0 + x(a_1 + x(a_2 + x(a_3 + a_4x)))$$

These contain, respectively, 2, 3, and 4 multiplications.

This is less than the second method, which would have need 3, 5, and 7 multiplications,

The first method will need 3, 6, and 10 multiplications.

# Evaluation of polynomials

For the general case, write

$$p(x) = a_0 + x(a_1 + x(a_2 + \dots + x(a_{n-1} + a_n x) \dots))$$

# Evaluation of polynomials

For the general case, write

$$p(x) = a_0 + x(a_1 + x(a_2 + \dots + x(a_{n-1} + a_n x) \dots))$$

This requires  $n$  multiplications, which is only about half that for the preceding method.

# Evaluation of polynomials

For the general case, write

$$p(x) = a_0 + x(a_1 + x(a_2 + \dots + x(a_{n-1} + a_n x) \dots))$$

This requires  $n$  multiplications, which is only about half that for the preceding method.

For an algorithm, write

```
poly = a(n)
for j = n-1 : -1 : 0
    poly = a(j) + x * poly
end
```

# Evaluation of polynomials

For the general case, write

$$p(x) = a_0 + x(a_1 + x(a_2 + \dots + x(a_{n-1} + a_n x) \dots))$$

This requires  $n$  multiplications, which is only about half that for the preceding method.

For an algorithm, write

```
poly = a(n)
for j = n-1 : -1 : 0
    poly = a(j) + x * poly
end
```

With all three methods, the number of additions is  $n$ ; but the number of multiplications can be dramatically different for large values of  $n$ .

# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt, \quad x \neq 0$$

# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt, \quad x \neq 0$$


We use Taylor polynomials to approximate this function, to obtain a way to compute it with accuracy and simplicity.



# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt, \quad x \neq 0$$

We use Taylor polynomials to approximate this function, to obtain a way to compute it with accuracy and simplicity.



# Approximating function $\text{SF}(x)$

As an example, begin with the degree 3 Taylor approximation to  $\sin t$ , expanded about  $t = 0$ :

$$\sin t = t - \frac{1}{6}t^3 + \frac{1}{120}t^5 \cos \zeta, \quad 0 \leq \zeta \leq t$$

# Approximating function $\text{SF}(x)$

As an example, begin with the degree 3 Taylor approximation to  $\sin t$ , expanded about  $t = 0$ :

$$\sin t = t - \frac{1}{6}t^3 + \frac{1}{120}t^5 \cos \zeta, \quad 0 \leq \zeta \leq t$$

$$\frac{\sin t}{t} = 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta$$

# Approximating function $SF(x)$

As an example, begin with the degree 3 Taylor approximation to  $\sin t$ , expanded about  $t = 0$ :

$$\sin t = t - \frac{1}{6}t^3 + \frac{1}{120}t^5 \cos \zeta, \quad 0 \leq \zeta \leq t$$

$$\frac{\sin t}{t} = 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta$$

$$\int_0^x \frac{\sin t}{t} dt = \int_0^x \left( 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta \right) dt$$

# Approximating function $\text{SF}(x)$

As an example, begin with the degree 3 Taylor approximation to  $\sin t$ , expanded about  $t = 0$ :

$$\sin t = t - \frac{1}{6}t^3 + \frac{1}{120}t^5 \cos \zeta, \quad 0 \leq \zeta \leq t$$

$$\frac{\sin t}{t} = 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta$$

$$\int_0^x \frac{\sin t}{t} dt = \int_0^x \left( 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta \right) dt$$

$$\int_0^x \frac{\sin t}{t} dt = \int_0^x 1 dt - \frac{1}{6} \int_0^x t^2 dt + \frac{1}{120} \int_0^x t^4 \cos \zeta dt$$

# Approximating function SF(x)

As an example, begin with the degree 3 Taylor approximation to  $\sin t$ , expanded about  $t = 0$ :

$$\sin t = t - \frac{1}{6}t^3 + \frac{1}{120}t^5 \cos \zeta, \quad 0 \leq \zeta \leq t$$

$$\frac{\sin t}{t} = 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta$$

$$\int_0^x \frac{\sin t}{t} dt = \int_0^x \left( 1 - \frac{1}{6}t^2 + \frac{1}{120}t^4 \cos \zeta \right) dt$$

$$\int_0^x \frac{\sin t}{t} dt = \int_0^x 1 dt - \frac{1}{6} \int_0^x t^2 dt + \frac{1}{120} \int_0^x t^4 \cos \zeta dt$$

$$\int_0^x \frac{\sin t}{t} dt = x - \frac{1}{18}x^3 + \frac{1}{120} \int_0^x t^4 \cos \zeta dt$$

# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt$$

# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt = 1 - \frac{1}{18}x^2 + R_2(x),$$



# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt = 1 - \frac{1}{18}x^2 + R_2(x),$$

where the remainder (error term) is given by

$$R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \zeta dt$$

# Approximating function $SF(x)$

$$SF(x) = \frac{1}{x} \int_0^x \frac{\sin t}{t} dt = 1 - \frac{1}{18}x^2 + R_2(x),$$

where the remainder (error term) is given by

$$R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \zeta dt$$

How large is the error in approximation

$$SF(x) \approx 1 - \frac{1}{18}x^2$$

on interval  $x \in [-1, 1]$ ?

# Approximating function $SF(x)$

For  $x > 0$  we have  $0 \leq \cos \tilde{\zeta} \leq 1$ ,

# Approximating function $SF(x)$

For  $x > 0$  we have  $0 \leq \cos \zeta \leq 1$ , therefore

$$0 \leq R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \zeta dt \leq \frac{1}{120} \frac{1}{x} \int_0^x t^4 dt = \frac{1}{600} x^4$$

# Approximating function $SF(x)$

For  $x > 0$  we have  $0 \leq \cos \xi \leq 1$ , therefore

$$0 \leq R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \xi dt \leq \frac{1}{120} \frac{1}{x} \int_0^x t^4 dt = \frac{1}{600} x^4$$

and the same result can be shown for  $x < 0$ .

# Approximating function $SF(x)$

For  $x > 0$  we have  $0 \leq \cos \zeta \leq 1$ , therefore

$$0 \leq R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \zeta dt \leq \frac{1}{120} \frac{1}{x} \int_0^x t^4 dt = \frac{1}{600} x^4$$

and the same result can be shown for  $x < 0$ .

Then for  $|x| \leq 1$ , we have

$$0 \leq R_2(x) \leq \frac{1}{600}$$

# Approximating function $\text{SF}(x)$

For  $x > 0$  we have  $0 \leq \cos \zeta \leq 1$ , therefore

$$0 \leq R_2(x) = \frac{1}{120} \frac{1}{x} \int_0^x t^4 \cos \zeta dt \leq \frac{1}{120} \frac{1}{x} \int_0^x t^4 dt = \frac{1}{600} x^4$$

and the same result can be shown for  $x < 0$ .

Then for  $|x| \leq 1$ , we have

$$0 \leq R_2(x) \leq \frac{1}{600}$$

To obtain a more accurate approximation, we can proceed exactly as above, but simply use a higher degree approximation to  $\sin t$ .