
a
GLASS: recurrence interval
one pair per patient (n = 173); Spearman rho = 0.31, p = 2.8e−05
Months to first recurrence
150
100
50
0
−1
0
1
2
Primary tumor WEBS
b
GLASS longitudinal pairs
179 pairs from 148 patients; paired Wilcoxon p = 0.09
WEBS
2
1
0
−1
Primary
Recurrence
Primary
Recurrence
Trend line (LOESS) is a visual aid; the test is Spearman correlation
c
Top WEBS−sensitizing single agents
GDSC2, FDR q < 0.1; negative rho = higher sensitivity
Piperlongumine
CHIR−99021
SL0101
Nelarabine
IOX2
Avagacestat
Nutlin−3a (−)
AZD5991
THR−103
GSK2830371A
−0.3
−0.2
−0.1
0.0
Spearman rho (WEBS vs ln IC50)
d
DepMap dependency (glioma lines)
Negative = stronger dependency in high−WEBS lines
HDAC2
SIRT2
SF3B1
SRSF1
MED1
n.s.
n.s.
nom.
nom.
FDR
−0.2
−0.1
0.0
Correlation with dependency (Chronos)
FDR q < 0.05
n.s.
nominal p < 0.05
Single−agent associations; combination predictions are separate