

a
Spliceosome acetylation (OR = 7.36; corrected 2.29)
HNRNPM−K239k
DDX39B−K188k
PRPF8−K43k
HNRNPK−K163k
HNRNPA1−K350k
HNRNPA2B1−K112k
HNRNPK−K60k
PRPF38B−K521k
HNRNPA1−K166k
HNRNPA2B1−K17k
DHX15−K17k
ALYREF−K93k
SNRNP200−K55k
DHX16−K210k
HNRNPM−K698k
HNRNPA2B1−K104k
HNRNPM−K672k
SNU13−K21k
HNRNPA2B1−K59k
DDX46−K776k
SNU13−K20k
SNRNP70−K162k
PRPF40B−K170k
PRPF40A−K169k
PRPF8−K36k
THOC5−K24k
SRSF11−K466k
HNRNPA2B1−K113k
HNRNPC−K243k
HNRNPA2B1−K46k
Acetyl.
2
1
0
−1
−2
WEBS group
LDTS
LHS
LLS
b
Phosphosites: 198/319 sig (corrected 124/319)
−log10(p)
15
10
5
0
−2
0
2
logFC (high− vs low−WEBS)
NS
Spliceosome NS
Spliceosome sig (up)
Other sig
Spliceosome sig (down)
c
Splicing−kinase abundance vs WEBS
CDK9
CDK13
CDK12
CLK3
CDK7
CLK1
SRPK1
CLK2
SRPK2
CLK4
0.0
0.2
0.4
Spearman rho vs WEBS
n.s.
p < 0.05
d
GENIE3 connectivity sensitivity
KAT2A
SIRT6
SIRT7
HDAC3
KAT5
CREBBP
EP300
KAT8
HDAC2
SIRT3
KAT2B
SIRT1
HDAC1
SIRT2
0.0
0.5
1.0
1.5
Relative |shift| (spliceosome / other)
Erasers
Writers