

Table 2: Pattern of results in follow up biopsies- comparative review of literature

Study	Cytology method used in the study	Total number of cases studied	Number of cases with biopsy				High grade dysplasia (CIN2 & above) on biopsy			
			LSIL	LSIL-H	ASC-H	HSIL	LSIL % (n)	LSIL-H % (n)	ASC-H % (n)	HSIL % (n)
1 Current study [11]	SurePath™ [¶]	77,979 LSIL-H 146 (0.19%) (12 months)	557	88	38	109	10% (55)	33% (29)	31% (12)	69% (75)
2 Nasser et al 2003 [1]	Not stated	Not stated (12 months)	150	144	X	X	15% (23)	29% (42)	X	X
3 Kir et al 2004 [2]	Not stated	21,342 (2 year)	27	13	X	X	11% (3)	61% (8)	X	X
4 McGrath et al 2000 [12]	Not stated	48,687 LSIL- 108 (0.2%) (14.5 months)	X	58	X	X	X	59% (34)	X	X
5 Elsheikh et al 2006 [14]	ThinPrep®*	129,911 LSIL- 194 (0.15%) (25 months)	575	59	110	289	13% (75)	41% (24)	45% (49)	74% (214)
6 Booth et al 2005 [13] (Abstract)	Not stated	21,082 LSIL-H 31 (0.15%)	X	X	X	X	10% (5)	45% (9)	X	69% (29)
7 D'Furio et al 2005 [17] (Abstract)	ThinPrep®*	Not stated	X	83	37	X	X	40%	62%	X
8 Underwood et al 2006 [15] (Abstract)	ThinPrep®&Conventional	130,761 A. ThinPrep®* (127,929) B. Conventional (2832) LSIL-H 270 (0.2%) (24 months)	X	X	X	X	13% (163)	36% (70)	38% (93)	66% (170)
9 O'Brien et al 2006 [16] (Abstract)	ThinPrep®*	A. Pre-imaging: 76,365 (50 LSIL-H 0.065%) B. Post-imaging: 63,812 (139 LSIL-H 0.22%)	X	A- 40 B- 107	X	X	X	A- 23%(9) B- 37%(39)	X	X
10 Jain et al 2005 [18] (Abstract)	Not stated	Total 67 LSIL-H A- Few : 3 or more ASC-H cells B- Rare: 1 to 2 ASC-H cells	X	48 A. 22 B. 26	X	X	X	A- 64% (14/22) B- 23% (6/26)	X	56% (9/16)

§This study used different terminology as mild to moderate dysplasia but implied non-definitive interpretation equivalent to LSIL-H; ¶SurePath™ (TriPath Imaging, Inc. Burlington, NC, USA)), * ThinPrep® (Cytoc Corporation, Marlborough, MA, USA), m, months; yrs, years. X Blanks represent lack of that information in the corresponding published data.

from HSIL, justifies LSIL-H as a separate group for optimal clinical management (Figure 5) with possible application of molecular events such as p16 in the future.

List of abbreviations

ASC-H, Atypical squamous cells Cannot exclude high-grade intraepithelial lesion; ASCCP, American Society for Colposcopy and Cervical Pathology; ASCUS, Atypical squamous cells of undetermined significance; Bethesda 2001, 2001 Bethesda System terminology; CIN, cervical intraepithelial neoplasia; HPV, human papilloma virus; HPV-T, HPV DNA testing; HSIL, high-grade squamous intraepithelial lesion; LSIL, low-grade squamous intraepithelial lesion; LSIL-H, low-grade squamous intraepithelial lesion, cannot exclude high-grade intraepithelial lesion; ND, negative for dysplasia; PPV, Positive predictive value; SAPK cells, small atypical parakeratotic cells.

Competing interests

The author(s) declare that they have no competing interests.

Authors' contributions

VS, Conceptual organization as senior author, cytological-histological evaluation, data analysis, executing IRB process, and writing the manuscript.

NK, Cytopathology fellow, organized IRB process, collected all the data, and review of manuscript.

RN & GB: Data analysis, review of manuscript, and analyze management algorithm.

Acknowledgements

This study was presented in part at the 53rd Annual Scientific Meeting, American Society of Cytopathology, 4–9 November 2005, San Diego, CA and was recipient of 2005 Warren R. Lang Resident Physician Award for the best scientific paper in cytology.

Authors thank the contribution by Drs. Cafaro, Parameswaran, Basir, Behmaram, Chavan, Rao, Markelova, Keating, and all our cytotechnologists for routine initial review of cases in this study. Our thanks to Maria Halaska for assisting in data retrieval. We also thank Dr. Laud, Professor of Biostatistics for evaluation and explanation of statistical analysis.