

Original research

Open Access

A simple and rapid method for detection of *Trypanosoma evansi* in the dromedary camel using a nested polymerase chain reaction

Imadeldin E Aradaib*^{1,2} and Ali A Majid³

Address: ¹Molecular Biology Research Unit, National Ribat University, Khartoum, Sudan, ²Molecular Biology laboratory (MBL), Department of Medicine, Pharmacology and Toxicology, Faculty of Veterinary Medicine, University of Khartoum, P. O. Box 32, Khartoum North, Sudan and ³National Council for Research, Khartoum, Republic of Sudan

Email: Imadeldin E Aradaib* - aradaib@yahoo.com; Ali A Majid - ali.majid@gmail.com

* Corresponding author

Published: 20 May 2006

Received: 04 November 2005

Kinetoplastid Biology and Disease 2006, 5:2 doi:10.1186/1475-9292-5-2

Accepted: 20 May 2006

This article is available from: <http://www.kinetoplastids.com/content/5/1/2>

© 2006 Aradaib and Majid; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

A nested polymerase chain reaction (nPCR)-based assay, was developed and evaluated for rapid detection of *Trypanosoma evansi* in experimentally infected mice and naturally infected camels (*Camelus dromedarius*). Four oligonucleotide primers (TE1, TE2, TE3 and TE4), selected from nuclear repetitive gene of *T. evansi*, were designed and used for PCR amplifications. The first amplification, using a pair of outer primers TE1 and TE2, produced a 821-bp primary PCR product from *T. evansi* DNA. The second amplification, using nested (internal) pair of primers TE3 and TE4, produced a 270-bp PCR product. *T. evansi* DNAs extracted from blood samples of experimentally infected mice and naturally infected Sudanese breed of dromedary camels were detected by this nested PCR-based assay. The nested primers TE3 and TE4 increased the sensitivity of the PCR assay and as little as 10 fg of *T. evansi* DNA (equivalent to a single copy of the putative gene of the parasite) was amplified and visualized onto ethidium bromide-stained agarose gels.

Amplification products were not detected when the PCR-based assay was applied to DNA from other blood parasites including *Thieleria annulata*, *Babesia bigemina* or nucleic acid free samples. Application of this nPCR-based assay to clinical samples resulted in direct detection of *T. evansi* from a variety of tissue samples collected from experimentally infected mice and blood from naturally infected camels. The described nPCR-based assay provides a valuable tool to study the epidemiology of *T. evansi* infection in camels and other susceptible animal populations.

Background

Trypanosoma evansi (*T. evansi*), the cause of trypanosomiasis (Surra), constitutes one of the major veterinary problems worldwide. The disease causes significant morbidity and mortality in camels in the Sudan, which has a population of over 3 million camels [1]. trypanosomiasis in camels occurs both in chronic and acute forms [2]. The chronic form of the disease is most common and is likely to be associated with secondary infections due to immu-

nosuppression [3]. Clinical signs and pathological lesions caused by *T. evansi* in camels are unreliable for definitive diagnosis [4]. In addition, detection of parasites in the blood is difficult because parasitaemia is intermittent [5]. Serological tests have been developed and evaluated for diagnosis of trypanosomiasis in camels. They include card agglutination test and enzyme-linked immunosorbent assay (Ab-ELISA) [6-8]. In general serological techniques are useful for detection of a past infection but not for