

Perspective

ADAPT: The Wrong Way to Stop a Clinical Trial

Steven E. Nissen

In an accompanying article published in *PLoS Clinical Trials*, the Alzheimer's Disease Anti-Inflammatory Prevention Trial (ADAPT) Research Group report the cardiovascular outcomes from their study [1]. The circumstances surrounding the termination of ADAPT were unusual and provide an important lesson for all clinical trialists, demonstrating the importance of following rigorous procedures for prematurely stopping clinical trials. In this case, stopping the trial before its intended completion resulted in data that cannot be reliably interpreted.

On September 30, 2004, Merck withdrew rofecoxib (Vioxx) from the market after the trial's data safety and monitoring board (DSMB) recommended termination of a placebo-controlled study of this agent in the prevention of colon polyps [2]. The reason for study termination was a statistically significant increase in adverse cardiovascular outcomes. On December 17, 2004, Pfizer announced that termination of a trial of celecoxib (Celebrex) in colon polyp prevention, because it also showed statistically significant evidence for increased cardiovascular event rates [3]. The results of these two trials were subsequently published in February 2005 in the *New England Journal of Medicine* [4,5]. The revelations about increased cardiovascular events with these "coxibs" generated enormous public attention and concern. However, in both cases, the decision-making leading to trial discontinuation was handled appropriately through the regular reviews conducted by an independent DSMB.

Three days following the announcement of the termination of the celecoxib colon polyp prevention study, the National Institute of Health (NIH), issued a press release entitled "Use of Non-Steroid Anti-Inflammatory Drugs Suspended in Large Alzheimer's Disease Prevention Trial" [6]. The NIH press release stated

that "data from the ADAPT trial indicated an apparent increase in cardiovascular and cerebrovascular events among participants taking naproxen when compared to placebo." In the press release, NIH Director Dr. Elias Zerhouni stated that "this step is being taken as a precautionary measure to ensure the safety of the study's participants" and that "the investigators made their decision based on the risk/benefit analysis specific to this trial." Shortly following the NIH announcement, the Food and Drug Administration (FDA) issued a public statement that "based on emerging information from a long-term prevention trial, the risk of cardiovascular and cerebrovascular events may increase among patients taking naproxen" [7].

These announcements generated front-page headlines such as "Heart Risk Seen in Naproxen" (*Wall Street Journal*), "Tough Choice: Pain or Risk" (*USA Today*), and "Patients, Doctors Agonize over Risks of Painkiller" (*Los Angeles Times*) [8]. Occurring immediately following the revelations about rofecoxib and celecoxib, the naproxen announcement generated considerable public apprehension [8]. Physicians received many urgent calls from worried patients. However, there was a major problem with the naproxen warning: it was not based upon the application of standard procedures for stopping an ongoing clinical trial. During the subsequent FDA hearings to set policy on nonsteroidal anti-inflammatory agents (NSAIDs) and COX-2 inhibitors, I described the warnings about naproxen as "the medical equivalent of yelling 'fire' in a crowded auditorium" [9].

The Data Finally Become Available

Now, nearly two years after the closure of the ADAPT study, we finally get to see the data that resulted in the public warning about naproxen [1]. For the standard composite endpoint of cardiovascular death, myocardial infarction, and stroke, there were 17 events in celecoxib treatment group, 23 in the naproxen arm, and 22 in the placebo. The hazard ratio for naproxen compared with placebo was

1.57 with 95% confidence intervals of 0.87 to 2.81, $p = 0.13$. A broader composite outcome that added heart failure and transient ischemic attack yielded a marginally significant p value when comparing naproxen with placebo.

We must ask whether a DSMB would stop an ongoing clinical trial for such findings. The answer, it appears, was that the DSMB did not stop the trial; NIH officials did [8]. There was no regularly scheduled DSMB safety review that resulted in study termination. It appears that NIH officials, concerned about revelations regarding the safety of coxibs in the colon polyps studies, simply decided to look at the cardiovascular event results in ADAPT [8,10]. Seeing a marginally significant difference between naproxen and placebo, they abruptly stopped the trial. Indeed, NIH officials went even further and issued the public warning [6].

Both the termination of this trial and the issuance of a public warning about naproxen were inappropriate and reflected faulty logic. These actions unnecessarily created public consternation

Funding: The author received no specific funding for this article.

Competing Interests: SEN is the principal investigator for a large clinical trial comparing cardiovascular event rates for three nonsteroidal anti-inflammatory agents: naproxen, ibuprofen, and celecoxib. Although the study is supported by Pfizer, SEN and the Executive Committee for the trial have stipulated that no member of the trial leadership may accept any compensation from the sponsor or any other maker of drugs in this class.

Citation: Nissen SE (2006) ADAPT: The wrong way to stop a clinical trial. *PLoS Clin Trials* 1(7): e35. doi:10.1371/journal.pctr.0010035

Copyright: © 2006 Steven E. Nissen. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abbreviations: ADAPT, Alzheimer's Disease Anti-Inflammatory Prevention Trial; DSMB, data safety and monitoring board; FDA, Food and Drug Administration; NIH, National Institutes of Health; NSAID, nonsteroidal anti-inflammatory agent. Steven Nissen is chairman of the Department of Cardiovascular Medicine at the Cleveland Clinic Foundation, Cleveland, Ohio, United States. E-mail: nissens@ccf.org

The Perspectives section is for experts to discuss the clinical practice or public health implications of a published study that is freely available online.