

Research

Open Access

Use of pIVEX plasmids for protein overproduction in *Escherichia coli*

Julie Rogé and Jean-Michel Betton*

Address: Unité de Repliement et Modélisation des Protéines Institut Pasteur CNRS-URA2185 28, rue du Docteur Roux 75724 Paris Cedex 15, France

Email: Julie Rogé - jroge.abag@wanadoo.fr; Jean-Michel Betton* - jmbetton@pasteur.fr

* Corresponding author

Published: 02 June 2005

Received: 21 April 2005

Microbial Cell Factories 2005, 4:18 doi:10.1186/1475-2859-4-18

Accepted: 02 June 2005

This article is available from: <http://www.microbialcellfactories.com/content/4/1/18>

© 2005 Rogé and Betton; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: The pIVEX plasmids are vectors optimized for expression in the Rapid Translation System (RTS) cell-free system under control of bacteriophage T7 transcription elements. Even if these plasmids are intended for use *in vitro*, it is usually worthwhile to compare both cell-free and bacterial expression from the same genetic construct. However, some RTS users encountered problems when they introduced these plasmids into *Escherichia coli* host strains producing the T7 RNA polymerase.

Results: We verified that difficulties in transforming the commonly used BL21(λDE3) strain with pIVEX arose from the presence of a strong T7 promoter combined with a high-copy number plasmid, independent of gene expression. When these vectors were introduced into this strain harboring a compatible plasmid carrying the lactose repressor (*lacI*), we improved the transformation efficiency by 4 orders of magnitude. Moreover, we designed a transformation protocol that allows, after induction, the overproduction of pIVEX-encoded proteins in the BL21(λDE3) strain.

Conclusion: Using the correct plasmid/host combination and transformation-expression protocol, we could directly compare overproduction of the same pIVEX-encoded proteins from both *in vivo* and *in vitro* expression systems.

Background

Recent developments in cell-free systems offer new and promising possibilities for producing recombinant proteins [1]. The RTS, commercialized by Roche, is an exchange cell-free system with improved productivity [2]. The continuous supply of consumable substrates and removal of reaction products provide a yield of several milligrams of protein. This system uses bacteriophage T7 RNA polymerase to perform transcription, while an enriched *E. coli* S30 extract provides the translational machinery. Thus, protein production in RTS requires a preliminary cloning step of the target gene into a vector,

downstream of the T7 promoter. For this purpose, the pIVEX family of expression plasmids has been optimized for *in vitro* use. They include a T7 promoter comprising the T7 gene 10 translation enhancer [3], an efficient prokaryotic Shine-Dalgarno sequence with an optimum distance to the start codon, a multiple cloning site, and a T7 terminator which prevents 3'-terminal exonucleolytic degradation of the mRNA. These vectors are very convenient since multiple cloning sites were designed to allow gene fusions with several tags either at the N- or C-terminal of target proteins by conserving the same restriction strategy and reading frame compatibility.