

Research

Open Access

Moderate exercise and chronic stress produce counteractive effects on different areas of the brain by acting through various neurotransmitter receptor subtypes: A hypothesis

Suptendra N Sarbadhikari^{*1} and Asit K Saha²

Address: ¹TIFAC-CORE in Biomedical Technology, Amrita Vishwa Vidyapeetham, Amritapuri 690525, India and ²School of Electrical and Information Engineering, University of South Australia, Mawson Lakes Campus, South Australia 5095, Australia

Email: Suptendra N Sarbadhikari^{*} - supten@gmail.com; Asit K Saha - draycott7@yahoo.com.au

^{*} Corresponding author

Published: 23 September 2006

Received: 13 July 2006

Theoretical Biology and Medical Modelling 2006, **3**:33 doi:10.1186/1742-4682-3-33

Accepted: 23 September 2006

This article is available from: <http://www.tbiomed.com/content/3/1/33>

© 2006 Sarbadhikari and Saha; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Regular, "moderate", physical exercise is an established non-pharmacological form of treatment for depressive disorders. Brain lateralization has a significant role in the progress of depression. External stimuli such as various stressors or exercise influence the higher functions of the brain (cognition and affect). These effects often do not follow a linear course. Therefore, nonlinear dynamics seem best suited for modeling many of the phenomena, and putative global pathways in the brain, attributable to such external influences.

Hypothesis: The general hypothesis presented here considers only the nonlinear aspects of the effects produced by "moderate" exercise and "chronic" stressors, but does not preclude the possibility of linear responses. In reality, both linear and nonlinear mechanisms may be involved in the final outcomes. The well-known neurotransmitters serotonin (5-HT), dopamine (D) and norepinephrine (NE) all have various receptor subtypes. The article hypothesizes that 'Stress' increases the activity/concentration of some particular subtypes of receptors (designated nt_s) for each of the known (and unknown) neurotransmitters in the right anterior (RA) and left posterior (LP) regions (cortical and subcortical) of the brain, and has the converse effects on a different set of receptor subtypes (designated nt_n). In contrast, 'Exercise' increases nt_n activity/concentration and/or reduces nt_s activity/concentration in the LA and RP areas of the brain. These effects may be initiated by the activation of Brain Derived Neurotrophic Factor (BDNF) (among others) in exercise and its suppression in stress.

Conclusion: On the basis of this hypothesis, a better understanding of brain neurodynamics might be achieved by considering the oscillations caused by single neurotransmitters acting on their different receptor subtypes, and the temporal pattern of recruitment of these subtypes. Further, appropriately designed and planned experiments will not only corroborate such theoretical models, but also shed more light on the underlying brain dynamics.

Background

Regular, "moderate", physical exercise is a non-pharmaco-

logical form of adjunctive treatment for depressive disorders. External stimuli such as various stressors or exercise