

Research article

Open Access

The p53-dependent apoptotic pathway of breast cancer cells (BC-M1) induced by the bis-type bio-reductive compound aziridinyl-naphthoquinone

Yu-Ping Yang¹, Hsien-Shou Kuo¹, Hsin-Da Tsai¹, Yi-Chen Peng¹ and Yuh-Ling Lin²¹Department of Biochemistry, College of Medicine, Taipei Medical University, Taiwan, ROC²Department of Medicine, College of Medicine, Fu-Jen Catholic University, Taipei Hsien, Taiwan, ROCCorresponding author: Yuh-Ling Lin, med0018@mails.fju.edu.tw

Received: 4 May 2004 Revisions requested: 27 May 2004 Revisions received: 4 Aug 2004 Accepted: 6 Sep 2004 Published: 04 Nov 2004

Breast Cancer Res 2005, **7**:R19-R27 (DOI 10.1186/bcr939)© 2004 Yang *et al.*, licensee BioMed Central Ltd.This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is cited.

Abstract

Introduction Several aziridinylbenzoquinone drugs have undergone clinical trials as potential antitumor drugs. These bio-reductive compounds are designed to kill cells preferentially within the hypoxia tumor microenvironment. The bio-reductive compound of bis-type naphthoquinone synthesized in our laboratory, 2-aziridin-1-yl-3-[(2-{2-[(3-aziridin-1-yl-1,4-dioxo-1,4-dihydronaphthalen-2-yl)thio]ethoxy}ethyl)thio]naphthoquinone (AZ-1), had the most potent death effect on the breast cancer cells BC-M1 in our previous screening. In the present study, we determined that the mechanism of the death effect of BC-M1 cells induced by AZ-1 was mediated by the apoptosis pathway.

Methods We evaluated the cytotoxicity of AZ-1 and the anti-breast cancer drugs tamoxifen and paclitaxel to BC-M1 cells and MCF-7 cells by the MTT assay and measured the apoptosis phenomena by Hoechst 33258 staining for apoptotic bodies. We also quantified the sub-G₁ peak area and the ratio of the CH₂/CH₃ peak area of the cell membrane in BC-M1 cells by flow cytometry and ¹H-NMR spectra, respectively. The apoptosis-related protein expressions, including p53, p21, the RNA-relating protein T-cell restricted intracellular antigen-related protein, cyclin-dependent kinase 2 (cell cycle regulating kinase) and pro-caspase 3, were detected by western blot, and the caspase-3 enzyme activity was also quantified by an assay kit.

Results AZ-1 induced two of the breast cancer cell lines, with IC₅₀ = 0.51 μM in BC-M1 cells and with IC₅₀ = 0.57 μM in MCF-7 cells, and showed less cytotoxicity to normal fibroblast cells (skin fibroblasts) with IC₅₀ = 5.6 μM. There was a 10-fold difference between two breast cancer cell lines and normal fibroblasts. Of the two anti-breast cancer drugs, tamoxifen showed IC₅₀ = 0.12 μM to BC-M1 cells and paclitaxel had much less sensitivity than AZ-1. The expression of p53 protein increased from 0.5 to 1.0 μM AZ-1 and decreased at 2.0 μM AZ-1. The p21 protein increased from 0.5 μM AZ-1, with the highest at 2 μM AZ-1. Regarding the AZ-1 compound-induced BC-M1 cells mediating the apoptosis pathway, the apoptotic body formation, the sub-G₁ peak area, the ratio of CH₂/CH₃ of phospholipids in the cell membrane and the enzyme activity of caspase-3 were all in direct proportion with the dose-dependent increase of the concentration of AZ-1. The death effect-related proteins, including T-cell restricted intracellular antigen-related protein, cyclin-dependent kinase 2, and pro-caspase-3, all dose-dependently decreased with AZ-1 concentration.

Conclusions The AZ-1-induced cell death of BC-M1 cells mediating the apoptosis pathway might be associated with p53 protein expression, and AZ-1 could have the chance to be a candidate drug for anti-breast cancer following more experimental evidence, such as animal models.

Keywords: apoptosis, bio-reductive compound, bis-type aziridinyl-naphthoquinone, breast cancer cells (BC-M1 and MCF-7)

Introduction

The bio-reductive drugs, aziridinylbenzoquinones, are a class of compounds designed to exploit one of the features of solid tumor biology caused by an inadequate blood sup-

ply to the solid tumor; namely, tumor hypoxia. Such regions generally are resistant to radiation and other oxygen-requiring treatment [1-4]. The ideal bio-reductive drug should be administered as an inactive prodrug that is only activated

AZ-1 = 2-aziridin-1-yl-3-[(2-{2-[(3-aziridin-1-yl-1,4-dioxo-1,4-dihydronaphthalen-2-yl)thio]ethoxy}ethyl)thio]naphthoquinone; cdk = cyclin-dependent kinase; DMEM = Dulbecco's modified Eagle's medium; IC₅₀ = inhibition concentration of 50% cell growth; NMR = nuclear magnetic resonance; PBS = phosphate-buffered saline; TIAR = T-cell restricted intracellular antigen-related protein.