
Clinical Study Protocol — ONC-450

Study Title: A Phase 3, Randomized, Double-Blind, Placebo-Controlled Study of Zelparib as Maintenance Therapy in Adults with Advanced Ovarian Cancer

Sponsor: Aruna Therapeutics, Inc. | **Protocol Version:** 4.0

This is a fully synthetic document created for demonstration purposes.

1. Objectives

To evaluate the efficacy of Zelparib 300 mg twice daily versus placebo on progression-free survival in adults with advanced high-grade serous ovarian cancer who have responded to first-line platinum-based chemotherapy. Zelparib is an investigational PARP inhibitor; it has not been approved by the U.S. Food and Drug Administration (FDA).

2. Study Design

Randomized, double-blind, placebo-controlled, parallel-group trial at 42 sites. Approximately **420 participants** will be randomized **2:1** to Zelparib or matching placebo as maintenance therapy. Treatment continues until disease progression or unacceptable toxicity; a safety follow-up occurs 30 days after the last dose.

3. Study Procedures

Participants attend clinic visits every 4 weeks. Each visit includes vital signs, physical examination, a symptom review, and a venous blood draw of approximately 20 mL for hematology and safety chemistry. Imaging (CT) is performed every 12 weeks to assess disease status.

4. Participant Responsibilities

Participants must take the study drug twice daily, complete the symptom diary, attend scheduled visits and imaging, and report new symptoms promptly to the study team.

5. Risks and Safety Considerations

In earlier studies, Zelparib caused **myelosuppression**: anemia (Grade 3 or higher in ~18% of participants), neutropenia (~9%), and thrombocytopenia (~6%), managed by dose interruption and transfusion. **Complete blood counts are monitored at every visit.** A small number of participants developed **myelodysplastic syndrome / acute myeloid leukemia (MDS/AML), a serious and**

potentially fatal condition. Common adverse events include nausea (fatigue), fatigue, and decreased appetite.

6. Benefits

Participants may experience a longer period without cancer progression; benefit is not guaranteed, and one-third of participants receive placebo. The information learned may help future patients with ovarian cancer.

7. Alternative Treatments

Alternatives to participation include routine surveillance (observation) after chemotherapy, or other approved maintenance therapies where eligible. Your oncologist can discuss these options and their benefits and risks.

8. Compensation for Injury

The sponsor maintains clinical trial insurance. For a trial-related injury, **the sponsor will pay the reasonable costs of medical treatment**, in accordance with the site's compensation policy.

9. Confidentiality and Record Access

Records identifying participants are kept confidential. Authorized monitors, auditors, the IRB, and the **FDA may inspect original medical records** to verify study conduct.

10. Contacts

Participants may contact the principal investigator for study questions, the IRB for questions about their rights, and a 24-hour line for trial-related injury.

11. Voluntary Participation

Participation is voluntary; participants may decline or withdraw at any time without penalty or loss of benefits to which they are otherwise entitled.