

Arithmetic Structure of Baryons and the Bost–Connes System

ShaoWu Hu

Independent Researcher, Shanghai, China

forest@xindong.com ORCID: 0009-0009-8300-4691

Abstract

Of the proton mass 938 MeV, the three valence quarks contribute only about 1%; the remaining 99% non-constituent mass, the absolute stability of the proton ($\tau_p > 10^{34}$ yr), color confinement, and the Regge regularity of the baryon excitation spectrum $N(939)$ – $\Lambda(1800)$ have long lacked a unified analytic explanation. Companion papers [19, 20] start from the finite Dirac operator layer D_F of the Bost–Connes (BC) spectral triple and provide an arithmetic description of the twelve fermion masses and the three gauge forces of the Standard Model, accounting for $\sim 1\%$ of the proton mass. The present work proves that all of the remaining phenomena listed above are uniquely described by the other layer of the BC spectral triple—the Hamiltonian layer $H = \log n$. The KMS partition function of this layer is identically the Riemann ζ function; its unique simple pole at $\beta = 1$ corresponds to a density-of-states divergence along the trivial character direction χ_0 , and this divergence is the arithmetic origin of color confinement and of the collective emergence of baryons. **Main Conjecture:** the emergent states at color-singlet levels $n = 3, 4, 5, \dots$ along χ_0 are precisely the baryons of the Standard Model, with masses given by the parameter-free formula

$$E_n = \sqrt{\zeta(\beta) \log^2 \zeta(\beta) / C^*}, \quad \beta = 1 + 1/\sqrt{n},$$

where $C^* = 1.954$ is the curvature constant of $\mathbb{N}^+$. For $n = 3$ we obtain $E_3 = 937$ MeV (deviation -0.2%); the range $n = 3, \dots, 10$ covers up to $\Lambda(1800)$ with all deviations $< 2\%$. Moreover: the neutron–proton mass difference 1.29 MeV (-0.2%), the magnetic moment ratio $\mu_p/\mu_n = -1.459$ (0.06%), the axial-vector coupling $g_A = 1.311$ ($+2.8\%$), the neutron lifetime $\tau_n = 930$ s (beam method 887.7, $+4.8\%$), the pion mass 137 MeV (-2.2%), and the nuclear force coupling constant $\alpha_{\text{nuclear}} = 14.1$ ($+0.8\%$).

Contents

1	Introduction	3
2	The Hamiltonian Layer	4
2.1	Quantum Mechanics of the Positive Integers and the ζ Partition Function	4
2.2	The ζ Pole, Color Confinement and Collective Emergence	4
2.3	Emergent States and Decoupling from the Three Gauge Forces	5
2.4	The Energy Scale of Emergent States	6
3	Properties of the Emergent States	7
3.1	Color Singlet	7
3.2	Ground-State Mass	8
3.3	Non-constituent Mass	8
3.4	Absolute Stability	9
3.5	Spin Classification	9
3.6	Excitation Spectrum	10
3.7	Summary of the Six Properties and Comparison with Experiment	11

4	Two-layer Structure of the Spectral Triple	12
4.1	Mathematical Basis of the Two-layer Structure: Tensor Product and Character Orthogonality	13
4.2	Mechanism of Quantum Number Transfer from Fermions to Baryons	14
4.3	Decomposition of the Spectral Action	14
4.4	Color Singlet Constraint vs. Character Projection	15
4.5	Summary Table of the Two-layer Structure	15
4.6	99% + 1%: Internal Decomposition, Not External Addition	15
4.7	Joint Observables	16
4.8	Fine Structure of the Color-Singlet Level	16
5	The Neutron–Proton Mass Difference	18
5.1	NO-GO Theorem: Holomorphic Functions Cannot Produce Flavor Splitting . . .	18
5.2	The Chern–Simons Phase Bypasses NO-GO	18
5.3	Two-layer Mass Difference Formula	19
6	Joint Two-layer Predictions	19
6.1	Magnetic Moment Ratio μ_p/μ_n	20
6.2	Axial-vector Coupling g_A	21
6.3	Neutron Lifetime τ_n	21
7	Mesons and the Nuclear Force	22
7.1	Pion Mass	22
7.2	Nuclear Force	23
7.2.1	Range	23
7.2.2	Coupling Constant	23
8	Discussion	24
8.1	Summary of Results	24
8.2	Open Questions	25
8.3	Concluding Remarks	25

1 Introduction

Protons and neutrons make up all of the visible matter in the universe, yet the Standard Model leaves several fundamental questions about their description unanswered. The combined mass of two up quarks and one down quark [25, 26] is about 9 MeV [1], while the proton mass is 938 MeV. The remaining 99% non-constituent mass is attributed in principle to the QCD vacuum energy and gluon-field contributions [27, 28, 2]; lattice QCD [29] can numerically reproduce it [3], but there is no analytic answer to why this value is precisely 938. The proton lifetime lower bound $\tau_p > 2.4 \times 10^{34}$ yr [5] is 24 orders of magnitude larger than the age of the universe, and the proton decays predicted by grand unified theories [10, 9] remain experimentally unconfirmed. Color confinement—the fact that quarks have never been observed as isolated states—is a non-perturbative phenomenon in the strong coupling regime of QCD; a rigorous mathematical proof is still lacking, and it is one of the Clay Millennium Prize Problems [34, 6]. Baryon excitations extend from $N(939)$ up to $N(2600)$ and exhibit clear Regge trajectories [35, 7, 8], suggesting a string-like intrinsic algebraic structure. The neutron–proton mass difference 1.293 MeV [1] amounts to only 0.14% of the proton mass; this small splitting determines that the universe is hydrogen-dominated, yet its analytic origin remains open [4].

Our companion papers have already provided a partial picture of the Standard Model on the finite Dirac operator layer D_F of the BC spectral triple [19, 20]. D_F acts on the dual group of $(\mathbb{Z}/7\mathbb{Z})^*$; the commutator $[D_F, T_a]$ along character directions yields the three gauge forces, and characters together with Dirichlet L -functions yield the identities and masses of the 12 fermions. This layer accounts for the internal quantum numbers of the proton (flavor, charge, spin) but cannot reach non-constituent mass, color confinement, or the baryon spectrum.

This paper turns to the other layer of the BC spectral triple—the integer quantization layer in which the Hamiltonian $H = \log n$ acts on $\ell^2(\mathbb{N}^*)$ [12]. The full Dirac operator is written as

$$D = H \otimes I_F + \gamma \otimes D_F, \quad (1)$$

where the two terms on the right-hand side carry distinct arithmetic content. We refer to the layer in which $H = \log n$ lives as the **Hamiltonian layer (Layer 1)**, and the layer of D_F as the **finite Dirac operator layer (Layer 2)**. We prove that the KMS partition function of Layer 1 is identically the Riemann ζ function, and that the unique simple pole of $\zeta(\beta)$ at $\beta = 1$ corresponds to a density-of-states divergence along the trivial character direction χ_0 . This divergence simultaneously characterizes color confinement (divergence only along χ_0 , finite along non-trivial directions) and the collective emergence of baryons. The mass of these emergent states is given by a parameter-free formula:

Main Conjecture. The collective emergent states of the BC system at color-singlet levels $n \in \mathbb{N}_{\geq 3}$ along χ_0 are precisely the baryons of the Standard Model. Their masses are

$$E_n = \sqrt{\zeta(\beta) \log^2 \zeta(\beta) / C^*}, \quad \beta = 1 + 1/\sqrt{n},$$

where $C^* = 1.954$ is the curvature constant of $\mathbb{N}^+$. For the derivation of the formula see [21].

The paper is organized as follows. Section 2 establishes the arithmetic skeleton of Layer 1: the spectrum of H , the ζ partition function, the pole along χ_0 under the Artin decomposition, the decoupling from Layer 2 in this direction, and the energy formula for emergent states. Section 3 rigorously deduces from these mathematical facts the six properties that the emergent states must possess—color singlet structure, mass, non-constituent nature, stability, spin classification, and excitation spectrum—and Section 3.7 unifies the comparison with experiment. Sections 4–7 derive, under the Main Conjecture, further testable predictions: the neutron–proton mass difference, the magnetic moment ratio, the axial-vector coupling, the neutron lifetime, and the pion mass. Section 8 discusses open questions.

2 The Hamiltonian Layer

2.1 Quantum Mechanics of the Positive Integers and the ζ Partition Function

The Hilbert space of the BC system, $\ell^2(\mathbb{N}^*)$, has the positive integers as an orthonormal basis, and the Hamiltonian acts as [12, 22, 23]

$$H|n\rangle = (\log n) |n\rangle. \quad (2)$$

Each prime p contributes the eigenvalue $\log p$ as a fundamental excitation; the energy of a composite integer is an integer linear combination of these fundamental excitations. The partition function $Z(\beta) = \text{Tr } e^{-\beta H}$ gives a mathematical identity on Layer 1:

Proposition 2.1 (KMS partition function equals ζ). *For $\text{Re } \beta > 1$,*

$$Z(\beta) = \text{Tr } e^{-\beta H} = \sum_{n=1}^{\infty} e^{-\beta \log n} = \sum_{n=1}^{\infty} n^{-\beta} = \zeta(\beta), \quad (3)$$

and ζ simultaneously admits the Euler product [15]

$$\zeta(\beta) = \prod_{p \text{ prime}} \frac{1}{1 - p^{-\beta}}. \quad (4)$$

The Euler product (4) implies that the partition function is a *product over all primes*: any physical quantity it describes cannot be localized at a single prime, but rather reflects the collective behavior of all primes. This non-localizability will translate into the arithmetic statement of non-constituent mass in Section 3.3.

2.2 The ζ Pole, Color Confinement and Collective Emergence

Quarks have never been observed as isolated states; in nature only color-singlet bound states appear, while colored states—single quarks or diquarks—cannot exist independently. The companion paper [20] proves color confinement at the C^* -algebra level: the complex characters of $(\mathbb{Z}/7\mathbb{Z})^*$ (χ_1, χ_5 , of order 6) do not admit self-adjoint observables, so colored states are not observable in the algebraic sense (Color Non-observability Theorem, [20] Theorem 4.5). This section provides a complementary thermodynamic argument: the comparison of the analytic behavior of ζ and the Dirichlet L -functions at $\beta = 1$ directly shows that colored states not only are not observable, but also cannot emerge thermodynamically as independent particles.

$\zeta(\beta)$ is holomorphic on $\text{Re } \beta > 1$, and has a unique simple pole at $\beta = 1$ with residue 1:

$$\zeta(\beta) = \frac{1}{\beta - 1} + \gamma + O(\beta - 1) \quad (\beta \rightarrow 1^+), \quad (5)$$

where γ is the Euler–Mascheroni constant. In statistical mechanics, the divergence of the partition function corresponds to a phase transition: the density of states diverges, infinitely many modes are excited simultaneously, and the system transitions from a single-particle description to a collective one (Bose–Einstein condensation and the Hagedorn limit are of this kind). The BC system therefore must exhibit collective emergence along some direction in Layer 1. Which direction? We place ζ in the algebraic environment carved out by $(\mathbb{Z}/7\mathbb{Z})^*$.

For $p = 7$, the Dedekind ζ function of the cyclotomic field $\mathbb{Q}(\zeta_7)$ admits the Artin decomposition [14]

$$\zeta_{\mathbb{Q}(\zeta_7)}(\beta) = \underbrace{\zeta(\beta)}_{\text{trivial character } \chi_0} \cdot \underbrace{\prod_{j=1}^5 L(\beta, \chi_j)}_{\text{five non-trivial characters}}. \quad (6)$$

The χ_0 direction contributes the ζ factor, while $\chi_1, \dots, \chi_5$ contribute Dirichlet L -function factors. By Dirichlet’s theorem [13], $L(1, \chi_j)$ takes finite non-zero values for all $j \neq 0$. Hence every divergence of (6) at $\beta = 1$ is carried by the $\zeta(\beta)$ factor:

Corollary 2.2 (Divergence strictly confined to χ_0). *In the decomposition (6), the pole at $\beta = 1$ is given entirely by $\zeta(\beta)$; all factors $L(\beta, \chi_j)$ ($j \neq 0$) are regular and finite. The density-of-states divergence of the BC system on Layer 1 is therefore strictly confined to the trivial character direction χ_0 ; along any non-trivial character direction the density of states is finite, and no collective emergence occurs.*

χ_0 is the identity character on $(\mathbb{Z}/7\mathbb{Z})^*$, indistinguishable in both color and flavor, and corresponds to the “color singlet direction”; the five non-trivial characters correspond to “colored” and “flavored” directions. Corollary 2.2 is therefore the arithmetic statement of color confinement: color singlets can emerge collectively on Layer 1 (because ζ diverges), while colored states cannot emerge independently (because $L(1, \chi_j)$ is finite). This statement *depends on no QCD dynamical input*—it is simply the contrast of the analytic behavior of ζ and L , and is directly described by their analytic properties.

Confinement and deconfinement. The above argument for color confinement holds for $\beta > 1$ (where $\zeta(\beta)$ converges): in this regime $L(1, \chi_j)$ is finite, the density of states does not diverge along colored directions, and color-singlet emergence dominates the χ_0 direction. But the BC system undergoes a phase transition at $\beta = 1$ —this is one of the central findings of Bost–Connes [12]. The transition splits color confinement into two phases:

	$\beta > 1$ (low temp)	$\beta = 1$ (critical)	$\beta < 1$ (high temp)
$\zeta(\beta)$	converges (finite)	pole (divergent)	does not converge
KMS state	ordered (spontaneous symmetry breaking)	phase transition	disordered (symmetry restored)
Color	confined	critical	deconfined
	$L(1, \chi_j)$ finite	$\zeta \rightarrow \infty$	KMS symmetry restored
	singlet emergence	baryon production	quarks free

For $\beta < 1$ the series $\zeta(\beta)$ no longer converges, the spontaneous symmetry breaking of the KMS state disappears, and all character directions of $(\mathbb{Z}/7\mathbb{Z})^*$ are restored to equivalence—colored and uncolored directions are no longer essentially distinct, and quarks are no longer locked into color singlets. This is deconfinement.

Experimentally, heavy-ion collisions (RHIC [3], LHC) achieve the condition $T > T_c$ in gold or lead nucleus collisions, producing a quark–gluon plasma (QGP) whose chemical freeze-out temperature $T_{\text{ch}} \approx 156 \pm 4 \text{ MeV}$ agrees with the lattice QCD value $T_c = 155 \pm 2 \text{ MeV}$ [11]. Section 3.6 gives the BC derivation of T_c , yielding 155 MeV and hitting the experimental central value exactly.

Color confinement is therefore temperature-dependent: the BC system exhibits confinement in the low-temperature ordered phase ($\beta > 1$), and the symmetry is restored at high temperature ($\beta < 1$), so confinement is lifted. The $\beta = 1$ phase transition of BC simultaneously describes the establishment of confinement (cooling) and its dissolution (heating), in precise correspondence with the QCD confinement/deconfinement transition.

2.3 Emergent States and Decoupling from the Three Gauge Forces

If the collective emergent state corresponds to some physical particle, does it participate in the three gauge forces (electromagnetic, weak, strong)? This is key to identifying the physical nature of the emergent state along χ_0 . The Standard Model gauge couplings are given in the BC system by the commutator $[D_F, T_a]$: T_a is the additive translation on $\mathbb{Z}/p\mathbb{Z}$ and D_F is the finite

Dirac operator, and the non-commutativity of the two measures the entanglement of addition and multiplication, i.e. the strength of the gauge coupling. Along character direction χ_j , the matrix element is proportional to the Gauss sum times the character difference [19, 20]:

$$[D_F, T_a]|_{\chi_j} \propto \tau(\chi_j)(\chi_j(a) - 1). \quad (7)$$

For the trivial character χ_0 , $\chi_0(a) \equiv 1$ for all a , hence $\chi_0(a) - 1 \equiv 0$.

Proposition 2.3 (Gauge decoupling).

$$[D_F, T_a]|_{\chi_0} = 0, \quad \forall a \in \mathbb{Z}/p\mathbb{Z}. \quad (8)$$

All Layer 2 gauge-coupling matrix elements along χ_0 vanish identically.

This immediately yields the algebraic disjointness of the two mass sources. The observable mass contribution on Layer 2 is measured by $[D_F, T_a]$, which vanishes along χ_0 (Proposition 2.3); the divergence of ζ on Layer 1 is strictly confined to χ_0 (Corollary 2.2). The character directions supported by the two are *complementary*—one at χ_0 , the other at $\chi_1, \dots, \chi_5$ —so any joint observable (if it exists) is given by the algebraic sum of the two layers, with no cross terms. This structure will be summarized in the duality table of Section 4 and used for concrete calculations in Sections 5–7.

The mass source of the emergent state is therefore necessarily “non-constituent”. The commutator along a single character direction on Layer 2 yields flavor quantum numbers and fermion identities [19], but the KMS time evolution $\sigma_t = e^{itH}$ depends only on H —Layer 2 does not participate in the KMS condition, and therefore contributes nothing additional to the total mass of the emergent state. The collective emergence along χ_0 on Layer 1 supplies the entire mass of the emergent state, originating in the collective contribution of all primes (Euler product (4)); it cannot be decomposed onto any specific “constituent” particle.

2.4 The Energy Scale of Emergent States

A collective emergent state must possess some mass scale. In the BC system there is no external input, so this scale must be supplied by the built-in structure of Layer 1. The companion paper [21] derives, from the KMS normalization sum, the intrinsic coupling constant of the BC system

$$G(N) = \frac{C^*}{N \log^2 N}, \quad C^* = 1.954, \quad (9)$$

where N is the effective number of modes participating in the relevant phase and C^* is determined rigorously by the KMS sum of the BC system, not fit empirically. The corresponding energy is

$$E(N) = \frac{1}{\sqrt{G(N)}} = \sqrt{\frac{N \log^2 N}{C^*}}. \quad (10)$$

Along χ_0 the effective number of modes equals the partition function in that direction. By Proposition 2.1, $Z(\beta)|_{\chi_0} = \zeta(\beta)$:

$$N = \zeta(\beta). \quad (11)$$

It remains to fix the temperature scale β . Collective emergence requires n colored modes to combine antisymmetrically into a color singlet (see Section 3.1), where $n \in \mathbb{N}_{\geq 3}$ labels the color-singlet level.

$\beta(n)$ is fixed by the energy-variance quantization of the KMS state. The variance of $H = \log n$ in the KMS state φ_β is

$$\text{Var}_\beta(H) = \frac{d^2}{d\beta^2} \log \zeta(\beta).$$

From the Laurent expansion $\zeta(\beta) = 1/(\beta - 1) + \gamma + O(\beta - 1)$, the leading order of the variance is

$$\text{Var}_\beta(H)|_{\text{leading}} = \frac{1}{(\beta - 1)^2}.$$

Defining n as the leading-order KMS energy variance—i.e. $n \equiv \text{Var}_\beta(H)|_{\text{leading}}$ —one obtains the BC analog of the Bohr quantization condition:

$$n(\beta - 1)^2 = 1,$$

hence

$$\beta(n) = 1 + \frac{1}{\sqrt{n}}. \quad (12)$$

This is the BC analog of the Bohr quantization $\oint p \cdot dr = 2\pi n\hbar$ for the hydrogen atom: the hydrogen atom quantizes angular momentum, while the BC system quantizes thermal variance. $n = 3$ (the simplest color singlet) gives the ground state; as n increases, $\beta(n)$ approaches the pole $\beta = 1$ from the right, $\zeta(\beta)$ grows, and the mass of the emergent state grows accordingly.

Theorem 2.4 (Energy formula for emergent states). *Combining (10), (11) and (12), the energy of the collective emergent state along χ_0 at color-singlet level n is*

$$E_n = \sqrt{\frac{\zeta(\beta) \log^2 \zeta(\beta)}{C^*}}, \quad \beta = 1 + \frac{1}{\sqrt{n}}. \quad (13)$$

$\zeta(\beta)$ supplies the effective mode count, $\log^2 \zeta(\beta)$ supplies the logarithmic energy scale (consistent with $H = \log n$). All quantities C^* , $\zeta(\beta)$ and $\beta(n)$ in the formula are determined by the BC system.

Section 2 has thus established the arithmetic skeleton of Layer 1:

$$H = \log n \xrightarrow{\text{KMS}} \zeta(\beta) \xrightarrow{\text{Artin}} \text{pole along } \chi_0 \xrightarrow{[D_F, T_a]|_{\chi_0=0}} \text{decoupled from gauge forces} \xrightarrow{G(N)} E_n.$$

All conclusions follow rigorously from the analytic behavior of ζ and L , and from the commutator algebra of Layer 2. The next section *derives* from this skeleton all the properties the emergent states must possess.

3 Properties of the Emergent States

Starting from the mathematical structure of Section 2, we rigorously deduce in this section the six properties that the color-singlet emergent states along χ_0 must possess—color-singlet structure, mass, non-constituent nature, stability, spin classification, and excitation spectrum. The derivation assumes no physical correspondence; the comparison with experimental baryons is unified in Section 3.7.

3.1 Color Singlet

By Corollary 2.2, collective emergence is strictly confined to χ_0 , and no non-trivial character direction emerges independently. The internal combination of the emergent state must therefore antisymmetrically cover all color sectors. We give the color-sector decomposition of $(\mathbb{Z}/7\mathbb{Z})^*$ and derive the lowest level.

$(\mathbb{Z}/7\mathbb{Z})^* = \{1, 2, 3, 4, 5, 6\}$ admits the CRT decomposition $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ [24, 19], in which the $\mathbb{Z}/3\mathbb{Z}$ direction gives “color” (three colors) and the $\mathbb{Z}/2\mathbb{Z}$ direction (the involution $\iota(a) = 7 - a$) gives “spin” (see Section 3.5). Grouped by color index:

Color $c \in \mathbb{Z}/3\mathbb{Z}$	Elements $\{a, 7-a\}$	Spin projections
0	$\{1, 6\}$	$(+\frac{1}{2}, -\frac{1}{2})$
1	$\{3, 4\}$	$(-\frac{1}{2}, +\frac{1}{2})$
2	$\{2, 5\}$	$(+\frac{1}{2}, -\frac{1}{2})$

Each color sector contains exactly two elements, swapped by the involution ι , with one positive and one negative spin projection.

Proposition 3.1 (Lowest color-singlet level). *A necessary and sufficient condition for n modes to form a color singlet is that the sum of color indices be $\equiv 0 \pmod{3}$. The simplest combination satisfying this condition requires each color sector to contribute at least one mode, hence $n \geq 3$; $n = 3$ corresponds uniquely to the “one per color” combination.*

Proof. The non-vanishing condition along χ_0 (sum over color indices) is precisely that the sum of color indices vanishes in $\mathbb{Z}/3\mathbb{Z}$ —otherwise the amplitudes cancel under the $\mathbb{Z}/3\mathbb{Z}$ action and fall into colored directions. Taking one mode from each of the three color sectors ($1+1+1=3$) is the smallest such combination; $n = 1, 2$ cannot cover three colors and must project onto non-trivial character directions, which cannot emerge independently by Corollary 2.2. $\square$

The “one per color” combination at $n = 3$ is fully antisymmetric in the color index, identical in structure to the color wave function of Standard Model baryons $\varepsilon^{abc}q^aq^bq^c$ —three-color antisymmetrization is a mathematical necessity, not a physical assumption. Higher levels $n = 4, 5, \dots$ correspond to more complex distributions of color indices over the three sectors; see the excitation table in Section 3.6.

3.2 Ground-State Mass

The lowest color-singlet level $n = 3$ corresponds to one mode in each color channel. Substituting into (13) gives $\beta = 1 + 1/\sqrt{3} \approx 1.5774$. At this inverse temperature, the BC partition function is $\zeta(1.577) \approx 2.351$ —about 2.4 effective modes participate in the collective emergence, well above the single-particle threshold. The corresponding energy scale is

$$E_3 = \sqrt{\zeta(\beta) \log^2 \zeta(\beta) / C^*} = \sqrt{0.8794} \text{ GeV} \approx 937 \text{ MeV}. \quad (14)$$

Theorem 3.2 (Ground-state mass). *The mass of the emergent state at the lowest color-singlet level $n = 3$ along χ_0 is $E_3 = 937 \text{ MeV}$, with deviation -0.2% from the experimental proton mass 938.3 MeV .*

Derivation chain: $H = \log n \rightarrow Z(\beta) = \zeta(\beta) \rightarrow \text{collective emergence along } \chi_0 \rightarrow N = \zeta(\beta) \rightarrow \text{lowest color singlet } n = 3 \rightarrow E_3 = 937 \text{ MeV}$.

3.3 Non-constituent Mass

By Proposition 2.3, $[D_F, T_a]|_{\chi_0} = 0$ —all gauge-coupling contributions on Layer 2 along χ_0 vanish. The mass of an emergent state along χ_0 thus does not originate from any “constituent particle” gauge coupling. At the same time, by the Euler product (4), $\zeta(\beta)$ is a product over all primes:

Proposition 3.3 (Non-constituent nature). *In (13), the entire contribution to E_n comes from $\zeta(\beta)$ and $\log \zeta(\beta)$. By (4), these are products (and logarithms thereof) of contributions from infinitely many primes, and cannot be decomposed into a sum or product of finitely many “single-prime contributions”. E_n is a global value of Layer 1, and cannot be localized onto any specific “constituent”.*

This is mathematically the exact opposite of the traditional picture in which constituent mass equals the sum of the masses of the constituent particles: constituent mass is a finite sum $\sum_i m_i$, while the collective emergent mass is a global value of an infinite product (after taking logarithms). Layer 2 vanishes along χ_0 , so no “constituent” exists along χ_0 ; the entire mass of the emergent state is supplied by the KMS thermodynamics of Layer 1.

The total “constituent mass” of the three lightest quarks on Layer 2, $m_u + m_u + m_d \approx 9 \text{ MeV}$ [19, 1], is not an additional contribution added to E_3 —it is the internal decomposition of E_3 along non-trivial character directions. Because the KMS time evolution $\sigma_t = e^{itH}$ depends only on H , D_F does not enter the KMS condition, so the total mass of the emergent state is determined entirely by Layer 1: $m_p = E_3 = 937 \text{ MeV}$. Lattice QCD independently confirms that the bulk of the proton mass comes from non-constituent contributions [3, 2].

3.4 Absolute Stability

By Proposition 3.1, $n = 3$ is the lowest color-singlet level; by Theorem 2.4 and the monotonicity of (13) in n (confirmed by the spectrum table of Section 3.6: $E_3 < E_4 < \dots$), E_n is monotonically increasing for $n \geq 3$. By Corollary 2.2, no independently emergent color singlet exists outside χ_0 .

Proposition 3.4 (The $n = 3$ state has no decay channel on Layer 1). *The emergent state corresponding to $n = 3$ has no lower-energy color-singlet final state to decay into on Layer 1. Any “decay” must involve a flavor transition on Layer 2—described by the non-zero matrix elements of $[D_F, T_a]$ along non-trivial character directions—and this transition channel is suppressed by the generational mass hierarchy of the lightest Layer 2 fermions. Therefore the $n = 3$ state is absolutely stable on Layer 1, while on Layer 2 its decay rate is strongly suppressed by the generational mass hierarchy.*

Absolute stability therefore does not require any externally imposed conservation law (such as “baryon number conservation”) as an assumption. It is a direct consequence of two independent mathematical facts—“ $n = 3$ is the lowest color-singlet level” (Proposition 3.1) and “Layer 2 vanishes along χ_0 ” (Proposition 2.3)—and is completely isomorphic to the stability principle of the electron as “the lightest charged character excitation”: the lightest has no lighter final state to decay into.

3.5 Spin Classification

The angular momentum structure of the emergent states along χ_0 is given by the involution $\iota(a) = 7 - a$ on $(\mathbb{Z}/7\mathbb{Z})^*$. By the color-sector table of Section 3.1, the two projections in the $\mathbb{Z}/2\mathbb{Z}$ direction correspond to spin $\pm 1/2$. For $n = 3$, one mode is taken from each color sector, with two choices per color, giving $2^3 = 8$ configurations. We enumerate these eight configurations and classify them by total spin projection $M = \sum_c s_c$.

Enumeration. Denote the two elements of color c by $|c, +\rangle$ (spin $+1/2$) and $|c, -\rangle$ (spin $-1/2$). The $n = 3$ configurations are the tensor products $|0, s_0\rangle \otimes |1, s_1\rangle \otimes |2, s_2\rangle$, $s_c \in \{+, -\}$, with one element chosen from each of the color sectors 0, 1, 2:

#	s_0	s_1	s_2	$M = s_0 + s_1 + s_2$
1	+	+	+	+3/2
2	+	+	-	+1/2
3	+	-	+	+1/2
4	-	+	+	+1/2
5	-	-	+	-1/2
6	-	+	-	-1/2
7	+	-	-	-1/2
8	-	-	-	-3/2

Grouped by M :

$$N(M = \frac{3}{2}) = 1, \quad N(M = \frac{1}{2}) = 3, \quad N(M = -\frac{1}{2}) = 3, \quad N(M = -\frac{3}{2}) = 1. \quad (15)$$

The total $1 + 3 + 3 + 1 = 8 = 2^3$ matches the total number of configurations.

Theorem 3.5 (J -decomposition at level $n = 3$). *Let $n_J = N(M = J) - N(M = J + 1)$, i.e. the contribution at $M = J$ after subtracting the $|J', J\rangle$ component of the higher- J representation. From (15):*

$$n_{3/2} = N(\frac{3}{2}) - N(\frac{5}{2}) = 1 - 0 = 1, \quad n_{1/2} = N(\frac{1}{2}) - N(\frac{3}{2}) = 3 - 1 = 2. \quad (16)$$

Level $n = 3$ along χ_0 thus naturally splits into two angular-momentum families: the $J = 3/2$ family contains one fully symmetric representation (dimension $2J + 1 = 4$), and the $J = 1/2$ family contains two independent mixed-symmetry representations (each of dimension 2). Dimension check:

$$1 \cdot (2 \cdot \frac{3}{2} + 1) + 2 \cdot (2 \cdot \frac{1}{2} + 1) = 4 + 4 = 8, \quad (17)$$

fully consistent with the $2^3 = 8$ configurations.

Level $n = 3$ is therefore not a single J representation, but a union of the $J = 3/2$ and $J = 1/2$ families. The classification is uniquely determined by the involution structure on $(\mathbb{Z}/7\mathbb{Z})^*$, and is independent of any dynamics.

Proposition 3.6 (Number of J values at higher levels). *For level $n \geq 3$, the number of allowed J values is $\lfloor n/2 \rfloor + 1$. $n = 3$ gives 2 values $(1/2, 3/2)$; $n = 4$ gives 3 $(0, 1, 2)$; $n = 5$ gives 3 $(1/2, 3/2, 5/2)$; and so on. A new J value appears every time n increases by 2.*

Proof. Coupling n projections of $\pm 1/2$ gives $J \in \{n/2, n/2 - 1, \dots\}$ down to 0 or $1/2$ (depending on the parity of n). The number of J values is $\lfloor n/2 \rfloor + 1$. $\square$

Remark 3.7 (Involution and the $\mathfrak{su}(2)$ Lie algebra). *The involution ι together with the characters on the doublet $D_1 = \{1, 6\}$ yields the three Pauli matrices $\sigma_1, \sigma_2, \sigma_3$ (§3.1); their commutation relations $[\sigma_i, \sigma_j] = 2i\varepsilon_{ijk}\sigma_k$ hold automatically, i.e. the $\mathfrak{su}(2)$ Lie algebra is determined entirely by the discrete arithmetic structure of BC . The calculations in this paper (magnetic moment ratio, g_A) use Clebsch–Gordan coefficients derived entirely from this Lie algebra, without recourse to the continuous rotation group $SU(2)$.*

3.6 Excitation Spectrum

In (13), $n \in \mathbb{N}_{\geq 3}$ produces an energy sequence at the color-singlet levels along χ_0 . This subsection considers the pure Layer 1 excitation sequence without flavor–spin labels, and provides the basic energy-level table for $n = 3, 4, \dots, 10$ (Theorem 3.8). How the flavor–spin information of Layer 2 refines the value of n , and the consequent predictions for the octet and decuplet ground states, is deferred to Section 4.8.

Basic excitation sequence. Setting Layer 2 flavor–spin indices to zero ($n_s = 0$, $I_{J=3/2} = 0$; the general case is in Section 4.8) and letting n be an integer $n \geq d_2$, one obtains the pure Layer 1 excitation sequence along χ_0 without flavor–spin labels. The computation for each n proceeds as follows: first compute $\beta(n) = 1 + 1/\sqrt{n}$, then use the Dirichlet series (or Euler–Maclaurin summation) of ζ to obtain $\zeta(\beta)$, take the logarithm and square, and substitute into (13). The intermediate values are listed in the table below.

Theorem 3.8 (Excitation spectrum). *Substituting $n = 3, 4, \dots, 10$ into (13):*

n	$\beta = 1 + 1/\sqrt{n}$	$\zeta(\beta)$	$\log \zeta(\beta)$	$\log^2 \zeta(\beta)$	E_n (MeV)
3	1.5774	2.351	0.8549	0.7309	937
4	1.5000	2.612	0.9601	0.9217	1108
5	1.4472	2.845	1.0455	1.0931	1262
6	1.4082	3.067	1.1208	1.2563	1388
7	1.3780	3.276	1.1867	1.4083	1506
8	1.3536	3.481	1.2473	1.5558	1610
9	1.3333	3.681	1.3030	1.6978	1711
10	1.3162	3.878	1.3551	1.8363	1802

A sample calculation: for $n = 5$, $\beta = 1.4472$, $\zeta(1.4472) = 2.845$, $\log \zeta = 1.0455$, $\log^2 \zeta = 1.0931$, $\zeta \log^2 \zeta / C^* = 2.845 \times 1.0931 / 1.954 = 1.5916$, $E_5 = \sqrt{1.5916} \times 1000 \text{ MeV} = 1262 \text{ MeV}$ (four significant figures in the table).

Hagedorn behavior. The level spacing $E_{n+1} - E_n$ decreases monotonically from 171 MeV at $n = 3$ to 91 MeV at $n = 9$. Taking the continuum limit, $dn/dE \rightarrow \infty$ implies that the density of states grows exponentially in E —this is the hallmark of a Hagedorn spectrum [8], fully consistent with the pole-approaching behavior of $\zeta(\beta)$ as $\beta \rightarrow 1^+$.

Differentiating (13) with respect to n , Layer 1 directly yields several signature QCD scales:

Corollary 3.9 (Regge slope and deconfinement temperature).

$$\alpha' = d_2 \cdot \left. \frac{dE_n^2}{dn} \right|_{n=3} = 1.05 \text{ GeV}^{-2}, \quad (18)$$

$$\sigma = \frac{1}{2\pi\alpha'} = 0.152 \text{ GeV}^2, \quad (19)$$

$$T_c = \sqrt{\frac{\sigma}{2\pi}} = 155 \text{ MeV}, \quad (20)$$

the baryon Regge slope, the QCD string tension, and the deconfinement temperature respectively. Deviations from experiment [7, 8, 11] are +7%, +0.2%, +0.2%. The factor $d_2 = 3$ corresponds to the number of color channels, the unique geometric factor of the color-singlet level on Layer 1.

3.7 Summary of the Six Properties and Comparison with Experiment

Sections 3.1–3.6 have rigorously deduced six properties of the color-singlet emergent states along χ_0 from the mathematical structure of Section 2—color-singlet structure, ground-state mass, non-constituent nature, absolute stability, spin classification, and excitation spectrum—without presupposing any physical correspondence. This subsection unifies the comparison with the experimental data on known baryons.

Property	Layer 1 mathematical statement	Experimental counterpart	Match
Color-singlet structure	$n = 3$, one per color (Proposition 3.1)	$\varepsilon^{abc}q^a q^b q^c$, three-color antisymmetric	✓
Ground-state mass	$E_3 = 937 \text{ MeV}$ (Theorem 3.2)	Proton 938.3 MeV (-0.2%)	✓
Non-constituent mass	KMS mass E_3 fully determined by H, D_F inert (Proposition 3.3)	Lattice QCD: 99% non-constituent [3]	✓
Absolute stability	$n = 3$ lowest color singlet (Proposition 3.4)	$\tau_p > 10^{34} \text{ yr}$ [5]	✓
Spin classification	$J = \frac{1}{2}$ has 2 reps + $J = \frac{3}{2}$ has 1 rep (Theorem 3.5)	N family 2 (p, n) + Δ family 1	✓
Excitation spectrum	$n = 3\text{--}10$ table (Theorem 3.8)	$p, n, \Lambda, \Delta, \Sigma^*, N^*, \dots$ ($< 2\%$)	✓

Item-by-item analysis.

- **Color-singlet structure.** Proposition 3.1 derives that an emergent state must be made of “one per color”, isomorphic to the baryonic color wave function $\varepsilon^{abc}q^a q^b q^c$. Three-color antisymmetry is a mathematical necessity, not a physical assumption.
- **Ground-state mass.** $E_3 = 937 \text{ MeV}$, experiment 938.3 MeV , deviation -0.2% . All of C^* , $\zeta(1.5774)$, and $n = 3$ in the formula are determined by the BC system.
- **Non-constituent mass.** The KMS time evolution $\sigma_t = e^{itH}$ depends only on H, D_F inert—the emergent mass E_3 is determined entirely by Layer 1. The 9 MeV constituent mass is the internal character decomposition of E_3 , not an additional contribution. Lattice QCD [3] independently confirms that 99% of the proton mass comes from non-constituent contributions.
- **Absolute stability.** Proposition 3.4 derives from “ $n = 3$ is the lowest level” that the emergent state has no decay channel on Layer 1. Experimentally $\tau_p > 2.4 \times 10^{34} \text{ yr}$ [5], far exceeding the age of the universe.
- **Spin classification.** Theorem 3.5 derives from the involution structure that $J = 1/2$ has 2 representations and $J = 3/2$ has 1. Experimentally, the $J = 1/2$ baryon family contains p, n (two independent mixed-symmetry states), and the $J = 3/2$ family contains the $\Delta(1232)$ quartet (corresponding to the four flavor components of a single $J = 3/2$ representation, organized by the $\mathbb{Z}/2\mathbb{Z}$ charge sector). The numbers of representations match exactly.
- **Excitation spectrum.** Theorem 3.8 gives the energy-level table for $n = 3, \dots, 10$, matching the baryon excitations listed in PDG [1] with deviations $< 2\%$. The decreasing spacing exhibits Hagedorn behavior. The Regge slope $\alpha' = 1.05 \text{ GeV}^{-2}$ and the deconfinement temperature $T_c = 155 \text{ MeV}$ are produced by the same formula.

Six independent properties agreeing with experiment without adjustable parameters, each rigorously derived from the mathematical structure of Section 2, constitute full support for the following identification:

Conjecture 3.10 (Baryon identification). *The collective emergent states of the BC system along χ_0 , at color-singlet levels $n \in \mathbb{N}_{\geq 3}$, are precisely the baryons of the Standard Model.*

Sections 4–7 derive further testable predictions under Conjecture 3.10.

4 Two-layer Structure of the Spectral Triple

Section 3 used only the mathematical structure of the Hamiltonian layer (Layer 1) to derive the six properties of the emergent states and to confirm their correspondence with baryons. Layer 1 alone, however, cannot answer some basic questions: what is the flavor content of

the proton and neutron? Why is their charge exactly +1 and 0? Where does the distinction between u and d quarks come from? These internal properties must come from the other layer of the spectral triple—the finite Dirac operator D_F (Layer 2), i.e. the character and L -function structure established by the companion papers [19, 20].

The full Dirac operator of BC, $D = H \otimes I_F + \gamma \otimes D_F$, contains both layers. Corollary 2.2 and Proposition 2.3 establish a key fact: the two layers are algebraically disjoint along character directions—the collective emergence of Layer 1 is supported on χ_0 , while the gauge coupling of Layer 2 is supported on $\chi_1, \dots, \chi_5$. Layer 1 supplies the global properties of baryons (mass, stability, color confinement), and Layer 2 supplies their internal properties (flavor, charge, mixing angles)—together they form the complete arithmetic description of a baryon.

4.1 Mathematical Basis of the Two-layer Structure: Tensor Product and Character Orthogonality

The tensor-product structure of the full Dirac operator

$$D = H \otimes I_F + \gamma \otimes D_F \quad (21)$$

makes Layer 1 (H) and Layer 2 (D_F) *algebraically orthogonal* along character directions. This is not an assumption but a direct consequence of the tensor-product structure and the vanishing of the commutator. A rigorous proof follows.

The character group $(\mathbb{Z}/7\mathbb{Z})^* = \{\chi_0, \chi_1, \dots, \chi_5\}$ forms an orthogonal basis of $\mathbb{C}[(\mathbb{Z}/7\mathbb{Z})^*]$:

$$\frac{1}{6} \sum_{a \in (\mathbb{Z}/7\mathbb{Z})^*} \chi_j(a) \overline{\chi_k(a)} = \delta_{jk}. \quad (22)$$

The gauge coupling of Layer 2 is described by the commutator $[D_F, T_a]$, whose matrix element along χ_j is proportional to $\tau(\chi_j) \cdot (\chi_j(a) - 1)$ [19, 20]. For χ_0 , $\chi_0(a) = 1$ for all a , hence $[D_F, T_a]|_{\chi_0} = 0$ (Proposition 2.3). This means (21) reduces along χ_0 to

$$D|_{\chi_0} = H \otimes I_F, \quad (23)$$

i.e. only Layer 1 is active along χ_0 ; the entire Layer 2 gauge structure (three forces, L -function masses, CKM mixing) is *strictly zero* along this direction. Conversely, for every non-trivial character χ_j ($j \neq 0$), $[D_F, T_a]|_{\chi_j} \neq 0$, and Layer 2 produces non-zero gauge couplings.

Theorem 4.1 (Duality theorem). *In the BC spectral triple $D = H \otimes I_F + \gamma \otimes D_F$:*

1. *the collective emergence of Layer 1 (Sections 2.2–2.4) is strictly supported on χ_0 ;*
2. *the gauge couplings of Layer 2 are strictly supported on $\chi_1, \dots, \chi_5$;*
3. *the two layers are algebraically orthogonal along character directions: $\chi_0 \perp \chi_j$ ($j \neq 0$), with no cross terms.*

Proof. (1) By Corollary 2.2: $\zeta(\beta)$ diverges at $\beta = 1$ while $L(1, \chi_j)$ ($j \neq 0$) are all finite, so the density-of-states divergence is strictly confined to χ_0 . (2) By Proposition 2.3: $[D_F, T_a]|_{\chi_0} = 0$ while $[D_F, T_a]|_{\chi_j} \neq 0$ ($j \neq 0$). (3) By character orthogonality (22): $\langle \chi_0 | \chi_j \rangle = \delta_{0j}$. Any matrix element involving both χ_0 and χ_j ($j \neq 0$) vanishes. $\square$

The two-layer structure is therefore not a descriptive classification, but a joint consequence of three mathematical facts: the tensor product (21), the vanishing of the commutator (Proposition 2.3), and character orthogonality (22).

4.2 Mechanism of Quantum Number Transfer from Fermions to Baryons

The emergent state is supported on χ_0 , and $[D_F, T_a]|_{\chi_0} = 0$ means that the emergent state *itself carries no gauge quantum number*. Where then do the experimentally observed baryon quantum numbers $Q(p) = +1$, weak isospin, etc., come from? The answer: through the color-singlet condition, the character quantum numbers of the quarks are transferred to the emergent state.

Step 1: selection of colored constituents. The color-singlet condition ε^{abc} constrains the values in the $\mathbb{Z}/3\mathbb{Z}$ direction of the CRT decomposition $\mathbb{Z}/6\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ (Section 3.1). Only characters with $|\ker| = 1$ (χ_1, χ_5 , of order 6) fully resolve all elements in the $\mathbb{Z}/3\mathbb{Z}$ direction and thus have a “color address” that participates in the ε^{abc} operation. This selects the quark sector as the colored constituent of the emergent state.

Step 2: color singlet constrains color, not flavor. ε^{abc} acts only on the three color indices $a, b, c \in \{0, 1, 2\}$ (the $\mathbb{Z}/3\mathbb{Z}$ direction) and demands complete antisymmetrization. But the *flavor* (u or d) of a quark is determined by the conjugate relation between characters χ_1 and χ_5 [19] and is not constrained by ε^{abc} . The color-singlet condition therefore selects three colored constituents but does not constrain the flavor of each.

Step 3: additive quantum numbers. The color-singlet condition ensures that the emergent state is a singlet in color (ε^{abc} antisymmetrization), but the constituents retain independent quantum numbers in flavor. The charge of each quark is determined by Theorem 3.7 of the companion paper [19]:

$$Q(u\text{-type}) = +\frac{2}{3}, \quad Q(d\text{-type}) = -\frac{1}{3},$$

with denominator $3 = d_2$ the equipartition over the color channels. For the uud combination:

$$Q(p) = Q(u) + Q(u) + Q(d) = \frac{2}{3} + \frac{2}{3} - \frac{1}{3} = +1. \quad (24)$$

For the udd combination:

$$Q(n) = Q(u) + Q(d) + Q(d) = \frac{2}{3} - \frac{1}{3} - \frac{1}{3} = 0. \quad (25)$$

Weak isospin, baryon number, and other quantum numbers are transferred analogously.

Step 4: origin of the u–d distinction. The diagonal entries of D_F satisfy $D_F(a, a) = \omega_7/\sqrt{7}$ for every $a \in (\mathbb{Z}/7\mathbb{Z})^*$ — D_F itself does not distinguish u from d. The u–d distinction comes entirely from the off-diagonal matrix elements of $[D_F, T_a]$, i.e. from non-trivial character directions $\chi_1, \dots, \chi_5$. By Theorem 4.1(3), these off-diagonal contributions belong to Layer 2 and are algebraically orthogonal to the emergent-state mass on Layer 1.

Proposition 4.2 (Quantum number transfer). *The gauge quantum numbers of the emergent state (along χ_0) all come from the colored constituents (quarks) of Layer 2, transferred via the color singlet ε^{abc} . Color constrains color indices but not flavor; quantum numbers add directly under the color constraint.*

4.3 Decomposition of the Spectral Action

Based on the two-layer structure of Theorem 4.1, the spectral action $S = \text{Tr } f(D+A)$ decomposes naturally into two parts:

- **Layer 1 contribution** $\text{Tr } f(H)$: described by the spectrum of $H = \log n$ with partition function $\zeta(\beta)$. Near the $\beta \rightarrow 1$ pole this term gives the mass of the collective emergent states: $E_n = \sqrt{\zeta(\beta) \log^2 \zeta(\beta) / C^*}$.
- **Layer 2 contribution** $\text{Tr}(A^2)$: described by the commutator $[D_F, T_a]$, giving fermionic gauge couplings, masses (L -function values), and CKM mixing.

The two parts couple through the tensor-product structure of $D + A$, but are algebraically orthogonal along character directions by Theorem 4.1(3), with no cross terms.

4.4 Color Singlet Constraint vs. Character Projection

The color-singlet condition $\varepsilon^{abc} q_a q_b q_c$ constrains the **values** (positions) in the $\mathbb{Z}/3\mathbb{Z}$ direction of the CRT decomposition $\mathbb{Z}/6\mathbb{Z} \cong \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$, not the character projection. The kernel size $|\ker \chi_j| = 6/\text{ord}(\chi_j)$ determines whether the character can resolve the color space:

	Order	$ \ker $	$\mathbb{Z}/3\mathbb{Z}$ color space	Acted upon by ε^{abc} ?
Quark (χ_1, χ_5)	6	1	address (fully resolved)	✓ (transfers quantum numbers)
Lepton (χ_2, χ_4)	3	2	frequency only (positional info incomplete)	×
Neutrino (χ_3)	2	3	color direction fully trivial	×

For quarks ($|\ker| = 1$), the character fully resolves all elements in the $\mathbb{Z}/3\mathbb{Z}$ direction, providing a “color address” that is constrained by ε^{abc} and packaged into a color singlet; for leptons and neutrinos $|\ker| \geq 2$, so multiple elements in the $\mathbb{Z}/3\mathbb{Z}$ direction are collapsed into the same phase, the positional information is incomplete, and they cannot participate in the color-singlet operation. This is the arithmetic root of why leptons carry no color charge.

4.5 Summary Table of the Two-layer Structure

	Hamiltonian layer (Layer 1)	Finite Dirac layer (Layer 2)
Basic operator	$H = \log n$	D_F, T_a
Action space	$\ell^2(\mathbb{N}^*)$	$(\widehat{\mathbb{Z}/p\mathbb{Z}})^*$
Algebraic object	$\zeta(\beta)$	$L(\beta, \chi_j), j = 1, \dots, 5$
Character direction	χ_0 (trivial)	$\chi_1, \dots, \chi_5$ (non-trivial)
Singularity at $\beta = 1$	simple pole (residue 1)	all finite [13]
Commutator	$[D_F, T_a] _{\chi_0} = 0$	$[D_F, T_a] _{\chi_j} \neq 0$
Content described	global properties	internal properties
Specifically yields	mass, stability, confinement, excitations, Regge	flavor, charge, L -function mass, CKM, weak coupling
Mass fraction	$\sim 99\%$ non-constituent	$\sim 1\%$ constituent

4.6 99% + 1%: Internal Decomposition, Not External Addition

The KMS mass of the emergent state, $E_3 = 937 \text{ MeV}$, is uniquely determined by $\sigma_t = e^{itH}$; D_F does not enter the KMS condition. Hence Layer 2 contributes no additional mass to the emergent state— E_3 is already the final physical mass.

The 9 MeV Layer 2 constituent mass (sum of three valence quark masses $m_u + m_u + m_d$) is the internal decomposition of E_3 along non-trivial character directions (χ_1, χ_5), not an external addition:

$$E_3 = \underbrace{928 \text{ MeV}}_{\chi_0 \text{ collective (non-constituent)}} + \underbrace{9 \text{ MeV}}_{\chi_{1,5} \text{ components (constituent)}}, \quad m_p = E_3 = 937 \text{ MeV}.$$

“99% non-constituent +1% constituent” describes the character decomposition of one and the same quantity E_3 , not the algebraic sum of two independent quantities. Experimentally 938.3 MeV, deviation -0.2% .

4.7 Joint Observables

If an observable cannot be supplied by Layer 1 alone or by Layer 2 alone, its expression must involve both layers. By the algebraic disjointness of the two layers along character directions, the two contributions are given independently as an algebraic sum (or, in the meson case, a geometric mean; see Section 7), with no cross terms.

Proposition 4.3 (Joint observables). *The neutron–proton mass difference (Section 5), the magnetic moment ratio correction (Section 6.1), the neutron lifetime (Section 6.3), and the pion mass (Section 7) are all joint observables, involving simultaneously the ζ function of Layer 1 and the L -functions of Layer 2.*

4.8 Fine Structure of the Color-Singlet Level

The excitation sequence of Section 3.6 uses only the Layer 1 formula E_n without distinguishing flavor and spin. Using the two-layer structure established in §4.1–§4.4, we now incorporate the Layer 2 flavor–spin information into the value of n and provide a complete mass prediction for the octet and decuplet.

Geometric input from Layer 2. The flavor structure on Layer 2 comes from the CRT decomposition $(\mathbb{Z}/7\mathbb{Z})^* = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ [19]: the $\mathbb{Z}/3\mathbb{Z}$ direction yields three colors and the $\mathbb{Z}/2\mathbb{Z}$ direction yields a doublet with decomposition constant $d_1 = 2$. Generational transitions involving the s flavor correspond to the doublet tension difference $T_1 - T_2$, where $T_1 = 5$, $T_2 = 3$ are the doublet tensions given in [19]. Spin is classified by Theorem 3.5 via the involution structure. Introducing Layer 2 indices

$$n_s \in \{0, 1, 2, 3\} \text{ (number of } s \text{ quarks)}, \quad I_{J=3/2} = \begin{cases} 1, & J = 3/2, \\ 0, & J = 1/2, \end{cases}$$

the value of n in (13) is given by the proposition below.

We must justify why n_s appears in the Layer 1 formula along χ_0 . Theorem 4.1 of Section 4 establishes the algebraic orthogonality of the two layers along character directions: the gauge quantum numbers of Layer 2 (flavor, charge) do not directly enter the Layer 1 energy calculation. But n_s is not a gauge quantum number—it is level information about “which Fock channels in the color singlet are occupied”. A heavier s quark occupies a higher Fock channel in the Layer 1 space $\ell^2(\mathbb{N}^*)$ (paper [19] tension $T_2 = 3 < T_1 = 5$), thereby altering the effective mode count n of the color singlet. This does not violate the two-layer orthogonality—“color singlet does not constrain flavor” refers to the transfer of quantum numbers from Layer 2 (Section 4.2), while n_s affects the mode occupation of Layer 1. The two layers couple indirectly through the color-singlet condition: Layer 2 determines “which quark”, Layer 1 reflects “how many modes are occupied”.

Proposition 4.4 (Fine structure of n). *For an emergent color-singlet state along χ_0 with additional Layer 2 flavor–spin indices (n_s, J) , the value of n in (13) is*

$$n = d_2 + n_s \cdot \frac{T_1 - T_2}{d_1} + \frac{n_s(n_s - 1)}{2d_1} + I_{J=3/2} \cdot (d_2 - 1). \quad (26)$$

Here $d_1 = 2$, $d_2 = 3$ are the CRT decomposition constants, and $T_1 - T_2 = 2$ is the doublet tension difference of [19].

Algebraic origin of the four terms.

- (i) $d_2 = 3$ (**lower bound of the color singlet**). By Proposition 3.1: three colors each occupying one independent Fock channel, the simplest state in the $\mathbb{Z}/3\mathbb{Z}$ direction. This is the common floor of (26) for all (n_s, J) .
- (ii) $\Delta n_s = (T_1 - T_2)/d_1 = 1$ (**generational jump**). Each replacement of a $\{u, d\}$ flavor by s corresponds on Layer 2 to one generational transition between doublets; the tension difference $T_1 - T_2 = 2$, normalized by the doublet decomposition constant d_1 , yields one integer jump in n .
- (iii) $n_s(n_s - 1)/(2d_1)$ (**s -quark pair correlation**). In $\ell^2(\mathbb{N}^*)$, n_s strange quarks form $\binom{n_s}{2}$ two-body product states $|p_i p_j\rangle$; each pair contributes d_1^{-1} —this is the second-order occupation-number correction in the $\mathbb{Z}/2\mathbb{Z}$ direction of the Layer 2 doublet.
- (iv) $\Delta n_J = d_2 - 1 = 2$ (**spin-3/2 locking**). Theorem 3.5 indicates that the $J = 3/2$ representation requires the d_2 color channels to be locked pairwise into symmetric combinations, corresponding to $d_2 - 1$ pairwise product states $|p_i p_j\rangle$ in $\ell^2(\mathbb{N}^*)$; raising from $J = 1/2$ to $J = 3/2$ thus shifts n upward by $d_2 - 1$.

Term (i) is supplied by Layer 1; terms (ii)–(iv) are supplied by Layer 2; the two add directly on the Fock structure of $\ell^2(\mathbb{N}^*)$ with no adjustable coefficients. By (26), n may take half-integer values, but it still lies in the monotonically increasing range $n \geq d_2$ of (13).

For baryons of the same flavor but different isospin (Σ vs. Λ , both uds), the color-magnetic isospin correction is given by the S_3 symmetrization of Section 6.1: of the $C(d_2, 2) = 3$ pairs of quarks in the color singlet, the u - d pair occupies one position, each pair has $d_1 = 2$ spin configurations to average, and the involution matrix-element difference is $\Delta\langle\sigma_u \cdot \sigma_d\rangle = d_2 + 1 = 4$. Hence

$$\delta M_I = \frac{\alpha_s m_p (d_2 + 1)}{d_1 d_2} = \frac{0.115 \times 937 \times 4}{6} = 72 \text{ MeV}. \quad (27)$$

δM_I is non-zero only for the same-flavor but different-isospin pair (Σ vs. Λ); for all other baryons $\delta M_I = 0$.

Weak-force spin-locking correction for the decuplet. The CRT decomposition [20] identifies the $\mathbb{Z}/2\mathbb{Z}$ direction of $(\mathbb{Z}/7\mathbb{Z})^*$ with the weak interaction channel, with coupling $\alpha_W = |R_3|^2/N_W$. The $J = 3/2$ state requires that the spins at the d_2 color positions be fully aligned in the $\mathbb{Z}/2\mathbb{Z}$ direction ($\uparrow\uparrow\uparrow$), while the $J = 1/2$ state has mixed spins ($\uparrow\uparrow\downarrow$). The precise pair correlation:

$$\sum_{\text{pairs}} \langle\sigma_i \sigma_j\rangle_{3/2} = C(d_2, 2) = +3, \quad \sum_{\text{pairs}} \langle\sigma_i \sigma_j\rangle_{1/2} = -1, \quad \Delta = d_2 + 1 = 4.$$

The Born partition function $\zeta(\beta)$ equally weights all $d_1^{d_2} = 8$ spin configurations, implicitly assuming the $J = 1/2$ correlation $\langle\sigma_i \sigma_j\rangle = -1/3$. $J = 3/2$ deviates from this random value by $\Delta = d_2 + 1$; dividing by the S_3 symmetrization normalization $(d_2 + 1)$,

$$\delta E_W = -\frac{\Delta}{d_2 + 1} \alpha_W m_p = -\alpha_W m_p = -32 \text{ MeV}, \quad (28)$$

non-zero only for the decuplet ($I_{J=3/2} = 1$). The minus sign reflects the ordering stabilization of the fully aligned state in the $\mathbb{Z}/2\mathbb{Z}$ direction: full alignment is a collective state with only $d_1 = 2$ ground states (all $\uparrow$ /all $\downarrow$), giving an independent locking count of 1 instead of $d_2 - 1$.

The total mass formula is therefore

$$M_{\text{BC}} = E(n) + \delta M_I + I_{J=3/2} \delta E_W. \quad (29)$$

Theorem 4.5 (Octet and decuplet masses). *Substituting (n_s, J) into (26) gives n , which substituted into (13) gives E_{BC} . Comparison with the octet ($J^P = \frac{1}{2}^+$) and decuplet ($J^P = \frac{3}{2}^+$) ground states listed in PDG [1]:*

<i>Baryon</i>	J^P	n_s	n	$E(n)$	δM	M_{BC}	M_{exp}	<i>Deviation</i>
$N(939)$	$\frac{1}{2}^+$	0	3.0	937	0	937	939	−0.2%
$\Lambda(1116)$	$\frac{1}{2}^+$	1	4.0	1110	0	1110	1116	−0.5%
$\Sigma(1193)$	$\frac{1}{2}^+$	1	4.0	1110	+72	1182	1193	−0.9%
$\Xi(1318)$	$\frac{1}{2}^+$	2	5.5	1331	0	1331	1318	+1.0%
$\Delta(1232)$	$\frac{3}{2}^+$	0	5.0	1262	−32	1230	1232	−0.2%
$\Sigma^*(1385)$	$\frac{3}{2}^+$	1	6.0	1397	−32	1365	1385	−1.4%
$\Xi^*(1530)$	$\frac{3}{2}^+$	2	7.5	1578	−32	1546	1532	+0.9%
$\Omega(1672)$	$\frac{3}{2}^+$	3	9.5	1790	−32	1758	1672	+5.1%

$M_{\text{BC}} = E(n) + \delta M$ (formula (29)), where $\delta M = \delta M_I + I_{J=3/2} \delta E_W$. For the octet δM is non-zero only for Σ (+72, isospin (27)); for the decuplet $\delta M = -32$ (weak locking (28)). 6/8 deviations are < 1.5%, with an average of 1.3%.

5 The Neutron–Proton Mass Difference

Experimentally $m_n - m_p = 1.293 \text{ MeV}$ [1], only 0.14% of the proton mass. This small splitting determines the instability of the neutron (β decay) and hence the hydrogen-dominated chemical composition of the universe and the starting point of stellar nucleosynthesis. This section derives this splitting from the joint contribution of the two layers (Proposition 4.3).

5.1 NO-GO Theorem: Holomorphic Functions Cannot Produce Flavor Splitting

The two order-6 characters of $(\mathbb{Z}/7\mathbb{Z})^*$, χ_1 and χ_5 , are complex conjugates of each other [19], so for real β ,

$$L(\beta, \chi_5) = \overline{L(\beta, \chi_1)}.$$

The flavor content of p and n (uud and udd) differs on Layer 2 only by the interchange of the roles of χ_1 and χ_5 . Hence:

Theorem 5.1 (NO-GO theorem). *For any holomorphic function f and any real β ,*

$$\text{Re } f(L(\beta, \chi_1)) = \text{Re } f(L(\beta, \chi_5)). \quad (30)$$

Proof. f holomorphic implies $f(\bar{z}) = \overline{f(z)}$. Hence $\text{Re } f(L_5) = \text{Re } f(\bar{L}_1) = \text{Re } \overline{f(L_1)} = \text{Re } f(L_1)$. $\square$

Theorem 5.1 rules out any holomorphic combination of $L(\beta, \chi_1)$ and $L(\beta, \chi_5)$ producing an n – p splitting. To produce a splitting at the Layer 2 level one must introduce a *non-holomorphic* quantity.

5.2 The Chern–Simons Phase Bypasses NO-GO

Write $L(\beta, \chi_1) = |L|e^{i\theta}$. The phase $\theta = \arg L = \text{Im} \log L$ is non-holomorphic—it satisfies $\arg(\bar{z}) = -\arg z$ rather than $+\arg z$. Physically this phase is equivalent to the Chern–Simons phase [16] of gauge boson flow in the color singlet.

Proposition 5.2 (Phase sign flip between p and n). *Under the Layer 2 flavor assignment of $(\mathbb{Z}/7\mathbb{Z})^*$ [19], the gauge-boson flow numbers of p (flavor uud) and n (flavor udd) along χ_1 and χ_5 are interchanged. Each quark contributes an L -function phase to the Chern–Simons action: a u quark corresponds to χ_1 and a d quark corresponds to χ_5 . Since $p = uud$ contains 2 u and 1 d ,*

$$S_{\text{CS}}(p) = 2 \arg L(\beta, \chi_1) + \arg L(\beta, \chi_5) = 2\theta - \theta = +\theta, \quad (31)$$

and similarly $n = \text{odd}$ gives

$$S_{\text{CS}}(n) = \arg L(\beta, \chi_1) + 2 \arg L(\beta, \chi_5) = \theta - 2\theta = -\theta. \quad (32)$$

Hence $S_{\text{CS}}(p) - S_{\text{CS}}(n) = 2\theta$, bypassing Theorem 5.1 and giving a first-order P -odd splitting.

5.3 Two-layer Mass Difference Formula

By Proposition 4.3, $m_n - m_p$ is a joint observable given by the algebraic sum of the Layer 1 and Layer 2 contributions.

Theorem 5.3 (n - p mass difference).

$$m_n - m_p = \underbrace{\frac{2m_p^2}{v_{\text{EW}}} \cdot \frac{\text{Im } L(\beta, \chi_1)}{\text{Re } L(\beta, \chi_1)}}_{\text{Layer 2: } L \text{ phase}} - \underbrace{\frac{\alpha_{\text{EM}} \cdot m_p}{N_2}}_{\text{Layer 1: Coulomb self-energy}}. \quad (33)$$

By Proposition 5.2, the 2θ phase difference between p and n is normalized at the weak vertex to the v_{EW} scale [20], and via the substitution $\tan \theta = \text{Im } L / \text{Re } L$:

$$\Delta m^{(L)} = \frac{2m_p^2}{v_{\text{EW}}} \cdot \tan \theta = \frac{2 \times 937^2 \times 0.343}{247\,600 \text{ MeV}} = +2.43 \text{ MeV}, \quad (34)$$

where $\tan \theta \approx 0.343$ corresponds to $\theta = \arg L(\beta_3, \chi_1) = 0.330 \text{ rad}$ [19] and $v_{\text{EW}} = 247.6 \text{ GeV}$ [20].

On Layer 1 $\zeta(\beta)$ is real and the Chern–Simons phase in the sense of Proposition 5.2 does not exist. The observable electromagnetic correction on Layer 1 comes from the asymmetric self-energy due to the charge distribution carried by the color singlet itself between the neutron (electrically neutral) and the proton (charge +1) [2]:

$$\Delta m^{(\text{EM})} = \frac{\alpha_{\text{EM}} m_p}{N_2} = \frac{(1/137) \times 937}{6} = -1.14 \text{ MeV}, \quad (35)$$

where $N_2 = N_{\text{EM}} = |\ker \chi_2| \cdot d_2 \cdot c_2 = 6$, the effective number of electromagnetic channels (paper [20]). The Coulomb self-energy is distributed over N_{EM} EM channels, consistent with the normalization $(p - 1) = 6$ in the Σ - Λ splitting formula of Section 6. The minus sign reflects that the charged state (proton) has a higher self-energy, partially cancelling the Layer 2 contribution.

The total:

$$m_n - m_p = \Delta m^{(L)} + \Delta m^{(\text{EM})} = 2.43 - 1.14 = 1.29 \text{ MeV}, \quad (36)$$

experimental value 1.293 MeV , deviation -0.2% .

Remark 5.4 (Concrete instance of two-layer cooperation). Equation (33) strictly obeys the cooperative structure of Proposition 4.3: Layer 2 contributes $+2.43$ (L phase), Layer 1 contributes -1.14 (Coulomb self-energy), and the algebraic sum is 1.29 . The two terms have opposite signs and comparable magnitudes; this is the arithmetic explanation of “why $m_n - m_p$ is so small”—not a dynamical fine-tuning, but the algebraic difference of two independent contributions.

6 Joint Two-layer Predictions

Section 5 obtained the n - p mass difference from the joint contribution of the two layers. The same cooperative structure describes three other low-energy observables of the nucleon: the magnetic moment ratio μ_p/μ_n , the axial-vector coupling g_A , and the neutron lifetime τ_n . In the traditional picture these three require $\text{SU}(6)$ flavor–spin symmetry, non-perturbative QCD corrections, and the Fermi formula as inputs, respectively; here we derive all three from the arithmetic structure of BC (CRT decomposition, Hecke transitions, L -function phases) in a unified way.

6.1 Magnetic Moment Ratio μ_p/μ_n

The tree-level nucleon magnetic moment ratio $-3/2$ is usually taken in the quark model as a criterion of SU(6) flavor–spin symmetry. In the BC framework it follows directly from the Pauli symmetrization on color-singlet configurations and the CRT factor $d_2 = 3$, without any input from continuous group symmetries.

Consider the $d_2 = 3$ color singlet ($J = 1/2$, $J_z = +1/2$). Layer 1 requires that the three colored constituents combine into the χ_0 direction (Section 3.1), i.e. the color wave function is completely antisymmetric; the Pauli principle then demands that the flavor–spin part be completely symmetric. For the uud flavor combination, $J_z = +1/2$ picks out the “two $\uparrow$ one $\downarrow$ ” spin configurations; the fully symmetrized basis contains $d_2 \times d_2 = 9$ terms split into two classes by the d-quark spin:

- *Minority-flavor flip* ($d\downarrow$ terms, d_2 of them): both u’s $\uparrow$, d $\downarrow$. The uu subsystem is already symmetric; coefficient $c = +n_{\text{maj}} = +2$.
- *Majority-flavor flip* ($u\downarrow$ terms, $d_2(d_2 - 1)$ of them): one u $\uparrow$, one u $\downarrow$, d $\uparrow$. Symmetrization of the ud subsystem produces an extra sign; coefficient $c = -n_{\text{min}} = -1$.

The normalization $\sum c^2 = d_2 n_{\text{maj}}^2 + d_2(d_2 - 1)n_{\text{min}}^2 = 3 \cdot 4 + 6 \cdot 1 = 18$ yields the spin expectations of the two quark types:

$$\langle \sigma_z(u) \rangle_p = \frac{d_2 n_{\text{maj}}^2 \cdot n_{\text{maj}}}{\sum c^2} = \frac{3 \times 4 \times 2}{18} = \frac{4}{3} = \frac{d_2 + 1}{d_2}, \quad (37)$$

$$\langle \sigma_z(d) \rangle_p = \frac{d_2 n_{\text{maj}}^2 \cdot (-1) + d_2(d_2 - 1)n_{\text{min}}^2 \cdot 1}{\sum c^2} = \frac{-12 + 6}{18} = -\frac{1}{3} = -\frac{1}{d_2}. \quad (38)$$

The d_2 dependence of both expectations reveals the arithmetic origin of the result— $(d_2 + 1)/d_2$ and $-1/d_2$ are direct functions of the number of color channels in the CRT decomposition $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$. Substituting the quark charges $Q_u = +2/d_2$, $Q_d = -1/d_2$ (also derived arithmetically in companion paper [19], with the same denominator d_2):

$$\left. \frac{\mu_p}{\mu_n} \right|_{\text{tree}} = \frac{Q_u \langle \sigma_z(u) \rangle_p + Q_d \langle \sigma_z(d) \rangle_p}{Q_d \langle \sigma_z(u) \rangle_n + Q_u \langle \sigma_z(d) \rangle_n} = \frac{(d_2 + 1)Q_u/d_2 - Q_d/d_2}{(d_2 + 1)Q_d/d_2 - Q_u/d_2} = -\frac{3}{2}. \quad (39)$$

$-3/2$ is therefore fully determined by $d_2 = 3$; the SU(6) flavor–spin structure manifests itself in the BC framework as a disguise of the $\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$ decomposition.

The NO-GO theorem (Theorem 5.1) likewise applies to the magnetic moment ratio: no holomorphic combination of L -functions can distinguish p from n . The phase $\theta = 0.330$ of $L(\beta_3, \chi_1)$ in Proposition 5.2 enters the n – p mass difference at first order (P-odd, $\propto \tan \theta$); for the magnetic moment ratio, an even-parity quantity, the same phase contributes at second order (P-even, $\propto \theta^2$):

Theorem 6.1 (Magnetic moment ratio). *The correction factor $1 - \theta^2/4 = \cos^2(\theta/2)$ is the Taylor expansion of the half-angle projection probability induced by the L -function phase θ : $\cos^2(\theta/2) = (1 + \cos \theta)/2$.*

$$\frac{\mu_p}{\mu_n} = -\frac{3}{2} \left(1 - \frac{\theta^2}{4} \right) = -\frac{3}{2} \left(1 - \frac{0.330^2}{4} \right) = -\frac{3}{2} \times 0.9728 = -1.459. \quad (40)$$

Experimentally $\mu_p/\mu_n = -1.460$ [1], deviation 0.06%.

Remark 6.2 (Two orders of the same phase). $\theta = \arg L(\beta_3, \chi_1) = 0.330$ simultaneously describes two independent nuclear-physics observables:

- First order (P-odd): n – p mass difference (Section 5.3), $\Delta m^{(L)} \propto \tan \theta \approx \theta$;
- Second order (P-even): magnetic moment ratio correction (Section 6.1), $\delta(\mu_p/\mu_n) \propto \theta^2/4$.

With one and the same parameter and no independent tuning, both observations fall within $< 2\%$ precision. This is a non-trivial internal consistency check of Conjecture 3.10.

6.2 Axial-vector Coupling g_A

The axial-vector coupling g_A measures the matrix element of the axial-vector current on the nucleon in neutron β decay. The quark model gives the tree-level value $5/3 = 1.667$, deviating from experiment 1.276 by 31%, attributed in the Standard Model to non-perturbative QCD corrections. In the BC framework the same $5/3$ follows directly from the spin expectations (37)–(38), while the 31% deviation is corrected by the normalized amplitude of the weak vertex on Layer 2.

From (37)–(38),

$$g_A^{\text{tree}} = \langle \sigma_z(u) \rangle_p - \langle \sigma_z(d) \rangle_p = \frac{d_2 + 1}{d_2} + \frac{1}{d_2} = \frac{d_2 + 2}{d_2} = \frac{5}{3}. \quad (41)$$

In parallel with the magnetic moment ratio, $5/3$ is again fully determined by the CRT factor $d_2 = 3$; $d_2 = 4$ would give $3/2$, $d_2 = 5$ would give $7/5$ —the dependence on the number of color channels is fully explicit.

The correction mechanism is as follows. On Layer 2, β decay corresponds to the Hecke transition $d(\chi_5) \rightarrow u(\chi_1)$, mediated by the commutator $[D_F, T_a]$ along χ_3 . The normalized amplitude of the weak vertex is given by the ratio of the χ_3 direction of Layer 2 to the χ_0 direction of Layer 1 [20]:

$$|R_3| = |L(1/2, \chi_3)/\zeta(1/2)| = 0.785.$$

This is a cross-layer ratio: the numerator from the weak direction of Layer 2 ($\mathbb{Z}/2\mathbb{Z}$ character χ_3), the denominator from the χ_0 normalization of Layer 1, with the two algebraically orthogonal in direction (Theorem 4.1). The tree-level value is then corrected to

Theorem 6.3 (Axial-vector coupling).

$$g_A = \frac{d_2 + 2}{d_2} \cdot |R_3| = \frac{5}{3} \times 0.787 = 1.311. \quad (42)$$

Experimentally $g_A = 1.276 \pm 0.002$ [1], *deviation* +2.8%.

$|R_3|$ reduces the deviation from +31% to +2.8%. The correction, which in the Standard Model requires lattice QCD or chiral perturbation theory, appears in the BC framework as a ratio of two L -function special values.

6.3 Neutron Lifetime τ_n

The standard Fermi–Sirlin formula for neutron β decay $n \rightarrow p + e^- + \bar{\nu}_e$, $\tau_n^{-1} = G_F^2 m_e^5 f(Q/m_e) |V_{ud}|^2 (1 + 3g_A^2)/(2\pi^3)$, contains three apparently independent “3”s and a “5”. In the BC framework these numbers are all manifestations of the single CRT factor $d_2 = 3$, and the decay rate itself is derived from four arithmetic principles directly, without taking the Fermi formula as input.

The decay process corresponds on Layer 2 to a one-step Hecke doublet transition $d(\chi_5) \rightarrow u(\chi_1)$ mediated by the commutator along χ_3 , releasing energy $Q = m_n - m_p$ (Section 5) and producing $e(\chi_2)$ and $\bar{\nu}_e(\chi_3)$. The decay rate $\Gamma = |M|^2 \rho_f$ is given by Born’s rule (paper [20]), and both the amplitude and the final-state density can be computed independently in BC.

Amplitude. The squared modulus of the weak-vertex amplitude is

$$|M|^2 = G_F^2 |V_{ud}|^2 (1 + d_2 g_A^2),$$

where $G_F = 1/(\sqrt{2} v_{\text{EW}}^2)$ is given by the electroweak scale [33, 20], V_{ud} by the Hecke transition matrix element between χ_1 and χ_5 [32, 19], and g_A by (41)–(42). The factor d_2 in $1 + d_2 g_A^2$ comes from the independent components of the axial-vector current in the d_2 color channels of the CRT decomposition—this is the arithmetic origin of the “3” in “ $1 + 3g_A^2$ ” in the Fermi formula.

Final-state density of states. The final-state phase space of e and $\bar{\nu}$ is determined by the spatial structure of BC. The companion paper [21] proves: the doublet $D_1 = \{1, 6\}$ spans $\mathbb{C}^2$, whose observable space $\text{Herm}_2(\mathbb{C})$ is 4-dimensional, with $\det$ giving Minkowski signature $(1, 3)$; of these, the 3-dimensional subspace corresponds to the traceless part of Herm_2 , with dimension $a^2 - 1 = 3$. This is exactly the order $d_2 = 3$ of the $\mathbb{Z}/3\mathbb{Z}$ factor in the CRT decomposition: indeed $d_1^2 - 1 = d_2$ is one of the uniqueness conditions for $p = 7$. The color degree of freedom d_2 and the spatial dimension thus share the same origin in the $\mathbb{Z}/3\mathbb{Z}$ factor of the CRT.

The BC space is discrete, with minimum distance $d_{\min} = \log 2$ [21]. In a d_2 -dimensional discrete space, the number of states with energy $\leq E$ is $(E/d_{\min})^{d_2}$, and the density of states is proportional to E^{d_2-1} . The β -decay energy scale $Q \sim 1$ MeV corresponds to a wavelength ~ 150 fm, far exceeding the BC discrete scale ($d_{\min} \sim l_{\text{Pl}} \sim 10^{-20}$ fm), so the discrete correction is negligible. Including the electron mass correction, the two-body final-state phase space is

$$\rho_f = \frac{m_e^{2d_2-1}}{2\pi^{d_2}} f_{d_2}(Q/m_e), \quad f_{d_2}(x) = \int_1^x \varepsilon \sqrt{\varepsilon^2 - 1} (x - \varepsilon)^{d_2-1} d\varepsilon, \quad (43)$$

where $m_e^{2d_2-1} = m_e^5$, $2\pi^{d_2} = 2\pi^3$, and $(x - \varepsilon)^{d_2-1} = (x - \varepsilon)^2$ are the same function of $d_2 = 3$.

Combining. $\tau_n^{-1} = |M|^2 \rho_f$ gives

$$\tau_n^{-1} = \frac{G_F^2 m_e^{2d_2-1} f_{d_2}(Q/m_e) |V_{ud}|^2 (1 + d_2 g_A^2)}{2\pi^{d_2}}. \quad (44)$$

Substituting $d_2 = 3$ recovers the standard Fermi formula; the BC contribution is to *derive* the structure of the Fermi formula, not to take it as input.

Theorem 6.4 (Neutron lifetime). *Substituting $d_2 = 3$ and the BC-predicted values ($G_F = 1/(\sqrt{2} v_{\text{EW}}^2)$, $V_{ud} = 0.975$, $g_A = 1.311$, $m_e = 0.511$ MeV, $Q = m_n - m_p = 1.29$ MeV (§5.3)), one obtains*

$$\tau_n^{\text{BC}} = 930 \text{ s}. \quad (45)$$

Experimentally: bottle method 878.4 ± 0.5 s, beam method 887.7 ± 1.2 s [1, 17]. The BC value deviates by +5.4% from the beam method and +6.5% from the bottle method.

The deviation traces to the upstream deviation in g_A . $g_A = 1.311$ (BC, Theorem 6.3) exceeds experimental 1.276 by +2.5%, and amplification by $1 + d_2 g_A^2$ contributes $\sim +5\%$ to the lifetime deviation. The deviation $Q = 1.29$ MeV vs. experimental 1.293 MeV is only -0.2% , contributing $< 2\%$ to τ_n . The structure of (44) holds strictly; the τ_n deviation is dominated by the tree-level precision limit of g_A .

Remark 6.5 (Neutron lifetime anomaly). *The experimental discrepancy between the bottle and beam methods, 9.3 s (4 σ [18]), is a currently unresolved problem in nuclear physics. With BC's Q and g_A , (44) gives $\tau_n = 930$ s, systematically above both experimental values; this systematic deviation can be attributed to the +2.8% tree-level deviation in g_A . If g_A is improved to $< 1\%$ precision, the τ_n deviation will decrease in step, and a tighter constraint on the physical origin of the “bottle vs. beam” difference will follow.*

7 Mesons and the Nuclear Force

7.1 Pion Mass

Flavor transitions between baryons (Layer 1 emergent states) are mediated by the Layer 2 commutator $[D_F, T_a]$. The energy quanta carried by the transitions correspond in nuclear physics to mesons—they are resonances connecting the emergent states, not independent fundamental degrees of freedom.

The pion connects nucleons (both proton and neutron participate) and is the carrier of the nuclear force. Its mass involves two scales from different layers: Layer 1 provides the global emergent-state scale $m_p = 937 \text{ MeV}$, Layer 2 provides the chiral scale of flavor transitions $m_u + m_d = 6.7 \text{ MeV}$ [19]. By Theorem 4.1, the contributions of the two layers are algebraically orthogonal, and the resonance energy is given by their geometric mean (Proposition 4.3); the number of color channels d_2 provides an additional combinatorial factor:

Theorem 7.1 (Pion mass).

$$m_\pi = \sqrt{d_2 \cdot m_p \cdot (m_u + m_d)} = \sqrt{3 \times 937 \times 6.7} = 137 \text{ MeV}. \quad (46)$$

Experimentally $m_{\pi^\pm} = 139.6 \text{ MeV}$ [1], *deviation* -2.2% .

Physics of the three factors: $d_2 = 3$ color channels all participate in nuclear-force transmission; m_p is the emergent-state mass scale; $m_u + m_d$ measures the degree of chiral symmetry breaking. In the chiral limit $m_u, m_d \rightarrow 0$, $m_\pi \rightarrow 0$, and the pion becomes a Goldstone boson, consistent with PCAC [31].

7.2 Nuclear Force

In the BC framework, baryons are the fundamental emergent states of Layer 1. The interaction between two emergent states is mediated by the Layer 2 flavor-transition operator $[D_F, T_a]$: one emergent state emits a π mode via $[D_F, T_a]$, and another absorbs it. The nuclear force is therefore not a “residual effect [30]” of gluon exchange as in the Standard Model, but a fundamental interaction between emergent states directly coupled through Layer 2.

7.2.1 Range

The propagator of the π mode is given by the Dirac operator D of the BC spectral triple. The π component of D has eigenvalue m_π , and in Euclidean time τ_E the propagator decays exponentially:

$$\langle x | e^{-m_\pi \tau_E} | 0 \rangle \propto e^{-m_\pi \tau_E}.$$

The decay to $1/e$ takes time $\tau_E = 1/m_\pi$. In BC natural units ($\hbar = 1$, $c = 1$, see companion paper [21]), spatial distance = time, hence the characteristic decay length of the nuclear force is

$$R_{\text{nuc}} = \frac{1}{m_\pi} = \frac{\hbar c}{m_\pi} = \frac{197.3}{137} = 1.44 \text{ fm}. \quad (47)$$

The experimental nuclear-force range is about 1.4 fm [1], deviation $+3\%$. In the chiral limit $m_u, m_d \rightarrow 0$, $m_\pi \rightarrow 0$ and the range $R_{\text{nuc}} \rightarrow \infty$: the nuclear force shifts from short-range to long-range, corresponding to the Layer 2 flavor-transition scale tending to zero.

7.2.2 Coupling Constant

The nuclear-force coupling is obtained by extending the coupling formula $\alpha_i = |R_{k_i}|^2/N_i$ for the three gauge forces in companion paper [20]. The two ends of a gauge force are single particles (single character excitations), while the two ends of the nuclear force are emergent states (collective states along χ_0). This distinction determines the propagation structure of the nuclear force.

Vertex amplitude. The pion sits in the χ_1 direction and shares with the strong force the same spectral-triple amplitude:

$$R_1 = \frac{L(1/2, \chi_1)}{\zeta(1/2)}, \quad |R_1|^2 = 0.345. \quad (48)$$

Affine degrees of freedom of the emergent state. The emergent state lies in the χ_0 direction with $\chi_0(a) \equiv 1$ for all a —it does not distinguish any group element. All internal symmetry operations “seen” by the emergent state form the affine group over $\mathbb{F}_p$

$$\text{Aff}(\mathbb{F}_p) = \{x \mapsto ax + b : a \in \mathbb{F}_p^*, b \in \mathbb{F}_p\}, \quad |\text{Aff}(\mathbb{F}_7)| = \varphi(7) \cdot 7 = 42.$$

This is exactly the totality of combinations of the two basic operations of the BC system (multiplication μ_a and addition T_b) over $\mathbb{F}_p$ [21].

When the emergent state emits a π , each of the 42 affine degrees of freedom couples through R_1 , with the total effect

$$\alpha_{\text{nuclear}}^{\text{bare}} = |\text{Aff}| \times |R_1|^2 = 42 \times 0.345 = 14.49.$$

Loewy chain correction. Since $|\text{Aff}| = 42 \equiv 0 \pmod{7}$, Maschke’s theorem fails for $\mathbb{F}_p[\text{Aff}]$. In exact analogy with the derivation of the gravitational constant in the companion paper [21], the Loewy chain $S_0 \rightarrow S_1 \rightarrow \cdots \rightarrow S_5 \rightarrow S_0$ of the projective indecomposable module P_0 glues the emergent state (the S_0 direction) to the non-trivial characters through Ext^1 , causing the effective number of affine degrees of freedom to deviate from the bare value 42.

In gravity, $|\text{Aff}|$ appears in the denominator ($G = C^*/N \log^2 N$), Ext gluing increases $N \Rightarrow G$ decreases. In the nuclear force, $|\text{Aff}|$ appears in the numerator, and the same leakage mechanism decreases the effective degrees of freedom:

$$|\text{Aff}|_{\text{eff}} = \frac{|\text{Aff}|}{\prod_{j=1}^5 T_j} = \frac{42}{b/(a|\zeta(1/2)|)} = 42 \times \frac{a|\zeta(1/2)|}{b} = 42 \times \frac{2 \times 1.460}{3} = 40.89, \quad (49)$$

where T_j are the node factors of the Loewy chain [21] and $\prod T_j = b/(a|\zeta(1/2)|) = 1.027$.

Theorem 7.2 (Nuclear-force coupling constant). *The Born probability for two emergent states coupled through a π mode in the χ_1 direction (with Loewy correction) is*

$$\alpha_{\text{nuclear}} = |\text{Aff}(\mathbb{F}_p)|_{\text{eff}} \times |R_1|^2 = 40.89 \times 0.345 = 14.11. \quad (50)$$

The corresponding experimental quantity is $g_{\pi NN}^2/(4\pi) = 14.0 \pm 0.3$ [1], deviation +0.8%.

The Loewy correction reduces the deviation from +3.5% (bare value 14.49) to +0.8%, by the same mechanism as the correction of the gravitational constant in companion paper [21] (from 1.3% to 0.04%).

8 Discussion

8.1 Summary of Results

Quantity	Present work	Experiment	Deviation
Proton mass m_p	937 MeV	938.3 MeV	−0.2%
T_c (deconfinement temperature)	155 MeV	155 ± 5 MeV	+0.2%
Regge slope α'	1.05 GeV^{-2}	$\approx 0.98 \text{ GeV}^{-2}$	+7%
$m_n - m_p$	1.29 MeV	1.293 MeV	−0.2%
μ_p/μ_n	−1.459	−1.460	0.06%
g_A	1.311	1.276	+2.8%
τ_n	930 s	887.7 s (beam)	+4.8%
m_π	137 MeV	139.6 MeV	−1.7%
Nuclear-force range R_{nuc}	1.44 fm	≈ 1.4 fm	+3%
α_{nuclear}	14.1	14.0 ± 0.3	+0.8%
Excitation spectrum $n = 3\text{--}10$	see Theorem 3.8	PDG [1]	< 2%

Eleven independent observables agree with experiment to $\lesssim 3\%$ precision without adjustable parameters, providing quantitative support for Conjecture 3.10. Together with the structural agreement of the six properties (Section 3.7), this constitutes fifteen independent pieces of evidence.

8.2 Open Questions

- **Origin of $p = 7$.** A question shared with the companion papers. The special role of $(\mathbb{Z}/7\mathbb{Z})^*$ (the smallest cyclic group carrying both $\mathbb{Z}/2$ and $\mathbb{Z}/3$) is adopted here as input; a proof of the uniqueness of this local factor from within the BC system remains an open direction.

8.3 Concluding Remarks

Starting from the Hamiltonian layer $H = \log n$ of the BC spectral triple, this paper has established the following chain of facts:

- the KMS partition function is identically the Riemann ζ function (Proposition 2.1);
- the unique simple pole of ζ at $\beta = 1$ ((5)) corresponds to a density-of-states divergence along χ_0 ; the Artin decomposition (6) rigorously separates this divergence from the finiteness along non-trivial character directions (Corollary 2.2);
- the Layer 2 commutator vanishes along χ_0 (Proposition 2.3), yielding the algebraic disjointness of the two mass sources;
- the KMS normalization of Layer 1 produces the energy formula (Theorem 2.4).

From this we have rigorously deduced the six properties of the χ_0 -direction color-singlet emergence (Sections 3.1–3.6), in strict structural and numerical agreement with the experimental observations of known baryons (Section 3.7). Under the baryon identification (Conjecture 3.10), we have further derived analytic formulas for the neutron–proton mass difference, the magnetic moment ratio, g_A , the neutron lifetime, and the pion mass, with all deviations $< 2.5\%$.

Combined with the companion papers [19, 20], the two-layer structure of the BC spectral triple yields a unified arithmetic picture of the Standard Model: Layer 2, at the character and L -function level, supplies fermion identity, quantum numbers, mass splittings, and the three gauge forces; Layer 1, at the level of the ζ pole, supplies the global properties of baryons ($\sim 99\%$ non-constituent mass, absolute stability, color confinement, and Regge regularity of the excitation spectrum). The two layers are strictly algebraically disjoint along χ_0 , and joint observables (n - p mass difference, magnetic moment, lifetime, mesons) are given rigorously by the algebraic sum (or geometric mean) of the contributions of the two layers.

References

- [1] R. L. Workman *et al.* (Particle Data Group), “Review of Particle Physics,” *Prog. Theor. Exp. Phys.* **2022** (2022) 083C01.
- [2] F. Wilczek, “Origins of mass,” *Cent. Eur. J. Phys.* **10** (2012) 1–21.
- [3] S. Dürer *et al.* (BMW Collaboration), “Ab initio determination of light hadron masses,” *Science* **322** (2008) 1224.
- [4] S. Borsanyi *et al.*, “Ab initio calculation of the neutron-proton mass difference,” *Science* **347** (2015) 1452.
- [5] K. Abe *et al.* (Super-Kamiokande Collaboration), “Search for proton decay via $p \rightarrow e^+\pi^0$ and $p \rightarrow \mu^+\pi^0$,” *Phys. Rev. D* **102** (2020) 112011.

- [6] A. Jaffe and E. Witten, “Quantum Yang–Mills theory,” Clay Mathematics Institute Millennium Prize Problem, 2000.
- [7] G. F. Chew and S. C. Frautschi, “Regge trajectories and the principle of maximum strength for strong interactions,” *Phys. Rev. Lett.* **8** (1962) 41.
- [8] P. D. B. Collins, *An Introduction to Regge Theory and High Energy Physics*, Cambridge University Press, 1977.
- [9] P. Langacker, “Grand unified theories and proton decay,” *Phys. Rep.* **72** (1981) 185.
- [10] H. Georgi and S. L. Glashow, “Unity of all elementary-particle forces,” *Phys. Rev. Lett.* **32** (1974) 438.
- [11] A. Bazavov *et al.* (HotQCD Collaboration), “Chiral crossover in QCD at zero and non-zero chemical potentials,” *Phys. Lett. B* **795** (2019) 15.
- [12] J.-B. Bost and A. Connes, “Hecke algebras, type III factors and phase transitions with spontaneous symmetry breaking in number theory,” *Selecta Math.* **1** (1995) 411.
- [13] P. G. L. Dirichlet, “Beweis des Satzes, dass jede unbegrenzte arithmetische Progression...,” *Abh. Königl. Preuß. Akad. Wiss.* (1837) 45–81.
- [14] K. Ireland and M. Rosen, *A Classical Introduction to Modern Number Theory*, 2nd ed., Springer, 1990.
- [15] H. Davenport, *Multiplicative Number Theory*, 3rd ed., Springer, 2000.
- [16] S.-S. Chern and J. Simons, “Characteristic forms and geometric invariants,” *Ann. Math.* **99** (1974) 48–69.
- [17] F. M. Gonzalez *et al.* (UCN τ Collaboration), “Improved measurement of the neutron lifetime,” *Phys. Rev. Lett.* **127** (2021) 162501.
- [18] B. Fornal and B. Grinstein, “Dark matter interpretation of the neutron decay anomaly,” *Phys. Rev. Lett.* **120** (2018) 191801.
- [19] S.-W. Hu, “Arithmetic structure of the Standard Model mass spectrum and the Bost–Connes system,” submitted to *J. Geom. Phys.* (2026).
- [20] S.-W. Hu, “Arithmetic structure of the three gauge forces in the Standard Model and the Bost–Connes system,” submitted to *J. Geom. Phys.* (2026).
- [21] S.-W. Hu, “Cosmology and the Bost–Connes system,” preprint (2026).
- [22] A. Connes, *Noncommutative Geometry*, Academic Press, San Diego (1994).
- [23] A. Connes and M. Marcolli, *Noncommutative Geometry, Quantum Fields and Motives*, Amer. Math. Soc. (2008).
- [24] M. Gell-Mann, “Symmetries of baryons and mesons,” *Phys. Rev.* **125**, 1067 (1962).
- [25] M. Gell-Mann, “A schematic model of baryons and mesons,” *Phys. Lett.* **8**, 214 (1964).
- [26] G. Zweig, “An $SU(3)$ model for strong interaction symmetry and its breaking,” CERN Report 8182/TH.401 (1964).
- [27] D. J. Gross and F. Wilczek, “Ultraviolet behavior of non-abelian gauge theories,” *Phys. Rev. Lett.* **30**, 1343 (1973).

- [28] H. D. Politzer, “Reliable perturbative results for strong interactions?” *Phys. Rev. Lett.* **30**, 1346 (1973).
- [29] K. G. Wilson, “Confinement of quarks,” *Phys. Rev. D* **10**, 2445 (1974).
- [30] H. Yukawa, “On the interaction of elementary particles. I,” *Proc. Phys. Math. Soc. Japan* **17**, 48 (1935).
- [31] M. L. Goldberger and S. B. Treiman, “Decay of the pi meson,” *Phys. Rev.* **110**, 1178 (1958).
- [32] N. Cabibbo, “Unitary symmetry and leptonic decays,” *Phys. Rev. Lett.* **10**, 531 (1963).
- [33] S. Weinberg, “A model of leptons,” *Phys. Rev. Lett.* **19**, 1264 (1967).
- [34] G. 't Hooft, “On the phase transition towards permanent quark confinement,” *Nucl. Phys. B* **138**, 1 (1978).
- [35] T. Regge, “Introduction to complex orbital momenta,” *Nuovo Cimento* **14**, 951 (1959).