

# L12: PIDs are UFDs and Polynomial Rings

## Definition 12.1: Ascending Chains, Noetherian Ring

Let  $R$  be a commutative ring with  $1 \neq 0$ .

An **ascending chain** of ideals in  $R$  is a sequence

$$I_1 \subset I_2 \subset I_3 \subset \dots \subset R$$

We say an ascending chain **stabilizes** if there exists  $N \in \mathbb{N}$  such that for all  $n, m \geq N$ ,  $I_n = I_m$ .

We say  $R$  satisfies the **ascending chain condition** (a.c.c.) if every ascending chain stabilizes.

If  $R$  satisfies the a.c.c., we say it is a **Noetherian ring**.

## Theorem 12.2: PID is Noetherian

If  $R$  is a PID, then  $R$  is Noetherian.

**Proof.** Let

$$I_1 \subset I_2 \subset I_3 \subset \dots \subset R$$

be an ascending chain in a PID.

Consider

$$I := \bigcup_{n \in \mathbb{N}} I_n$$

which is an ideal. Then since  $R$  is a PID,  $I = (a)$  for some  $a \in R$ .

Therefore

$$a \in I = \bigcup_{n \in \mathbb{N}} I_n \implies a \in I_N$$

for some  $N \in \mathbb{N}$ .

Hence  $(a) \subset I_n$  implying  $I \subset I_N$  and so we deduce

$$I = I_N = I_{N+1} = I_{N+2} = \dots$$

■

## Theorem 12.3: PID IS UFD

Every PID is a UFD.

Let  $R$  be a PID.

We want to show if  $R \in R \setminus \{0\}$ ,  $r \notin R^\times$ .

Then  $r$  admits a **unique** expression as a product of irreducibles.

## Lemma 12.4

$R$  has **some** expression as a product of irreducibles

**Proof.** If  $r$  is irreducible, then  $r = r$ .

If not, then  $r = r_1 \cdot r_2$ ,  $r_1, r_2 \notin R^\times$ . Then  $r \in (r_1)$  but  $(r) \neq (r_1)$ , therefore  $(r) \subsetneq (r_1)$ .

If  $r_1, r_2$  are irreducibles, then we are done.

If not,

$$r_1 = r_{11} \cdot r_{12}$$

$$r_2 = r_{21} \cdot r_{22}$$

where  $r_{ij} \in R^\times$ ,  $i, j \in \{1, 2\}$ . Again,  $r_1 \in (r_{11})$  but  $(r_1) \neq (r_{11})$ , therefore  $(r) \subsetneq (r_1) \subsetneq (r_{11})$ .

Since  $R$  is a PID, it is also Noetherian, and so this chain stabilizes eventually. Hence

$$r = (r_{111\dots 1} \cdot r_{111\dots 2}) \cdot \dots \cdot (r_{222\dots 1} \cdot r_{222\dots 2})$$

where each term on the right is irreducible. ■

### Lemma 12.5

The factorization into irreducibles is **unique** (up to reordering and associates).

**Proof.** Say  $r = p_1 \cdot p_2 \cdot \dots \cdot p_n$ . Let's induct on  $n$ .

If  $n = 1$ , then  $r = \underbrace{p_1}_{\text{irred.}}$  implies  $r$  is irreducible.

Suppose

$$r = q_1 \cdot q_2 \cdot \dots \cdot q_n, \quad n \geq 2, \quad q_i \text{ irreducible } \forall i \in \{1, \dots, n\}$$

But then  $q_1, (q_2 \cdot \dots \cdot q_n) \notin R^\times$  implying  $r$  is not irreducible, which is a contradiction.

Therefore  $r = r$  is the unique way to write  $r$  as the product of irreducibles.

Now suppose if  $r$  admits a factorization into at most  $n - 1$  irreducibles, then the factorization is unique. If

$$\begin{aligned} r &= p_1 \cdot p_2 \cdot \dots \cdot p_n, \quad p_i \text{'s irreducible} \\ &= q_1 \cdot q_2 \cdot \dots \cdot q_m, \quad q_j \text{'s irreducible for } m \geq n \end{aligned}$$

Then  $p_1 | q_1 \cdot (q_2 \cdot \dots \cdot q_m)$  and recall irreducibles are prime in a PID. Since  $p_1$  is irreducible either  $p_1 | q_1$  or  $p_1 | (q_2 \cdot \dots \cdot q_m)$ , so w.l.o.g assume  $p_1 | q_1$  i.e  $q_1 = u \cdot p_1$ ,  $u \in R$ .

Since  $q_1$  is irreducible, then  $u \in R^\times$  or  $p_1 \in R^\times$ . But  $p_1$  is irreducible, so it can be not an element of  $R^\times$ , therefore  $u \in R^\times$  and so  $p_1$  and  $q_1$  associate.

So we write

$$\begin{aligned} r &= p_1 \cdot p_2 \cdot \dots \cdot p_n = q_1 \cdot q_2 \cdot \dots \cdot q_m \\ &= (u \cdot p_1) \cdot q_2 \cdot \dots \cdot q_m \end{aligned}$$

Since  $R$  is an integral domain, we can cancel  $p_1$  from both sides to get

$$\underbrace{p_2 \cdot \dots \cdot p_n}_{\text{product of } (n-1) \text{ irred.}} = (u \cdot q_2) \cdot q_3 \cdot \dots \cdot q_m$$


Now by our induction hypothesis,  $r$  admits a factorization into at most  $(n - 1)$  irre-

ducibles, which implies

$$\{(u \cdot q_2), q_3, q_4, \dots, q_m\} = \{p_2, p_3, \dots, p_n\}$$

and so  $m = n$  and the  $p_i$ 's are unique. ■

We can now see a hierarchy for the specific structures we have discussed thus far



## Polynomial Rings (Again)

Let  $R$  be a commutative integral domain with  $1 \neq 0$ . Recall some facts we've already proven

- (1)  $R[X]$  is an integral domain.
- (2)  $R[X]^\times = R^\times$  e.g.  $\mathbb{Z}[X]$ , the only units are  $\{\pm 1\}$ .
- (3)  $\deg[p(X) \cdot q(X)] = \deg p(X) + \deg q(X)$
- (4) The field of fractions of  $R[X]$  is the field of rational functions

$$R(X) := \left\{ \frac{p(X)}{q(X)} \mid p, q \in R[X], q \neq 0 \right\}$$

- (5) If  $F$  is a field, then  $F[X]$  is a Euclidean Domain.

### Corollary 12.6: $F[X]$ is PID, UFD, and Noetherian

If  $F$  is a field,  $F[X]$  is a PID, UFD, and Noetherian.

- (6) Let  $I \subset R$  be an ideal, and define

$$(I) := I[X] := \{p(X) \in R[X] \mid \text{coeffs. are in } I\}$$

Then

$$R[X]/(I) \cong (R/I)[X]$$

#### **Proof.**

Consider the map

$$\begin{aligned} \phi: R[X] &\rightarrow (R/I)[X] \\ a_0 + a_1X + \cdots + a_nX^n &\mapsto \bar{a} + \bar{a}_1X + \bar{a}_2X^2 + \cdots + \bar{a}_nX^n \end{aligned}$$

for example

$$\begin{aligned} \phi: \mathbb{Z}[X] &\rightarrow (\mathbb{Z}/3\mathbb{Z})[X] \\ 1 + 2X + 4X^3 &\mapsto \bar{1} + \bar{2}X + \bar{4}X^3 = \bar{1} + \bar{2}X + X^3 \end{aligned}$$

"Clearly"  $\phi$  is a surjective ring homomorphism, so

$$(R/I)[X] \cong R[X]/\text{Ker } \phi$$

But  $\text{Ker } \phi := \{a_0 + a_1X + \cdots + a_nX^n \mid a_i \in I\} = (I)$ . ■

### Corollary 12.7

If  $I \subset R$  is prime, then  $(I) \subset R[X]$  is prime.

**Example 12.1.** Consider  $3\mathbb{Z} := \{0, 3, -3, 6, -6, \dots\}$  and

$$(3\mathbb{Z}) := \{a_0 + a_1X + a_2X^2 + \cdots + a_nX^n \mid a_i \in 3\mathbb{Z}\} \implies \mathbb{Z}[X]/(3\mathbb{Z}) \cong (\mathbb{Z}/3\mathbb{Z})[x]$$

e.g

$$1 + 2X + 4X^3 = 1 + 2X + X^3 + \underbrace{3X^3}_{\in 3\mathbb{Z}}$$

we can think about the coefficients

$$\underbrace{1, 2, 4}_{\in \mathbb{Z}} \rightarrow \underbrace{\bar{1}, \bar{2}, \bar{1}}_{\in \mathbb{Z}/3\mathbb{Z}}$$

**Theorem 12.8:**  $F[X]$  satisfies euclidean condition

If  $a(X), b(X) \in F[X]$  where  $F$  is a field. Then there exist unique  $q(X), r(X) \in F[X]$  such that  $\deg(r(X)) < \deg(b(X))$  (or  $r(X) = 0$ ) for which

$$a(X) = q(X) \cdot b(X) + r(X)$$

*Note:* Recall  $\mathbb{Z}$  are a Euclidean Domain with  $N(n) = |n|$ , e.g

$$7 = 3 \cdot 2 + 1 \quad N(1) = 1 < N(2)$$

$$7 = 4 \cdot 2 - 1 \quad N(-1) = 1 < N(2)$$

**Proof.** Suppose  $a(X) = q(X) \cdot b(X) + r(X) = q'(X)b(X) + r'(X)$ , then

$$r(X) = a(X) - q(X) \cdot b(X)$$

$$r'(X) = a(X) - q'(X) \cdot b(X)$$

and  $\deg(r), \deg(r') < \deg(b)$ .

Consider

$$r(X) - r'(X) = q'(X) \cdot b(X) - q(X) \cdot b(X) = [q'(X) - q(X)] \cdot b(X)$$

If  $q' - q, b \neq 0$ , then

$$\begin{aligned} \deg[(q' - q) \cdot b] &= \deg(q' - q) + \deg(b) \\ &= \deg[r - r'] < \deg b \end{aligned}$$

Then  $\deg q - q'$  must be 0 and so  $q' - q = 0 \implies q' = q \implies r = r'$ . ■

**Corollary 12.9**

Suppose  $F, K$  are fields with  $F \subset K$  and  $a(X), b(X) \in F[X]$ .

Then the quotient and remainder polynomials of  $a$  by  $b$  are independent of field.

**Proof.** There exist  $q(X), r(X) \in F[X]$  and  $Q(X), R(X) \in K[X]$  with  $\deg r < \deg b$  and  $\deg R < \deg b$ , such that

$$a(X) = q(X) \cdot b(X) + r(X) \quad a(X) = Q(X) \cdot b(X) + R(X)$$

But there is uniqueness since  $q, r \in K[X]$  it must mean that

$$q(X) = Q(X) \quad r(X) = R(X) \quad \text{■}$$

**Corollary 12.10**

$b(X) | a(X)$  in  $K[X]$  iff  $b(X) | a(X)$  in  $F[X]$

**Example 12.2.**

$$(X - 1) \mid X^2 - 1 \text{ in } \mathbb{R}[X], \mathbb{C}[X]$$

However,

$$(X - i) \mid X^2 + 1 \text{ in } \mathbb{C}[X] \text{ but not } \mathbb{R}[X]$$

Since  $X^2 + 1$  has no nontrivial factors in  $\mathbb{R}[X]$ .

## Polynomial Rings with Multiple Variables

### Definition 12.11: Multivariable Polynomial Ring

Let  $R$  be a commutative ring with  $1 \neq 0$ .

The **polynomial ring in the variables**  $X_1, \dots, X_n$  **with coefficients in**  $\mathbf{R}$  is defined inductively as

$$R[X_1, X_2, \dots, X_n] := R[X_1, X_2, \dots, X_{n-1}][X_n]$$

Concretely, think of  $R[X_1, \dots, X_n]$  as finite sums of **monomials**, i.e

$$aX_1^{d_1}X_2^{d_2}\dots X_n^{d_n}, \quad d_i \in \mathbb{Z}, d_i \geq 0$$

e.g

$$1 + 2XY + Y^2, 2X - 7X^3y + 2XY^4 + 1 \in \mathbb{Z}[X, Y]$$

### Definition 12.12: Multi-Degree

The **degree** of a monomial

$$aX_1^{d_1}X_2^{d_2}\dots X_n^{d_n}$$

is  $d = d_1 + d_2 + \dots + d_n$ .

The **multi-degree** is  $(d_1, d_2, d_3, \dots, d_n)$ .

The **degree** of a polynomial is the highest degree of any monomial in it.

### Proposition 12.13

Let  $R$  be an integral domain and

$$p(X_1, \dots, X_n), q(X_1, \dots, X_n) \in R[X_1, X_2, \dots, X_n] \setminus \{0\}$$

then

- (1)  $R[X_1, X_2, \dots, X_n]$  is an integral domain.
- (2)  $R[X_1, X_2, \dots, X_n]^\times = R^\times$
- (3)  $\deg[p \cdot q] = \deg p + \deg q$