

Linear Programming

Linear Programming, LP Duality Theorem, LP-Relaxation

Linear Program (LP)

- A set of linear equations/inequalities.
- Maximize or minimize a given linear objective function.

$$\text{maximize } c_1x_1 + c_2x_2 + \cdots c_nx_n$$

$$\text{subject to } a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \leq b_1$$

$$a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \leq b_2$$

$$\vdots$$

$$a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \leq b_m$$

$$x_1, x_2, \dots, x_n \geq 0$$


Example

- Suppose a factory can produce two kinds of products: oil and sugar.
- Profit for 1 tons of sugar: 1
- Profit for 1 tons of oil: 6
- Limited resources, can produce at most
 - 200 tons of sugar
 - 300 tons of oil
 - Overall weight is at most 400 tons
- Problem: maximize the profit

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$


Feasible Region

maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$




Maximizing the Objective

maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$



Maximizing the Objective


maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$



Maximizing the Objective

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$

Optimum is obtained at vertex A , where $(x_1, x_2) = (100, 300)$ and $c = 1900$.



Another Example with Three variables

maximize $x_1 + 6x_2 + 13x_3$


subject to $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 + x_3 \leq 400$

$x_2 + 3x_3 \leq 600$

$x_1, x_2, x_3 \geq 0$



Another Example with Three variables

maximize $x_1 + 6x_2 + 13x_3$


subject to $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 + x_3 \leq 400$


$x_2 + 3x_3 \leq 600$

$x_1, x_2, x_3 \geq 0$



Another Example with Three variables

maximize $x_1 + 6x_2 + 13x_3$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 + x_3 \leq 400$
 $x_2 + 3x_3 \leq 600$
 $x_1, x_2, x_3 \geq 0$



Important Observations

1. There always exists an optimum $x = (x_1, \dots, x_n)$ at a **vertex** of the polytope.
 - Linear objective $\Rightarrow c = c_1x_1 + \dots + c_nx_n$ is a **hyperplane**.
 - Optimum is obtained only when the whole feasible region is below the hyperplane and the hyperplane “barely” intersect the region by a point.
2. The feasible region is always convex.
 - Linear Constraints \Rightarrow feasible region is bounded by **hyperplanes**.
3. A local maximum is also a global maximum.
 - By the convexity of the feasible region...


Simplex Method

- Choose an arbitrary starting **vertex**.
- Iteratively move to an adjacent **vertex** along an **edge** if such movement increase the objective.
- Terminate when we reach a local maximum.

Simplex Method

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$


Starting from vertex O .



Simplex Method

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$


Moving from O to A
increases the objective.



Simplex Method

maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$


Moving from A to B
increases the objective.



Simplex Method

maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$


Moving from B to C
increases the objective.



Simplex Method

maximize $x_1 + 6x_2$
subject to $x_1 \leq 200$
 $x_2 \leq 300$
 $x_1 + x_2 \leq 400$
 $x_1, x_2 \geq 0$

C is a local maximum:
Moving to either D or B
decreases the objective.



Simplex Method

maximize $x_1 + 6x_2 + 13x_3$


subject to $x_1 \leq 200$

$x_2 \leq 300$

$x_1 + x_2 + x_3 \leq 400$

$x_2 + 3x_3 \leq 600$

$x_1, x_2, x_3 \geq 0$




Time Complexity for Simplex Method


- There are exponentially many vertices: $\binom{m}{n}$ for m constraints and n variables.
- Worst-case running time: exponential
 - Many attempts have failed.
 - e.g., choose neighbors with highest objective value, choose neighbors randomly, etc.
- [Teng & Spielman] Smoothed analysis
 - Average case polynomial time if add random Gaussian noise to the constraints.
- Runs fast in practice, and most commonly used

Polynomial Time Algorithms for LP

- Ellipsoid Method



- Interior Point Method



Standard Form LP

- Maximization as objective with " \leq " constraints and non-negative variables.

$$\begin{aligned} \text{maximize} \quad & c_1x_1 + c_2x_2 + \cdots c_nx_n \\ \text{subject to} \quad & a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \leq b_1 \\ & a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \leq b_2 \\ & \vdots \\ & a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \leq b_m \\ & x_1, x_2, \dots, x_n \geq 0 \end{aligned}$$

$$\begin{aligned} \text{maximize} \quad & \mathbf{c}^T \mathbf{x} \\ \text{subject to} \quad & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq \mathbf{0} \end{aligned}$$

Other Forms Reduce to Standard Form

- Minimization to Maximization

- $\min c_1x_1 + \cdots + c_nx_n \iff \max -c_1x_1 - \cdots - c_nx_n$

- \geq -inequalities

- $a_1x_1 + \cdots + a_nx_n \geq b \iff -a_1x_1 - \cdots - a_nx_n \leq -b$

- Inequality \iff Equality

- $a_1x_1 + \cdots + a_nx_n = b \iff \begin{cases} a_1x_1 + \cdots + a_nx_n \leq b \\ a_1x_1 + \cdots + a_nx_n \geq b \end{cases}$

- $a_1x_1 + \cdots + a_nx_n \leq b \iff a_1x_1 + \cdots + a_nx_n + s = b$

- Variable with unrestricted signs

- Introduce two variables x^+ and x^- with standard constraints $x^+, x^- \geq 0$
 - Replace x with $x^+ - x^-$

Take-Home Message

- A linear program can be solved in a polynomial time.
- Whenever a problem can be formulated by a linear program, it is polynomial-time solvable.

Formulation as Linear Program

- The **maximum flow problem** can be formulated by a linear program.

$$\begin{aligned} &\text{maximize} && \sum_{u:(s,u) \in E} f_{su} \\ &\text{subject to} && 0 \leq f_{uv} \leq c_{uv} && \forall (u,v) \in E \\ &&& \sum_{v:(v,u) \in E} f_{vu} = \sum_{w:(v,w) \in E} f_{vw} && \forall u \in V \setminus \{s,t\} \end{aligned}$$

- Ford-Fulkerson Method implements the simplex method.

Part II: LP Duality

Motivation

- We have seen that the optimal solution for the LP below is $(x_1, x_2) = (100, 300)$, with value 1900.
 - Geometric argument, argument based on simplex method
- Let's try to prove it by some simple observations from the LP itself!

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$

Motivation

$$\begin{array}{ll} \text{maximize} & x_1 + 6x_2 \\ \text{subject to} & x_1 \leq 200 \quad \text{(i)} \\ & x_2 \leq 300 \quad \text{(ii)} \\ & x_1 + x_2 \leq 400 \quad \text{(iii)} \\ & x_1, x_2 \geq 0 \end{array}$$

- Let's try adding (i) to 6 times (ii): $x_1 + 6x_2 \leq 200 + 6 \times 300 = 2000$
- We know that any solution (x_1, x_2) cannot yield objective value greater than 2000.
- Can we combine the inequality in a better way to show that the objective value is at most 1900?

Motivation

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 && \text{(i)} \\ & & x_2 \leq 300 && \text{(ii)} \\ & & x_1 + x_2 \leq 400 && \text{(iii)} \\ & & x_1, x_2 \geq 0 \end{aligned}$$

- Can we combine the inequality in a better way to show that the objective value is at most 1900?
- Yes, we can:
 - Multiple (ii) by 5 and add to (iii): $x_1 + 6x_2 \leq 300 \times 5 + 400 = 1900$.
- This proves that $(x_1, x_2) = (100, 300)$ with objective value 1900 is optimal!

Let's try this one...

- Suppose we multiple (i) by y_1 , (ii) by y_2 , (iii) by y_3 , and (iv) by y_4 .
- We have $(y_1 + y_3)x_1 + (y_2 + y_3 + y_4)x_2 + (y_3 + 3y_4)x_3 \leq 200y_1 + 300y_2 + 400y_3 + 600y_4$.
- We need $y_1, y_2, y_3, y_4 \geq 0$ to keep the inequality.
- To find an upper bound to the objective $x_1 + 6x_2 + 13x_3$, we need to make sure $x_1 + 6x_2 + 13x_3 \leq (y_1 + y_3)x_1 + (y_2 + y_3 + y_4)x_2 + (y_3 + 3y_4)x_3$ holds for every (x_1, x_2, x_3) .
- Since $x_1, x_2, x_3 \geq 0$, we must have:
 - $y_1 + y_3 \geq 1$
 - $y_2 + y_3 + y_4 \geq 6$
 - $y_3 + 3y_4 \geq 13$

$$\text{maximize } x_1 + 6x_2 + 13x_3$$

$$\text{subject to } x_1 \leq 200 \quad (\text{i})$$

$$x_2 \leq 300 \quad (\text{ii})$$

$$x_1 + x_2 + x_3 \leq 400 \quad (\text{iii})$$

$$x_2 + 3x_3 \leq 600 \quad (\text{iv})$$

$$x_1, x_2, x_3 \geq 0$$

Let's try this one...

- $(y_1 + y_3)x_1 + (y_2 + y_3 + y_4)x_2 + (y_3 + 3y_4)x_3 \leq 200y_1 + 300y_2 + 400y_3 + 600y_4.$
- Since $x_1, x_2, x_3 \geq 0$, we must have:
 - $y_1 + y_3 \geq 1$
 - $y_2 + y_3 + y_4 \geq 6$
 - $y_3 + 3y_4 \geq 13$
- Now, we want to find the tightest possible upper-bound to $x_1 + 6x_2 + 13x_3$.
- This means we want to **minimize** $200y_1 + 300y_2 + 400y_3 + 600y_4$.

$$\text{maximize } x_1 + 6x_2 + 13x_3$$

$$\text{subject to } x_1 \leq 200 \quad (\text{i})$$

$$x_2 \leq 300 \quad (\text{ii})$$

$$x_1 + x_2 + x_3 \leq 400 \quad (\text{iii})$$

$$x_2 + 3x_3 \leq 600 \quad (\text{iv})$$

$$x_1, x_2, x_3 \geq 0$$

Dual Program

- The problem of finding the tightest upper-bound can be formulated by another linear program!
- This linear program is called the **dual** program, and the original one is called the **primal** program.

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 + 13x_3 \\ &\text{subject to } x_1 \leq 200 \\ &\quad x_2 \leq 300 \\ &\quad x_1 + x_2 + x_3 \leq 400 \\ &\quad x_2 + 3x_3 \leq 600 \\ &\quad x_1, x_2, x_3 \geq 0 \end{aligned}$$

$$\begin{aligned} &\text{minimize } 200y_1 + 300y_2 + 400y_3 + 600y_4 \\ &\text{subject to } y_1 + y_3 \geq 1 \\ &\quad y_2 + y_3 + y_4 \geq 6 \\ &\quad y_3 + 3y_4 \geq 13 \\ &\quad y_1, y_2, y_3, y_4 \geq 0 \end{aligned}$$

Dual Program

- Factory Example:

$$\begin{aligned} &\text{maximize } x_1 + 6x_2 \\ &\text{subject to } x_1 \leq 200 \\ &\quad \quad \quad x_2 \leq 300 \\ &\quad \quad \quad x_1 + x_2 \leq 400 \\ &\quad \quad \quad x_1, x_2 \geq 0 \end{aligned}$$

$$\begin{aligned} &\text{minimize } 200y_1 + 300y_2 + 400y_3 \\ &\text{subject to } y_1 + y_3 \geq 1 \\ &\quad \quad \quad y_2 + y_3 \geq 6 \\ &\quad \quad \quad y_1, y_2, y_3 \geq 0 \end{aligned}$$

- Dual program for standard form:

$$\begin{aligned} &\text{maximize } \mathbf{c}^T \mathbf{x} \\ &\text{subject to } A\mathbf{x} \leq \mathbf{b} \\ &\quad \quad \quad \mathbf{x} \geq \mathbf{0} \end{aligned}$$

$$\begin{aligned} &\text{minimize } \mathbf{b}^T \mathbf{y} \\ &\text{subject to } A^T \mathbf{y} \geq \mathbf{c} \\ &\quad \quad \quad \mathbf{y} \geq \mathbf{0} \end{aligned}$$

Weak Duality Theorem

- By our motivation of dual program, we obtain the following theorem.
- Theorem [Weak Duality Theorem]. If $\hat{\mathbf{x}}$ is a feasible solution to (a) and $\hat{\mathbf{y}}$ is a feasible solution to (b), then $\mathbf{c}^T \hat{\mathbf{x}} \leq \mathbf{b}^T \hat{\mathbf{y}}$.

$$\begin{array}{ll} \text{maximize} & \mathbf{c}^T \mathbf{x} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq \mathbf{0} \end{array} \quad (a)$$

$$\begin{array}{ll} \text{minimize} & \mathbf{b}^T \mathbf{y} \\ \text{subject to} & A^T \mathbf{y} \geq \mathbf{c} \\ & \mathbf{y} \geq \mathbf{0} \end{array} \quad (b)$$



Strong Duality Theorem: This gap is always closed!

Strong Duality Theorem

- Theorem [Strong Duality Theorem]. Let \mathbf{x}^* be the optimal solution to (a) and \mathbf{y}^* be the optimal solution to (b), then $\mathbf{c}^\top \mathbf{x}^* = \mathbf{b}^\top \mathbf{y}^*$.

$$\begin{array}{ll} \text{maximize} & \mathbf{c}^\top \mathbf{x} \\ \text{subject to} & A\mathbf{x} \leq \mathbf{b} \\ & \mathbf{x} \geq \mathbf{0} \end{array} \quad (\text{a})$$

$$\begin{array}{ll} \text{minimize} & \mathbf{b}^\top \mathbf{y} \\ \text{subject to} & A^\top \mathbf{y} \geq \mathbf{c} \\ & \mathbf{y} \geq \mathbf{0} \end{array} \quad (\text{b})$$

Primal feasible

Primal OPT = Dual OPT

Dual feasible



Application of Strong Duality Theorem

- Max-Flow-Min-Cut Theorem
- Minimax Theorem
- König-Egerváry Theorem
- Design approximation algorithms:
 - Dual fitting
 - Primal-Dual Schema
- Economic interpretation: "resource allocation"- "resource valuation"

Today's Lecture

- Introduction to Linear Programming
- LP Duality Theorem


Proof of Strong Duality Theorem

Not presented in the class.
You may want to read it if you are interested.

Proof of Strong Duality Theorem

- **Theorem [Farkas Lemma].** Exactly one of the followings holds for matrix $A \in \mathbb{R}^{m \times n}$ and vector $\mathbf{b} \in \mathbb{R}^m$:
 1. There exists $\mathbf{x} \in \mathbb{R}^n$ with $\mathbf{x} \geq 0$ such that $A\mathbf{x} = \mathbf{b}$.
 2. There exists $\mathbf{y} \in \mathbb{R}^m$ such that $A^T\mathbf{y} \geq 0$ and $\mathbf{b}^T\mathbf{y} < 0$.
- $\{A\mathbf{x} \mid \mathbf{x} \geq 0\}$ is the grey area.
- 1 says that \mathbf{b} is inside the grey area.
- 2 says that we can separate the grey area and \mathbf{b} by a hyperplane (defined by the normal vector \mathbf{y}).
 - In this case \mathbf{b} must be outside the grey area.


Illustration for $A = [\mathbf{c}_1 \ \mathbf{c}_2]$



A Corollary to Farkas Lemma


- **Corollary.** Exactly one of the followings holds for matrix $A \in \mathbb{R}^{m \times n}$ and vector $\mathbf{b} \in \mathbb{R}^m$:
 1. There exists $\mathbf{x} \in \mathbb{R}^n$ with $\mathbf{x} \geq \mathbf{0}$ such that $A\mathbf{x} \geq \mathbf{b}$.
 2. There exists $\mathbf{y} \in \mathbb{R}^m$ with $\mathbf{y} \leq \mathbf{0}$ such that $A^T\mathbf{y} \geq \mathbf{0}$ and $\mathbf{b}^T\mathbf{y} < 0$.

Case 1 of the Corollary



- $\{Ax \mid x \geq 0\}$ is the dark grey area.
- $\{x \mid x \geq b\}$ is the light grey area.
- 1 says that the two areas intersect.

Case 2 of the Corollary



- $\{A\mathbf{x} \mid \mathbf{x} \geq \mathbf{0}\}$ is the dark grey area.
- $\{\mathbf{x} \mid \mathbf{x} \geq \mathbf{b}\}$ is the light grey area.
- 2 describes that the two areas do not intersect.
- We can find a separating plane with normal vector \mathbf{y} .
 - Thus, $A^T \mathbf{y} \geq \mathbf{0}$ and $\mathbf{b}^T \mathbf{y} < 0$
- We must have $\mathbf{y} \leq \mathbf{0}$:
 - If this fails for one entry: $y_i > 0$
 - $\mathbf{z} = (\varepsilon, \dots, \varepsilon, z_i = 1, \varepsilon, \dots, \varepsilon)$ and \mathbf{y} on same side
 - \mathbf{z} is in the first quadrant, and it will eventually intersect the light grey area after extension.
 - The two areas are on the same side with \mathbf{y} .

Proof of the Corollary

- Define $A' \in \mathbb{R}^{m \times (n+m)}$ by $A' = [A \ -I]$.
- Apply Farkas Lemma on A' and b .
- Let P1 and P2 be 1 and 2 in Farkas Lemma; Q1 and Q2 be 1 and 2 in the corollary.
- We aim to show $P1 \Leftrightarrow P2$ and $Q1 \Leftrightarrow Q2$.

Proof of the Corollary

- Define $A' \in \mathbb{R}^{m \times (n+m)}$ by $A' = [A \ -I]$.
- $P1 \Leftrightarrow \exists \mathbf{x}' \in \mathbb{R}^{n+m}$ s.t. $\mathbf{x}' \geq \mathbf{0}$ and $A'\mathbf{x}' = \mathbf{b}$.
- (by writing $\mathbf{x}' = \begin{bmatrix} \mathbf{x} \\ \bar{\mathbf{x}} \end{bmatrix}$) $\Leftrightarrow [A \ -I] \begin{bmatrix} \mathbf{x} \\ \bar{\mathbf{x}} \end{bmatrix} = \mathbf{b}$ (where $\mathbf{x} \geq \mathbf{0}, \bar{\mathbf{x}} \geq \mathbf{0}$)
- $\Leftrightarrow A\mathbf{x} - \bar{\mathbf{x}} = \mathbf{b} \Leftrightarrow A\mathbf{x} \geq \mathbf{b}$ (since $\bar{\mathbf{x}} \geq \mathbf{0}$)
- $\Leftrightarrow Q1$

Proof of the Corollary

- Define $A' \in \mathbb{R}^{m \times (n+m)}$ by $A' = [A \ -I]$.
- $P2 \Leftrightarrow \exists \mathbf{y} \in \mathbb{R}^m$ s.t. $A'^T \mathbf{y} \geq \mathbf{0}$ and $\mathbf{b}^T \mathbf{y} < 0$.
- $\Leftrightarrow \begin{bmatrix} A^T \\ -I \end{bmatrix} \mathbf{y} \geq \mathbf{0}$ and $\mathbf{b}^T \mathbf{y} < 0$
- $\Leftrightarrow A^T \mathbf{y} \geq \mathbf{0}, \quad -\mathbf{y} \geq \mathbf{0}, \quad \text{and } \mathbf{b}^T \mathbf{y} < 0$
- $\Leftrightarrow Q2$

Now we are ready to prove strong duality theorem...

- Weak duality: $\mathbf{c}^\top \mathbf{x} \leq \mathbf{b}^\top \mathbf{y}^*$ holds for any $\mathbf{x} \geq \mathbf{0}$.
- Suppose strong duality fails: $\mathbf{c}^\top \mathbf{x} < \mathbf{b}^\top \mathbf{y}^*$.
- There does not exist $\mathbf{x} \geq \mathbf{0}$ satisfying $A\mathbf{x} \leq \mathbf{b}$ and $\mathbf{c}^\top \mathbf{x} \geq \mathbf{b}^\top \mathbf{y}^*$.
- We cannot have $\begin{bmatrix} -A \\ \mathbf{c}^\top \end{bmatrix} \mathbf{x} \geq \begin{bmatrix} -\mathbf{b} \\ \mathbf{b}^\top \mathbf{y}^* \end{bmatrix}$ and $\mathbf{x} \geq \mathbf{0}$.
- Q1 in corollary fails for matrix $\begin{bmatrix} -A \\ \mathbf{c}^\top \end{bmatrix}$ and vector $\begin{bmatrix} -\mathbf{b} \\ \mathbf{b}^\top \mathbf{y}^* \end{bmatrix}$.
- Thus, Q2 must be true.

Now we are ready to prove strong duality theorem...

- Q2 is true for matrix $\begin{bmatrix} -A \\ \mathbf{c}^\top \end{bmatrix}$ and vector $\begin{bmatrix} -\mathbf{b} \\ \mathbf{b}^\top \mathbf{y}^* \end{bmatrix}$.

- There exist $\mathbf{y} \in \mathbb{R}^m$ and $w \in \mathbb{R}$ such that

$$\begin{bmatrix} -A^\top & \mathbf{c} \end{bmatrix} \begin{bmatrix} \mathbf{y} \\ w \end{bmatrix} \geq \mathbf{0}, \quad \begin{bmatrix} -\mathbf{b}^\top & \mathbf{b}^\top \mathbf{y}^* \end{bmatrix} \begin{bmatrix} \mathbf{y} \\ w \end{bmatrix} < 0, \quad \text{and} \quad \begin{bmatrix} \mathbf{y} \\ w \end{bmatrix} \leq \mathbf{0}.$$

- After matrix multiplications,

$$\begin{cases} -A^\top \mathbf{y} + w\mathbf{c} \geq \mathbf{0} \\ -\mathbf{b}^\top \mathbf{y} + w\mathbf{b}^\top \mathbf{y}^* < 0 \\ \mathbf{y} \leq \mathbf{0} \\ w \leq 0 \end{cases}$$

Proof of Strong Duality Theorem

$$\begin{cases} -A^T \mathbf{y} + w\mathbf{c} \geq \mathbf{0} \\ -\mathbf{b}^T \mathbf{y} + w\mathbf{b}^T \mathbf{y}^* < 0 \\ \mathbf{y} \leq \mathbf{0} \\ w \leq 0 \end{cases}$$

- Suppose $w < 0$. We divide w on both sides:

$$\begin{cases} -A^T \left(\frac{\mathbf{y}}{w} \right) + \mathbf{c} \leq \mathbf{0} \\ -\mathbf{b}^T \left(\frac{\mathbf{y}}{w} \right) + \mathbf{b}^T \mathbf{y}^* > 0 \\ \left(\frac{\mathbf{y}}{w} \right) \geq \mathbf{0} \end{cases}$$

- $\left(\frac{\mathbf{y}}{w} \right)$ is a better solution than \mathbf{y}^* in the dual LP, contradiction!

Proof of Strong Duality Theorem

$$\begin{cases} -A^T \mathbf{y} + w\mathbf{c} \geq \mathbf{0} \\ -\mathbf{b}^T \mathbf{y} + w\mathbf{b}^T \mathbf{y}^* < 0 \\ \mathbf{y} \leq \mathbf{0} \\ w \leq 0 \end{cases}$$

- Let's then do the case $w = 0$.
- We have $-A^T \mathbf{y} \geq \mathbf{0}$, $-\mathbf{b}^T \mathbf{y} < 0$, and $\mathbf{y} \leq \mathbf{0}$.
- Q2 in Corollary holds for $-A$ and $-\mathbf{b}$.
- So Q1 must be false: $\nexists \mathbf{x} \geq \mathbf{0}: (-A)\mathbf{x} \geq -\mathbf{b}$.
- The feasible region for the primal LP is empty!