

From Lowe Syndrome to Dent Disease: Correlations between Mutations of the *OCRL1* Gene and Clinical and Biochemical Phenotypes

Haifa Hichri,¹ John Rendu,^{1,2} Nicole Monnier,^{1,2} Charles Coutton,³ Olivier Dorseuil,⁴ Rosa Vargas Poussou,⁵ Geneviève Baujat,⁶ Anne Blanchard,⁷ François Nobili,⁸ Bruno Ranchin,⁹ Michel Remesy,¹⁰ Rémi Salomon,¹¹ Véronique Satre,³ and Joel Lunardi^{1,2*}

¹CHU Grenoble, Laboratoire de Biochimie et Génétique Moléculaire, Grenoble, France; ²INSERM U836, Grenoble Institut des Neurosciences, Grenoble, France; ³CHU Grenoble, Laboratoire de Génétique Chromosomique, Grenoble, France; ⁴Institut Cochin, Département de Génétique, Développement et Pathologie Moléculaire, INSERM U567-CNRS UMR8104-Université Paris V, Paris, France; ⁵Hôpital Européen Georges Pompidou, Département de Génétique et Inserm U 970, Paris, France; ⁶Hôpital Necker-Enfants Malades, Département de Génétique, Paris, France; ⁷Hôpital Européen Georges Pompidou, Centre d'Investigation Clinique, Paris, France; ⁸CHU Besançon, Néphrologie Pédiatrique, Besançon, France; ⁹Hôpital Femme, Mère, Enfant, Hospices Civils de Lyon, Centre de Référence des Maladies, Rénales Rares, Bron, France; ¹⁰CHU Toulouse, Hôpital des Enfants, Service de Néphrologie, Toulouse, France; ¹¹Hôpital Necker-Enfants Malades, Centre de Référence des Maladies Rénales; Héritaires de l'Enfant et de l'Adulte, Paris, France

Communicated by Arnold Munch

Received 6 August 2010; accepted revised manuscript 6 October 2010.

Published online 28 October 2010 in Wiley Online Library (www.wiley.com/humanmutation). DOI 10.1002/humu.21391

ABSTRACT: Mutations of *OCRL1* are associated with both the Lowe oculocerebrorenal syndrome, a multi-systemic and Dent-2 disease, a renal tubulopathy. We have identified a mutation in 130 Lowe syndrome families and 6 affected by Dent-2 disease with 51 of these mutations being novel. No founding effect was evidenced for recurrent mutations. Two mutations initially reported as causing Dent-2 disease were identified in patients, including two brothers, presenting with Lowe syndrome thus extending the clinical variability of *OCRL1* mutations. mRNA levels, protein content, and PiP₂-ase activities were analyzed in patient's fibroblasts. Although mRNA levels were normal in cells harboring a missense mutation, the *OCRL1* content was markedly lowered, suggesting that enzymatic deficiency resulted mainly from protein degradation rather than from a catalytic inactivation. Analysis of a splicing mutation that led to the elimination of the initiation codon evidenced the presence of shortened forms of *OCRL1* that might result from the use of alternative initiation codons. The specific mapping of the frameshift and nonsense mutations, exclusively identified in exons 1–7 and exons 8–23, respectively, for Dent disease and Lowe syndrome together with the possible use of alternative initiation codons might be related to their clinical expression, that is, Lowe syndrome or Dent-2 disease.

Hum Mutat 32:379–388, 2011. © 2010 Wiley-Liss, Inc.

KEY WORDS: *OCRL1*; Lowe syndrome; Dent 2 disease; phosphatidylinositol 4, 5 biphosphate homeostasis

Introduction

Lowe Syndrome

The oculocerebrorenal syndrome of Lowe (*OCRL*; MIM# 309000) is a rare X-linked multisystem disorder presenting with major abnormalities in the eyes, the kidneys, and the central nervous system [Loi, 2006; Lowe et al., 1952]. *OCRL* is a rare disease with a prevalence estimated between 1 and 2 boys per million people. Ocular abnormalities include a constant prenatal development of cataracts and frequent associated signs such as glaucoma, microphthalmos, decreased visual acuity, and corneal keloid formation. Neonatal hypotonia, intellectual impairment, and areflexia are also cardinal features. The majority of patients have a cognitive delay and behavioral troubles including temper tantrums and aggressiveness are frequently noted. Brain magnetic resonance imaging (MRI) may show periventricular cystic lesions [Loi, 2006; Schneider et al., 2001]. Fanconi syndrome, a generalized impairment of the proximal tubular cells functions, is a major feature.

Onset of the tubular dysfunction can vary between patients, and the severity tends to worsen with age. Low molecular weight proteinuria (LWMP) is invariably present and aminoaciduria, hypercalciuria, and bicarbonaturia are frequently included. Progressive glomerular dysfunction leads usually to renal failure. Skeletal muscle abnormalities may develop as secondary consequences of hypotonia or renal dysfunction. Nontender joint swelling and subcutaneous nodules are also frequently described in affected patients and may reflect a primary abnormality of connective tissue growth. Lowe syndrome results from mutations of the *OCRL1* gene (MIM# 300535) that encodes a phosphatidylinositol 4,5 biphosphate (PI(4,5)P₂) phosphatase.

Dent Disease

Dent disease (Dent-1; MIM# 300009), is a X-linked proximal renal tubulopathy, characterized by LWMP, hypercalciuria, and

Additional Supporting Information may be found in the online version of this article.

*Correspondence to: Joel Lunardi, Laboratoire de Biochimie et Génétique Moléculaire, Inserm U836, CHU Grenoble BP217X, 38043 Grenoble Cedex, France. E-mail: jlunardi@chu-grenoble.fr