

SUMMARY KEYWORDS

leatherback turtles, dives, leather

bags, turtles, jellyfish, animals, leatherback, diving, recorded, nested, scientists, tissue, layer, female, deep diver, surface, sea, leathery skin, captivity, migrates

SPEAKERS

Speaker 1 (100%)

1

Speaker 1

0:03

Secrets are called leatherback turtle. Jenny the mannequin of Florida Atlantic University has shown that the advantages of leatherback streamlined shape. Swimmy, leatherback turtles have a lower drag more efficient than other species, indicating more efficient the force to swim at the same distance to other sea turtles of the comparable size and weight. leatherback turtles spend on leverage 20% less energy among all the deep diving animals. The champions have always been assumed to be the marine mammals, the great whales and also CEOs. Our recent investigations however, suggest that leatherback sea turtles may also be ranked among the oceans greatest air breathing diver reptiles. Well Mary the dives of the leatherback sea turtles near St. Crocs. In the US Virgin Islands, scientists recorded as 650 pounds of female that sank to more than 3330 feet and remained there for 37 minutes. High pressure also poses a challenge to deep diver in terms of strong forces could compress the chest, causing broken bones or tissue damage online is hot shout relatives leatherbacks has soft a shell made up from widely separated the ribs and buried in a thick tissue and whole structure is overlaid with a leathery skin. How do the deep diving animals whales CEOs of penguins and leatherback turtles avoid a decompression sickness and other hazards while they dive to the great depths even sleep for long period of time. Many of the deepest diving marine mammals have a small loss and forcefully exhale before diving. In addition to reducing the belly and sea so that they can dive more easily. This severely limits the amount of nitrogen in their bloodstream. One reason the leatherback ranges so far, maybe specialized the diet. This giant of the sea fleas mainly on jellyfish full of a high protein, and the other useful minerals that require during the day I was on a variety of

jellyfish and related prey are abandoned. They're eaten by few other animals, as one might suspect. A lifespan eating mostly jellyfish requires some special adaptations to make the job easier. During the 1994 and 1985 nesting seasons, with the assistance of scientists the record is to 10 Female leatherbacks had just laid eggs. As a result, we were able to monitor the pattern of migration in Atlantic Ocean when they returned to the sea. Some of the longest travel recorded for many animal have been made by this marine turtle, a 9070. A female leatherback was tagged when she nested in Cerami on the northeast coast of South America. Less than one year later, she was captured on the coast of Ghana, West Africa, having traveled 3700 miles across the Atlantic Ocean. Upon returning to the surface, the Turtles change a quick breath and immediately heading straight down again. Later on no time was given to sleep or rest. We then begin to track and track the turtles when they swim back to the surface. On closer examination, we noticed a difference in dives depending on the time of the day, nine dives are shallower, with a less in depth than day dives. Leatherbacks are such swimmers, they rarely stop moving a behavior that has made it impossible to tip them in captivity. The fur on the flippers are more than half the length of those on the body and the generated power from huge muscles, which can account for 30% of the animal's total body weight. Says leatherbacks cannot easily be studied in captivity, researchers have been developing methods of studying them and see the use the recorder and the instrument capable of recording location and adapts in C. Developed by Gerald acumen on psychological research laboratory. The device has been used to study diving marine animals. Why they keep diving scientists as soon realized that the turtles were probably costing much of the energy to follow their food source. In tropical waters. Jellyfish are most common at Creek gaps in a biological zone called the diapausing layer, described shortly after development of the Sona this turns consists of layer of zooplankton, the Hoover's below 1800 feet during the day and migrates to the surface and night