

# NG and ADF test (excess return)

*Shuofan Zhang*

*December 06, 2018*

## Data

Data: 131-8=123 series in total, 480 observations, from Jan 1964 to Dec 2003.

The “spread” series (difference between two I(1) series) are removed.

log() is done.

ADF test suggest I(0) for all four bond returns, whether use “trend” or “drift” specification.

## Lasso 1

I(2) is first differenced, others are original.

$$\begin{aligned} y_t = & I(0)_{t-1} + I(0)_{t-2} + I(0)_{t-3} + I(0)_{t-4} \\ & + I(1)_{t-1} + I(1)_{t-2} + I(1)_{t-3} + I(1)_{t-4} \\ & + \Delta I(2)_{t-1} + \Delta I(2)_{t-2} + \Delta I(2)_{t-3} + \Delta I(2)_{t-4} \end{aligned}$$

# Lasso 2


All stationary.

$$\begin{aligned} y_t = & I(0)_{t-1} + I(0)_{t-2} + I(0)_{t-3} + I(0)_{t-4} \\ & + \Delta I(1)_{t-1} + \Delta I(1)_{t-2} + \Delta I(1)_{t-3} + \Delta I(1)_{t-4} \\ & + \Delta^2 I(2)_{t-1} + \Delta^2 I(2)_{t-2} + \Delta^2 I(2)_{t-3} + \Delta^2 I(2)_{t-4} \end{aligned}$$

# Lasso 3

Combination of Lasso 1 and 2.

$$\begin{aligned} y_t = & I(0)_{t-1} + I(0)_{t-2} + I(0)_{t-3} + I(0)_{t-4} \\ & + \Delta I(1)_{t-1} + \Delta I(1)_{t-2} + \Delta I(1)_{t-3} + \Delta I(1)_{t-4} \\ & + \Delta^2 I(2)_{t-1} + \Delta^2 I(2)_{t-2} + \Delta^2 I(2)_{t-3} + \Delta^2 I(2)_{t-4} \\ & + I(1)_{t-1} + \Delta I(2)_{t-1} \end{aligned}$$



Coefficients trace plot

