


Gaussian generative models

Topics we'll cover

- ① Classification using multivariate Gaussian generative modeling
- ② The form of the decision boundaries


Back to the winery data

Go from 1 to 2 features: test error goes from 29% to 8%.



With all 13 features: test error rate goes to zero.

The multivariate Gaussian



$N(\mu, \Sigma)$: Gaussian in \mathbb{R}^d

- mean: $\mu \in \mathbb{R}^d$
- covariance: $d \times d$ matrix Σ

Density
$$p(x) = \frac{1}{(2\pi)^{d/2} |\Sigma|^{1/2}} \exp \left(-\frac{1}{2} (x - \mu)^T \Sigma^{-1} (x - \mu) \right)$$

If we write $S = \Sigma^{-1}$ then S is a $d \times d$ matrix and

$$(x - \mu)^T \Sigma^{-1} (x - \mu) = \sum_{i,j} S_{ij} (x_i - \mu_i) (x_j - \mu_j),$$

a **quadratic function** of x .

Binary classification with Gaussian generative model

- Estimate class probabilities π_1, π_2
- Fit a Gaussian to each class: $P_1 = N(\mu_1, \Sigma_1)$, $P_2 = N(\mu_2, \Sigma_2)$

Given a new point x , predict class 1 if

$$\pi_1 P_1(x) > \pi_2 P_2(x) \Leftrightarrow x^T M x + 2w^T x \geq \theta,$$

where:

$$M = \frac{1}{2}(\Sigma_2^{-1} - \Sigma_1^{-1})$$
$$w = \Sigma_1^{-1}\mu_1 - \Sigma_2^{-1}\mu_2$$

and θ is a threshold depending on the various parameters.


Linear or **quadratic** decision boundary.

Common covariance: $\Sigma_1 = \Sigma_2 = \Sigma$


Linear decision boundary: choose class 1 if

$$x \cdot \underbrace{\Sigma^{-1}(\mu_1 - \mu_2)}_w \geq \theta.$$


Example 1: Spherical Gaussians with $\Sigma = I_d$ and $\pi_1 = \pi_2$.



Example 2: Again spherical, but now $\pi_1 > \pi_2$.



Example 3: Non-spherical.



Classification rule: $w \cdot x \geq \theta$

- Choose w as above
- Common practice: fit θ to minimize training or validation error


Different covariances: $\Sigma_1 \neq \Sigma_2$

Quadratic boundary: choose class 1 if $x^T M x + 2w^T x \geq \theta$, where:


$$M = \frac{1}{2}(\Sigma_2^{-1} - \Sigma_1^{-1})$$

$$w = \Sigma_1^{-1}\mu_1 - \Sigma_2^{-1}\mu_2$$


Example 1: $\Sigma_1 = \sigma_1^2 I_d$ and $\Sigma_2 = \sigma_2^2 I_d$ with $\sigma_1 > \sigma_2$



Example 2: Same thing in 1-d. $\mathcal{X} = \mathbb{R}$.



Example 3: A parabolic boundary.



Multiclass discriminant analysis

k classes: weights π_j , class-conditional densities $P_j = N(\mu_j, \Sigma_j)$.

Each class has an associated **quadratic** function

$$f_j(x) = \log(\pi_j P_j(x))$$

To classify point x , pick $\arg \max_j f_j(x)$.

If $\Sigma_1 = \dots = \Sigma_k$, the boundaries are **linear**.

