

```

1  /*
2      | Point Class |
3      Desc: A generic point class with some helper funcs
4      Source: KawakiMeido
5      State: Untested and VERY buggy lmao
6  */
7
8  #define X real()
9  #define Y imag()
10
11 template <typename T>
12 class Point {
13 public:
14     static constexpr double EPS = 1e-6;
15
16     std::complex<T> p;
17
18     // Constructors
19     Point(T x = 0, T y = 0) : p(x, y) {}
20     explicit Point(const std::complex<T>& val) : p(val) {}
21
22     // Accessors
23     // T real() { return p.real(); }
24     T real() const { return p.real(); }
25     // T imag() { return p.imag(); }
26     T imag() const { return p.imag(); }
27
28     void setX(int x) {
29         p.real(x);
30     }
31
32     void setY(int y) {
33         p.imag(y);
34     }
35
36     // Comparisons
37     bool operator==(const Point& other) const {
38         if constexpr (std::is_floating_point_v<T>) {
39             return (std::abs(p.real() - other.p.real()) < EPS) &&
40                 (std::abs(p.imag() - other.p.imag()) < EPS);
41         } else {
42             return p == other.p;
43         }
44     }
45     bool operator!=(const Point& other) const { return !(*this == other); }
46
47     // Arithmetics
48     Point& operator+=(const Point& other) { p += other.p; return *this; }
49     Point& operator-=(const Point& other) { p -= other.p; return *this; }

```

```
50     friend Point operator+(Point a, const Point& b) { a += b; return a; }
51     friend Point operator-(Point a, const Point& b) { a -= b; return a; }
52
53     // Helper Functions
54     friend T dot(const Point& a, const Point& b)    { return (std::conj(a.p) *
55 b.p).real(); }
56     friend T cross(const Point& a, const Point& b) { return (std::conj(a.p) *
57 b.p).imag(); }
58     friend T sqdist(const Point& a, const Point& b){ return std::norm(a.p -
59 b.p); }
60     friend T dist(const Point& a, const Point& b)  { return std::abs(a.p - b.p);
61 }
62     friend long double angle(const Point& a, const Point& b) { return
63 std::arg(b.p - a.p); }
64     friend long double slope(const Point& a, const Point& b) { return
65 std::tan(std::arg(a.p - b.p)); }
66 };
```