


Transport of Respiratory Gases

Transport of Oxygen in Blood



Air \rightarrow Lungs \rightarrow Blood (RBC) \rightarrow Aerobic tissues

- RBC \rightarrow Chemical union
97-98%
- Blood Plasma \rightarrow Dissolved
2-3%

EXTERNAL RESPIRATION

Pulmonary Artery
(P_{CO_2} 45 mm Hg)

Pulmonary Veins
(P_{O_2} 100 mm Hg)

Systemic Veins

Systemic Arteries

INTERNAL RESPIRATION


$P_{O_2} = 40$ mm Hg $\downarrow\downarrow$
 $P_{CO_2} = 45$ mm Hg $\uparrow\uparrow$

AFFINITY of Hb:



$CO > CO_2 > O_2$

The same videos and lecture notes provided by Physics In Seconds are for educational and informational purposes only and are not to be used for any commercial purpose. By accessing and using these materials, you agree to use them solely for personal, non-commercial use and will not hold the copyright holder liable for any damages.

By accessing and using the materials, you also agree to abide by all local copyright laws.



RED BLOOD CELL




\rightarrow Bicarbonate $\rightarrow 70\%$

\rightarrow CO₂ more soluble than O₂

CO_2 → Bicarbonate — 70%
 → Blood Plasma — 7%
 → Carb-Amino Hb (23%)

Gaseous Exchange at Cell Levels


→ CO_2 more soluble than O_2 .
 → CO_2 is more important.
 → P.P of O_2 in Blood ↓ stimulates CO_2




Harnsburger phenomenon (chloride shift)

H^+ ↑↑
 O_2 - release

Gaseous Exchange at Lungs



ALVEOLI



CHLORIDE SHIFT/
Harnsburger Phenomenon

The course videos and lecture notes provided by Physics In Seconds are for educational and informational purposes only and protected by local copyright laws.

Unauthorized reproduction or distribution is strictly prohibited. By accessing and using these materials, you agree to use them solely for personal, non-commercial use and will not hold the copyright holder liable for any damages.

By accessing and using the materials, you also agree to abide by all local copyright laws.

O_2 → Pressure derived
 Max. O_2 → 20 ml / 100 ml (100% Saturation)
 115 mm Hg → 19.6 ml / 100 ml (98% Saturation)

$$\frac{19.6}{20} \times 100 = 98\%$$

100 ml of Blood → 15 gm of Hb (1g Hb → 1.34 ml O_2)
 → $15 \times 1.34 = 20 \text{ ml}$

VENOUS BLOOD

→ 14.4 ml of O_2
 → 54 ml of CO_2
 Pulmonary Artery

ARTERIAL BLOOD

19.6 ml O_2 / 100 ml
 50 ml CO_2 / 100 ml
 Pulmonary Vein

→ 54 ml of CO_2

Pulmonary Artery

50 ml CO_2 / 100 ml

Pulmonary Vein

FACTORS

- CO_2 ↑↑
- Temp. ↑↑
- pH ↑↑
- H^+ concentration ↑↑

O_2 - CARRYING CAPACITY

- ↓↓
- ↓↓
- ↑↑
- ↓↓

O_2 - RELEASE

- ↑↑
- ↑↑
- ↓↓
- ↑↑

HIGH METABOLIC RATE

- Muscles → O_2 ↑↑
- Production of ATP ↑↑
- Large no. of O_2 dissociate from Hb.
- O_2 release ↑↑ → Tissue level

LOW METABOLIC RATE

- Less O_2 utilization
- Production of ATP ↓↓
- Less O_2 molecule dissociate.

The course videos and lecture notes provided by Physics In Seconds are for educational and informational purposes only and protected by local copyright laws.

Unauthorised reproduction or distribution is strictly prohibited. By accessing and using these materials, you agree to use them solely for personal, non-commercial use and will not hold the copyright holder liable for any damages.

By accessing and using the materials, you also agree to abide by all local copyright laws.