




# W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere<sup>\*1</sup>, Daniel Argüeso<sup>2</sup>, and Andrea Zonato<sup>3</sup>

<sup>1</sup> Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany <sup>2</sup> Physics Department, University of the Balearic Islands, Palma, Spain <sup>3</sup> Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Editor Name](#) 

Submitted: 01 January XXXX

Published: 01 January XXXX

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

An important objective of WUDAPT, the World Urban Database and Access Portals Tools community project, is to 1) to acquire and make accessible coherent and consistent information on form and function of urban morphology relevant to climate weather, and environment studies, and 2) to provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

The Python-based WUDAPT-to-WRF (W2W) package is developed in this context, and translates Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model. It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides a more simple, efficient and improved procedure to use LCZ information in WRF. Some important changes include a direct manipulation of the geogrid files (without the creation of temporary files), and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

## Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue, because of two recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the online LCZ Generator (Demuzere et al., 2021), and 2) WRF versions > 4.3 (Skamarock et al., 2021) are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato et al. (Under Review) and Zonato & Chen (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here, so as to make the translation of LCZ-based parameters better and more simple.

<sup>\*</sup>corresponding author

## Initial data requirements

In order to use the tool, two input files are required:

1. A **geo\_em.d0X** (.nc) file for the inner WRF model domain in which one would like to use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool as part of the WRF Preprocessing System (WPS), without additional modifications of the standard procedure.
2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of the `geo_em.d0X.nc` file. There are a number of ways to obtain an LCZ map for your region of interest (ROI):
  - Extract your ROI from the continental-scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et al., 2020) (see [here](#) for more info).
  - Check if your ROI is already covered by the many LCZ maps available in the [submission table](#) of the LCZ Generator.
  - Use the [LCZ Generator](#) to make an LCZ map for your ROI. See also [here](#) for more information.

## Workflow

The goal of the Python-based W2W tool is to obtain a WRF domain file (`geo_em.d0X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters relevant for all urban parameterizations embedded in WRF: the single layer urban canopy model Noah/SLUCM (Kusaka et al. (2001)), the Building Environment Parameterization (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al. (2010)).

To get to that point, a number of sequential steps are followed:

- *Step 1: Remove the default urban land cover*

The default urban land cover from MODIS is replaced with the dominant surrounding vegetation category, as is done in Li et al. (2020). This procedure affects WRF's parameters `LU_INDEX`, `LANDUSEF` and `GREENFRAC`. `LU_INDEX` is selected as the dominant category from the *nlus* (default = 45) nearest grid points (excluding ocean, urban and lakes). `LANDUSEF` and `GREENFRAC` are calculated as the mean over all grid points with that category among the *nlus* nearest points. @ DANIEL: CORRECT??

Resulting output: **geo\_em.d0X\_NoUrban.nc**

- *Step 2: Define the LCZ-based urban extent*

LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF resolution. Areas with `FRC_URB2D` < 0.2 (*frc*) are currently considered non-urban. This choice has been made to avoid to employ the urban schemes in areas where the majority of the landuse is vegetated, since the impact of the impervious surfaces is low. The `FRC_URB2D` field is also used to mask all other urban parameter fields, so that their extent is consistent.

Resulting output: **geo\_em.d0X\_LCZ\_extent.nc**

- *Step 3: Introduce modal built LCZ classes*

For each WRF grid cell, the mode of the underlying built LCZ classes is added to LU\_INDEX (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default considers LCZ classes 1-10 as built classes (*bc*). Sometimes, also LCZ E (or 15 - Bare rock or paved) can be considered as a built LCZ classes, as it might reflect large asphalt surfaces such as big parking lots or airstrips. In that case, make sure to set the *bc* argument appropriately.

▪ *Step 4: Assign urban canopy parameters*

Two pathways are followed when assigning the various urban canopy parameters to the LCZ map, and translating this information onto WRF's grid:

**Pathway 1: Morphological parameters** are assigned directly to the high-resolution LCZ map, and are afterwards aggregated to the lower-resolution WRF grid. In this way, the method produces a unique urban morphology parameter value for each WRF grid cell. This was found to be more efficient in reproducing urban boundary layer features, especially in the outskirts of the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#) routine (Varentsov et al., 2020).

Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv`, and are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014). In addition:

- Building width (BW), available in `LCZ_UCP_default.csv`, is taken from `URBPARAM_LCZ.TBL` (stored in WRF's run/ folder).
- While `URBPARAM_LCZ.TBL` also has values on street width, W2W derives street width from the mean building height (`MH_URB2D`) and the Height-to-Width ratio (`H2W`), to have these fields consistent.
- Plan (`LP_URB2D`), frontal (`LF_URB2D`) and total (`LB_URB2D`) area indices are based on formulas in Zonato et al. (2020).
- `HI_URB2D` is obtained by fitting a bounded normal distribution to the minimum (`MH_URB2D_MIN`), mean (`MH_URB2D`), and maximum (`MH_URB2D_MAX`) building height, as provided in `LCZ_UCP_default.csv`. The building height standard deviation is also required, and is approximated as  $(MH\_URB2D\_MAX - MH\_URB2D\_MIN) / 4$ .
- For computational efficiency, `HI_URB2D` values lower than 5% were set to 0 after resampling, the remaining `HI_URB2D` percentages are re-scaled to 100%.

**Pathway 2:** In line with the former Fortran-based W2W procedure, **radiative and thermal parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell (see *Step 3*). These parameter values are not stored in the netcdf output, but are read from `URBPARAM_LCZ.TBL` and assigned automatically to the modal LCZ class when running the model.

▪ *Step 5: Adjust global attributes*


In a final step, some global attributes are adjusted in the resulting netcdf files:

- `NBUI_MAX` is added as a global attribute, reflecting the maximum amount of `HI_URB2D` classes that are not 0 across the model domain. This parameter can be used when compiling WRF, to optimize memory storage.
- `NUM_LAND_CAT` is set to 41, to reflect the addition of 10 (or 11) built LCZ classes. This is not only done for the highest resolution domain file (e.g. `d04`), but also for **all of its parent domain files (e.g. `d01`, `d02`, `d03`)**. As such, make sure these files are also available in the input data directory.

Resulting output: `geo_em.d0X_LCZ_params.nc`

## Integration in WRF's preprocessing

The current tool is designed to work with the `geo_em.d0X` files produced by the `geogrid.exe`, in the context of the WRF Preprocessing System (WPS). The user should standardly run the `geogrid.exe`, which provide the various `geo_em.d0X.nc` containing the static data fields. No additional variable are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL`. The `w2w` tool reads the standard `geo_em.d0X.nc` files (for all the domains) and produces the aforementioned **`geo_em.d0X_LCZ_params.nc`**. The user should then simply rename those files with its standard name for each domain, respectively. Figure (i don't know hoy to ref in github) shows the modified workflow to set-up and run a WRF simulations including urban parameters derived from LCZs.



## Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_NoUrban.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019), Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2021)).

## Important notes

- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the values according to the characteristics of your ROI.

148     ▪ It is advised to use this tool with urban parameterization options BEP or BEP+BEM  
 149       (sf\_urban\_physics = 2 or 3). In case you use this tool with the SLUCM model  
 150       (sf\_urban\_physics = 1), make sure your lowest model level is above the highest  
 151       building height. If not, real.exe will provide the following error message: ZDC + ZOC +  
 152       2m is larger than the 1st WRF level - Stop in subroutine urban - chan  
 153       ge ZDC and ZOC

## 154 Acknowledgements

155 We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez  
 156 Sanchez and Oscar Brousse.

## 157 References

- 158 Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of  
 159 heat island and thermal comfort over an urban airshed in India under varying land use/land  
 160 cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- 161 Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient  
 162 land use producing data tool for mesoscale models? Integration of urban LCZ in WRF  
 163 over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- 164 Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O.,  
 165 Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley,  
 166 M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018).  
 167 WUDAPT: An urban weather, climate, and environmental modeling infrastructure for  
 168 the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924.  
 169 <https://doi.org/10.1175/BAMS-D-16-0236.1>
- 170 Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate  
 171 zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- 172 Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining ex-  
 173 pert and crowd-sourced training data to map urban form and functions for the continental  
 174 US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- 175 Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to  
 176 Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- 177
- 178 Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing  
 179 detailed urban canopy parameters for mesoscale climate modelling: a comparison between  
 180 WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*,  
 181 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- 182 Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler,  
 183 W. (2021). Resolving the influence of local flows on urban heat amplification during  
 184 heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>
- 185
- 186 Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Ur-  
 187 ban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab  
 188 Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>
- 189

- Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X.,  
Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall  
Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>
- Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to con-  
temporary land-use datasets across the contiguous united states using dynamical down-  
scaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>
- Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WU-  
DAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. [https://www.wudapt.org/wp-content/  
uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.  
pdf%20\(Accessed%20on%2011%20August%202021](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)
- Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parame-  
terisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>
- Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into  
WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged,  
Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>
- Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M., &  
Niyogi, D. (2021). Largescale Urban Impacts on a European Heat Wave Event. *Journal  
of Geophysical Research : Atmospheres (Under Review)*, 1–33.
- Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy  
rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban  
Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved  
WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*,  
248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model  
coupled with an urban canopy parameterization for urban climate simulations—part I.  
formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied  
Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>
- Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W.,  
Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of  
the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric  
Research. <https://doi.org/10.5065/1dfh-6p97>
- Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies.  
*Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>
- Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the 'local climate  
zone' scheme using temperature observations and model simulations [Journal Article].  
*International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>
- Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters  
on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>
- Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang,  
R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on



- 238 WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018),  
239 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- 240 Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones,*  
241 *Mitigation Strategies, building materials permeability and new buildings drag coefficient.*  
242 <http://www.wudapt.org/wudapt-to-wrf/>. [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)  
243 [product-tool/urban-canopy-model/WRF\\_urban\\_update\\_Readme\\_file\\_WRF4.3.pdf](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
- 244 Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the  
245 performance of a novel WUDAPT averaging technique to define urban morphology with  
246 mesoscale models. *Urban Climate*, 31(May 2019), 100584. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.uclim.2020.100584)  
247 [uclim.2020.100584](https://doi.org/10.1016/j.uclim.2020.100584)
- 248 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., & Barlage, M. (Under Review).  
249 Exploring the effects of rooftop mitigation strategies on urban temperatures and energy  
250 consumption. *Journal of Advances in Modeling Earth Systems*, 1–29.

DRAFT