




# W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere<sup>\*1</sup>, Daniel Argüeso<sup>2</sup>, and Andrea Zonato<sup>3</sup>

<sup>1</sup> Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany <sup>2</sup> Physics Department, University of the Balearic Islands, Palma, Spain <sup>3</sup> Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

## Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Editor Name](#) 

Submitted: 01 January XXXX

Published: 01 January XXXX

## License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

## Summary

An important objective of WUDAPT, the World Urban Database and Access Portals Tools community project, is to 1) to acquire and make accessible coherent and consistent information on form and function of urban morphology relevant to climate weather, and environment studies, and 2) to provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

The Python-based WUDAPT-to-WRF (W2W) package is developed in this context, and translates Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model. It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides a more simple, efficient and improved procedure to use LCZ information in WRF.

## Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue, because of two recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the online LCZ Generator (Demuzere et al., 2021), and 2) WRF versions > 4.x (Skamarock et al., 2021) are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato et al. (Under Review) and Zonato & Chen (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here, so as to make the translation of LCZ-based parameters better and more simple.

## Initial data requirements

In order to use the tool, two input files are required:

<sup>\*</sup>corresponding author

1. A **geo\_em.d0X** (.nc) file for the inner WRF model domain in which one would like to use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool as part of the WRF Preprocessing System (WPS). **@ ANDREA: does a user needs to use specific settings here to create this file?? Please extend this section if needed.**
2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of the `geo_em.d0X.nc` file. There are a number of ways to obtain an LCZ map for your region of interest (ROI):
  - Extract your ROI from the continental-scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et al., 2020) (see [here](#) for more info).
  - Check if your ROI is already covered by the many LCZ maps available in the [submission table](#) of the LCZ Generator.
  - Use the [LCZ Generator](#) to make an LCZ map for your ROI. See also [here](#) for more information.

## Workflow

The goal of the Python-based W2W tool is to obtain a WRF domain file (`geo_em.d0X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters relevant for all urban parameterizations embedded in WRF: the single layer urban canopy model Noah/SLUCM (Kusaka et al. (2001)), the Building Environment Parameterization (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al. (2010)).

To get to that point, a number of sequential steps are followed:

- *Step 1: Remove the default urban land cover*

The default urban land cover from MODIS is replaced with the dominant surrounding vegetation category, as is done in Li et al. (2020). This procedure affects WRF's parameters `LU_INDEX`, `LANDUSEF` and `GREENFRAC`. `LU_INDEX` is selected as the dominant category from the *nlus* (default = 45) nearest grid points (excluding ocean, urban and lakes). `LANDUSEF` and `GREENFRAC` are calculated as the mean over all grid points with that category among the *nlus* nearest points. **@ DANIEL: CORRECT??**

Resulting output: **geo\_em.d0X\_NoUrban.nc**

- *Step 2: Define the LCZ-based urban extent*

LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF resolution. Areas with `FRC_URB2D` < 0.2 (*frc*) are currently considered non-urban **@ ANDREA - ADD SMALL SENTENCE TO STATE WHY THAT IS**. The `FRC_URB2D` field is also used to mask all other urban parameter fields, so that their extent is consistent.

Resulting output: **geo\_em.d0X\_LCZ\_extent.nc**

- *Step 3: Introduce modal built LCZ classes*

For each WRF grid cell, the mode of the underlying built LCZ classes is added to `LU_INDEX` (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default considers LCZ classes 1-10 as built classes (*bc*). Sometimes, also LCZ E (or 15 - Bare rock or paved) can be considered as a built LCZ classes, as it might reflect large asphalt surfaces such as big parking lots or airstrips. In that case, make sure to set the *bc* argument appropriately.

76     ▪ *Step 4: Assign urban canopy parameters*

77     Two pathways are followed when assigning the various urban canopy parameters to the LCZ  
78     map, and translating this information onto WRF's grid:

79     **Pathway 1: Morphological parameters** are assigned directly to the high-resolution LCZ  
80     map, and are afterwards aggregated to the lower-resolution WRF grid. In this way, the  
81     method produces a unique urban morphology parameter value for each WRF grid cell. This  
82     was found to be more efficient in reproducing urban boundary layer features, especially in  
83     the outskirts of the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#)  
84     routine (Varentsov et al., 2020).

85     Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv`, and  
86     are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014).  
87     In addition:

- 88     ▪ Building width (BW), available in `LCZ_UCP_default.csv`, is taken from `URBPARAM_LCZ.TBL` (stored in WRF's `run/` folder).
- 89     ▪ While `URBPARAM_LCZ.TBL` also has values on street width, `W2W` derives street width from the mean building height (`MH_URB2D`) and the Height-to-Width ratio (`H2W`), to have these fields consistent.
- 90     ▪ Plan (`LP_URB2D`), frontal (`LF_URB2D`) and total (`LB_URB2D`) area indices are based on formulas in Zonato et al. (2020).
- 91     ▪ `HI_URB2D` is obtained by fitting a bounded normal distribution to the minimum (`MH_URB2D_MIN`), mean (`MH_URB2D`), and maximum (`MH_URB2D_MAX`) building height, as provided in `LCZ_UCP_default.csv`. The building height standard deviation is also required, and is approximated as  $(MH\_URB2D\_MAX - MH\_URB2D\_MIN) / 4$ .
- 92     ▪ For computational efficiency, `HI_URB2D` values lower than 5% were set to 0 after resampling, the remaining `HI_URB2D` percentages are re-scaled to 100%.

102    **Pathway 2:** In line with the former Fortran-based `W2W` procedure, **radiative and thermal parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell (see *Step 3*). These parameter values are not stored in the netcdf output, but are read from `URBPARAM_LCZ.TBL` and assigned automatically to the modal LCZ class when running the model.

107    ▪ *Step 5: Adjust global attributes*

108    In a final step, some global attributes are adjusted in the resulting netcdf files:

- 109    ▪ `NBUI_MAX` is added as a global attribute, reflecting the maximum amount of `HI_URB2D` classes that are not 0 across the model domain. This parameter can be used when compiling WRF, to optimize memory storage.
- 110    ▪ `NUM_LAND_CAT` is set to 41, to reflect the addition of 10 (or 11) built LCZ classes. This is not only done for the highest resolution domain file (e.g. `d04`), but also for **all of its parent domain files (e.g. `d01`, `d02`, `d03`)**. As such, make sure these files are also available in the input data directory.

116    Resulting output: `geo_em.d0X_LCZ_params.nc`

## Integration in WRF's preprocessing

**@ ANDREA:** Can you write a small text to describe 1) where this tool sits within WRF's typical workflow and 2) what needs to be done afterwards? E.g.:

- what a user needs to do after running the tool (eg. renaming netcdf outputs, what WRF tools to launch etc ...)
- indicate what namelist settings are required?

Perhaps it would make sense to have a schematic, similar to the first half of this Figure [Figure 1](#), available from [MeyerandRiechert\(2019\)](#)?

*D. Meyer, M. Riechert*

*Environmental Modelling and Software 112 (2019) 166–178*




Fig. 1. Typical WRF workflow and artefacts.

Figure 1: WRF workflow example.

## Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019), Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2021)).

## Important notes

- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the values according to the characteristics of your ROI.
- It is advised to use this tool with urban parameterization options BEP or BEP+BEM (`sf_urban_physics = 2` or `3`). In case you use this tool with the SLUCM model (`sf_urban_physics = 1`), make sure your lowest model level is above the highest building height. If not, `real.exe` will provide the following error message: `ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine urban - change ZDC and ZOC` **@ ANDREA: Correct??**

## Acknowledgements

We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez Sanchez and Oscar Brousse.

## References

- Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of heat island and thermal comfort over an urban airshed in India under varying land use/land cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient land use producing data tool for mesoscale models? Integration of urban LCZ in WRF over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O., Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley, M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018). WUDAPT: An urban weather, climate, and environmental modeling infrastructure for the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924. <https://doi.org/10.1175/BAMS-D-16-0236.1>
- Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert and crowd-sourced training data to map urban form and functions for the continental US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing detailed urban canopy parameters for mesoscale climate modelling: a comparison between WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*, 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler, W. (2021). Resolving the influence of local flows on urban heat amplification during heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>
- Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Urban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>
- Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X., Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>
- Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to contemporary land-use datasets across the contiguous united states using dynamical downscaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>

- 192 Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WU-*  
 193 *DAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. [https://www.wudapt.org/wp-content/](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)  
 194 [uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)  
 195 [pdf%20\(Accessed%20on%2011%20August%202021](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)
- 196 Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parame-  
 197 terisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>  
 198 <https://doi.org/10.1023/A:1016099921195>
- 199 Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into  
 200 WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged,  
 201 Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>  
 202 <https://doi.org/10.1007/s00704-019-02881-1>
- 203 Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M., &  
 204 Niyogi, D. (2021). Largescale Urban Impacts on a European Heat Wave Event. *Journal*  
 205 *of Geophysical Research : Atmospheres (Under Review)*, 1–33.
- 206 Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy  
 207 rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban*  
 208 *Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>  
 209 <https://doi.org/10.1016/j.uclim.2020.100616>
- 209 Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved  
 210 WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*,  
 211 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>  
 212 <https://doi.org/10.1016/j.atmosres.2020.105220>
- 212 Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model  
 213 coupled with an urban canopy parameterization for urban climate simulations—part I.  
 214 formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied*  
 215 *Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>  
 216 <https://doi.org/10.1007/s00704-009-0142-9>
- 216 Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W.,  
 217 Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of*  
 218 *the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric  
 219 Research. <https://doi.org/10.5065/1dfh-6p97>  
 220 <https://doi.org/10.5065/1dfh-6p97>
- 220 Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies.  
 221 *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>  
 222 <https://doi.org/10.1175/BAMS-D-11-00019.1>
- 223 Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the 'local climate  
 224 zone' scheme using temperature observations and model simulations [Journal Article].  
 225 *International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>  
 226 <https://doi.org/10.1002/joc.3746>
- 227 Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters  
 228 on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>  
 229 <https://doi.org/10.3390/atmos11121349>
- 230 Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang,  
 231 R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on  
 232 WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018),  
 233 100460. <https://doi.org/10.1016/j.uclim.2019.100460>  
 234 <https://doi.org/10.1016/j.uclim.2019.100460>
- 234 Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones,*  
 235 *Mitigation Strategies, building materials permeability and new buildings drag coefficient.*  
 236 <http://www.wudapt.org/wudapt-to-wrf/>. [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)  
 237 [product-tool/urban-canopy-model/WRF\\_urban\\_update\\_Readme\\_file\\_WRF4.3.pdf](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)  
 238 [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
- 238 Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the  
 239 performance of a novel WUDAPT averaging technique to define urban morphology with



240 mesoscale models. *Urban Climate*, 31(May 2019), 100584. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.uclim.2020.100584)  
241 [uclim.2020.100584](https://doi.org/10.1016/j.uclim.2020.100584)  
242 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., & Barlage, M. (Under Review).  
243 Exploring the effects of rooftop mitigation strategies on urban temperatures and energy  
244 consumption. *Journal of Advances in Modeling Earth Systems*, 1–29.

DRAFT