




W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere^{*1}, Daniel Argüeso², and Andrea Zonato³

¹ Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany ² Physics Department, University of the Balearic Islands, Palma, Spain ³ Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

Software

- [Review](#) 
- [Repository](#) 
- [Archive](#) 

Editor: [Editor Name](#) 

Submitted: 01 January XXXX

Published: 01 January XXXX

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

An important objective of WUDAPT, the World Urban Database and Access Portals Tools community project, is 1) to acquire and make accessible coherent and consistent information on form and function of urban morphology relevant to climate weather, and environment studies, and 2) to provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

The Python-based WUDAPT-to-WRF (W2W) package is developed in this context, and translates Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model (Skamarock et al., 2021). It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides an improved, more simple, and more efficient procedure to use LCZ information in WRF. Some important changes include a direct manipulation of the geogrid files (without the creation of temporary files), and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue, because of two recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the LCZ Generator web application (Demuzere et al., 2021), and 2) WRF versions > 4.3 (Skamarock et al., 2021) are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato et al. (Under Review) and Zonato & Chen (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here, so as to make the translation of LCZ-based parameters better and more simple.

^{*}corresponding author

Initial data requirements

In order to use the tool, two input files are required:

1. A **geo_em.d0X** (.nc) file for the inner WRF model domain in which one would like to use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool as part of the WRF Preprocessing System (WPS), without additional modifications of the standard procedure.
2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of the `geo_em.d0X.nc` file. There are a number of ways to obtain an LCZ map for your region of interest (ROI):
 - Extract your ROI from the continental-scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et al., 2020) (see [here](#) for more info).
 - Check if your ROI is already covered by the many LCZ maps available in the [submission table](#) of the LCZ Generator.
 - Use the [LCZ Generator](#) to make an LCZ map for your ROI. See also [here](#) for more information.

Workflow

The goal of the Python-based W2W tool is to obtain a WRF domain file (`geo_em.d0X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters relevant for all urban parameterizations embedded in WRF: the single layer urban canopy model (Noah/SLUCM, Kusaka et al. (2001)), the Building Environment Parameterization (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al. (2010)).

To get to that point, a number of sequential steps are required:

- *Step 1: Remove the default urban land cover*

The default urban land cover from MODIS is replaced with the dominant surrounding vegetation category, as done in Li et al. (2020). This procedure affects WRF's parameters `LU_INDEX`, `LANDUSEF` and `GREENFRAC`. `LU_INDEX` is selected as the dominant category from the `nlus` (default = 45) nearest grid points (excluding ocean, urban and lakes). `GREENFRAC` is calculated as the mean over all grid points with that category among the `nlus` nearest points. For each grid point, if `LANDUSEF` had any percentage of urban, it is set to zero and the percentage is added to the dominant category assigned to that grid point.

Resulting output: **geo_em.d0X_NoUrban.nc**

- *Step 2: Define the LCZ-based urban extent*

LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF resolution. Areas with `FRC_URB2D` < 0.2 (*frc*) are currently considered non-urban. This choice has been made to avoid to employ the urban schemes in areas where the majority of the landuse is vegetated, since the impact of the impervious surfaces is low. The `FRC_URB2D` field is also used to mask all other urban parameter fields, so that their extent is consistent.

Resulting output: **geo_em.d0X_LCZ_extent.nc**

74 ▪ *Step 3: Introduce modal built LCZ classes*

75 For each WRF grid cell, the mode of the underlying built LCZ classes is added to LU_INDEX
 76 (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default considers
 77 LCZ classes 1-10 as built classes (*bc*). In some cases, also LCZ E (or 15 - Bare rock or paved)
 78 can be considered as a built LCZ classes, as it might reflect large asphalt surfaces such as
 79 big parking lots or airstrips. In that case, the user must make sure the *bc* argument is set
 80 appropriately.

81 ▪ *Step 4: Assign urban canopy parameters*

82 Two pathways **THIS COULD BE UNDERSTOOD AS EXCLUDING PATHWAYS, BUT**
 83 **BOTH ARE ACTUALLY HAPPENING. PERHAPS CLARIFY** are followed when assign-
 84 ing the various urban canopy parameters to the LCZ map, and translating this information
 85 onto WRF's grid:

86 **Pathway 1: Morphological parameters** are assigned directly to the high-resolution LCZ
 87 map, and are afterwards aggregated to the lower-resolution WRF grid. In this way, the
 88 method produces a unique urban morphology parameter value for each WRF grid cell. This
 89 was found to be more efficient in reproducing urban boundary layer features, especially in
 90 the outskirts of the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#)
 91 routine (Varentsov et al., 2020).

92 Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv`, and
 93 are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014).
 94 In addition:

- 95 ▪ Building width (BW), available in `LCZ_UCP_default.csv`, is taken from `URBPARM_LC`
 96 `Z.TBL` (stored in WRF's run/ folder).
- 97 ▪ While `URBPARM_LCZ.TBL` also has values on street width, W2W derives street width from
 98 the mean building height (`MH_URB2D`) and the Height-to-Width ratio (`H2W`), to have
 99 these fields consistent.
- 100 ▪ Plan (`LP_URB2D`), frontal (`LF_URB2D`) and total (`LB_URB2D`) area indices are based
 101 on formulas in Zonato et al. (2020).
- 102 ▪ `HI_URB2D` is obtained by fitting a bounded normal distribution to the minimum
 103 (`MH_URB2D_MIN`), mean (`MH_URB2D`), and maximum (`MH_URB2D_MAX`)
 104 building height, as provided in `LCZ_UCP_default.csv`. The building height
 105 standard deviation is also required, and is approximated as $(MH_URB2D_MAX -$
 106 $MH_URB2D_MIN) / 4$.
- 107 ▪ For computational efficiency, `HI_URB2D` values lower than 5% were set to 0 after
 108 resampling, the remaining `HI_URB2D` percentages are re-scaled to 100%.

109 **Pathway 2:** In line with the former Fortran-based W2W procedure, **radiative and thermal**
 110 **parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell
 111 (see *Step 3*). These parameter values are not stored in the netcdf output, but are read from
 112 `URBPARM_LCZ.TBL` and assigned automatically to the modal LCZ class when running the
 113 model.

114 ▪ *Step 5: Adjust global attributes*

115 In a final step, some global attributes are adjusted in the resulting netcdf files:

- 116 ▪ `NBUI_MAX` is added as a global attribute, reflecting the maximum amount of
 117 `HI_URB2D` classes that are not 0 across the model domain. This parameter can be
 118 used when compiling WRF, to optimize memory storage.

- NUM_LAND_CAT is set to 41, to reflect the addition of 10 (or 11) built LCZ classes. This is not only done for the highest resolution domain file (e.g. d04), but also for **all of its parent domain files (e.g. d01, d02, d03)**. As such, make sure these files are also available in the input data directory.

Resulting output: `geo_em.d0X_LCZ_params.nc`

Integration in WRF's preprocessing

The current tool is designed to work with the `geo_em.d0X` files produced by `geogrid.exe`, which is available in the WRF Preprocessing System (WPS). The user should run `geogrid.exe` using its default settings, which will provide the various `geo_em.d0X.nc` files containing the static data fields. No additional variable are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL` table. The `W2W` tool (Figure 1) reads the standard `geo_em.d0X.nc` files (for all the domains) and produces the aforementioned `geo_em.d0X_LCZ_params.nc` files. The user should then simply rename these files to the standard name for each of the domains.




Figure 1: Modified workflow to set-up and run a WRF simulations including urban parameters derived from LCZs using W2W.

Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_NoUrban.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019),

Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2021)).

Important notes

- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the urban canopy parameters values according to the characteristics of your ROI.
- It is advised to use this tool with urban parameterization options BEP or BEP+BEM (`sf_urban_physics` = 2 or 3, respectively). In case you use this tool with the SLUCM model (`sf_urban_physics` = 1), make sure your lowest model level is above the highest building height. If not, `real.exe` will provide the following error message: `ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine urban - change ZDC and ZOC`

Acknowledgements

We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez Sanchez and Oscar Brousse.

References

- Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of heat island and thermal comfort over an urban airshed in India under varying land use/land cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient land use producing data tool for mesoscale models? Integration of urban LCZ in WRF over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O., Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley, M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018). WUDAPT: An urban weather, climate, and environmental modeling infrastructure for the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924. <https://doi.org/10.1175/BAMS-D-16-0236.1>
- Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert and crowd-sourced training data to map urban form and functions for the continental US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing detailed urban canopy parameters for mesoscale climate modelling: a comparison between WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*, 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>

- Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler, W. (2021). Resolving the influence of local flows on urban heat amplification during heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>
- Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Urban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>
- Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X., Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>
- Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to contemporary land-use datasets across the contiguous united states using dynamical downscaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>
- Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WUDAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. [https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20\(Accessed%20on%2011%20August%202021](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)
- Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parameterisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>
- Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged, Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>
- Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M., & Niyogi, D. (2021). Largescale Urban Impacts on a European Heat Wave Event. *Journal of Geophysical Research : Atmospheres (Under Review)*, 1–33.
- Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*, 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model coupled with an urban canopy parameterization for urban climate simulations—part I. formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>
- Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric Research. <https://doi.org/10.5065/1dfh-6p97>
- Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies. *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>
- Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the 'local climate zone' scheme using temperature observations and model simulations [Journal Article].

- 233 *International Journal of Climatology*, 34(4), 1062–1080. [https://doi.org/10.1002/joc.](https://doi.org/10.1002/joc.3746)
234 3746
- 235 Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters
236 on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>
237
- 238 Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang,
239 R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on
240 WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018),
241 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- 242 Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones,*
243 *Mitigation Strategies, building materials permeability and new buildings drag coefficient.*
244 <http://www.wudapt.org/wudapt-to-wrf/>. [https://ral.ucar.edu/sites/default/files/public/](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
245 [product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf](https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf)
- 246 Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the
247 performance of a novel WUDAPT averaging technique to define urban morphology with
248 mesoscale models. *Urban Climate*, 31(May 2019), 100584. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.uclim.2020.100584)
249 [uclim.2020.100584](https://doi.org/10.1016/j.uclim.2020.100584)
- 250 Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., & Barlage, M. (Under Review).
251 Exploring the effects of rooftop mitigation strategies on urban temperatures and energy
252 consumption. *Journal of Advances in Modeling Earth Systems*, 1–29.