

Genetic Architecture of Transcriptome Regulation

Heather E. Wheeler¹, Nicholas Knoblauch², GTEx Consortium, Nancy J. Cox³, Dan L. Nicolae¹, Hae Kyung Im¹

2015-05-13 12:26:40 ¹Department of Medicine, University of Chicago, ²Committee on Genetics, Genomics, and Systems Biology, University of Chicago, ³Division of Genetic Medicine, Vanderbilt University

1 Abstract

Lorem ipsum dolor sit amet, est ad doctus eligendi scriptorem. Mel erat falli ut. Feugiat legendos adipisci vix at, usu at laoreet argumentum suscipiantur. An eos adhuc aliquip scriptorem, te adhuc dolor liberavisse sea. Ponderum vivendum te nec, id agam brute disputando mei.

2 Introduction

Lorem ipsum dolor sit amet, est ad doctus eligendi scriptorem. Mel erat falli ut. Feugiat legendos adipisci vix at, usu at laoreet argumentum suscipiantur. An eos adhuc aliquip scriptorem, te adhuc dolor liberavisse sea. Ponderum vivendum te nec, id agam brute disputando mei.

Putant numquam tacimates at eum. Aliquip torquatos ex vis, mei et quando debitis appareat, impetus accumsan corrumpit in usu. Nam mucius facilis singulis id, duo ei autem imperdiet instructor. Cu ceteros alienum mel, id vix putant impedit, ex idque eruditi forensibus eum. Posse dicunt id usu. Ei iracundia constituto sed, duo ne exerci ignota, an eum unum conceptam.

Has audire salutandi no, ut eam dicat libris dicunt. Pri hendrerit quaerendum adversarium ea, dicat atqui munere et sea. Illum insolens eos ne, eu enim graece rationibus mea. At postea utamur mel, eius nonumes percipitur at vis. Numquam similique in per, te quo saepe utroque pericula.

Ea nonumy volumus usu, no mel inermis dissentias. Dico partiendo vituperatoribus eum et. Mea accusam convenire te, usu populo qualisque gloriatur ut. Eu eum oratio altera option, ad mea ignota scriptorem. Ne suas latine vix, eos oblique sanctus pertinax cu.

3 Methods

Lorem ipsum dolor sit amet, est ad doctus eligendi scriptorem. Mel erat falli ut. Feugiat legendos adipisci vix at, usu at laoreet argumentum suscipiantur. An eos adhuc aliquip scriptorem, te adhuc dolor liberavisse sea. Ponderum vivendum te nec, id agam brute disputando mei.

Putant numquam tacimates at eum. Aliquip torquatos ex vis, mei et quando debitis appareat, impetus accumsan corrumpit in usu. Nam mucius facilis singulis id, duo ei autem imperdiet instructor. Cu ceteros alienum mel, id vix putant impedit, ex idque eruditi forensibus eum. Posse dicunt id usu. Ei iracundia constituto sed, duo ne exerci ignota, an eum unum conceptam.

3.1 Equations

The deterministic part of the model is defined by this **in-line equation** as $\mu_i = \beta_0 + \beta_1 x$, and the stochastic part by the **centered equation**:

$$\frac{1}{\sqrt{2\pi}\sigma}e^{-(x-\mu_i)^2/(2\sigma^2)}$$

3.2 Tables

Warning: package 'knitr' was built under R version 3.1.2

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-0.09	0.1	-0.89	0.38
x	2.19	0.1	22.51	0.00

Table 1: This is a GLM summary table.

3.3 Plots

3.4 Citations

The relationship was first described by Reference 1. However, there are also opinions that the relationship is spurious [2]. We used R for our calculations [3], and we used package `knitcitations` [4] to make the bibliography.

4 Results and discussion

Lorem ipsum dolor sit amet, est ad doctus eligendi scriptorem. Mel erat falli ut. Feugiat legendos adipisci vix at, usu at laoreet argumentum suscipiantur. An eos adhuc aliquip scriptorem, te adhuc dolor liberavisse sea. Ponderum vivendum te nec, id agam brute disputando mei.

Putant numquam tacimates at eum. Aliquip torquatos ex vis, mei et quando debitis appareat, impetus accumsan corrumpit in usu. Nam mucius facilis singulis id, duo ei autem imperdiet instructor. Cu ceteros alienum mel, id vix putant impedit, ex idque eruditi forensibus eum. Posse dicunt id usu. Ei iracundia constituto sed, duo ne exerci ignota, an eum unum conceptam.

Has audire salutandi no, ut eam dicat libris dicunt. Pri hendrerit quaerendum adversarium ea, dicat atqui munere et sea. Illum insolens eos ne, eu enim graece rationibus mea. At postea utamur mel, eius nonumes percipitur at vis. Numquam similique in per, te quo saepe utroque pericula.

Ea nonumy volumus usu, no mel inermis dissentias. Dico partiendo vituperatoribus eum et. Mea accusam convenire te, usu populo qualisque gloriatur ut. Eu eum oratio altera option, ad mea ignota scriptorem. Ne suas latine vix, eos oblique sanctus pertinax cu.

References

1. Halpern BS, Regan HM, Possingham HP, McCarthy MA. Accounting for uncertainty in marine reserve design. *Ecol Letters*. Wiley-Blackwell; 2006;9: 2–11. doi:[10.1111/j.1461-0248.2005.00827.x](https://doi.org/10.1111/j.1461-0248.2005.00827.x)
2. Keil P, Belmaker J, Wilson AM, Unitt P, Jetz W. Downscaling of species distribution models: a hierarchical approach. *Freckleton R*, editor. *Methods Ecol Evol*. Wiley-Blackwell; 2012;4: 82–94. doi:[10.1111/j.2041-210x.2012.00264.x](https://doi.org/10.1111/j.2041-210x.2012.00264.x)



Figure 1: DGN-WB joint heritability (h^2). Local h^2 is estimated with SNPs within 1 Mb of each gene. Global h^2 is estimated with SNPs that are eQTLs in the Framingham Heart Study on other chromosomes (FDR < 0.05).




Figure 2: DGN-WB joint heritability (h^2) with known trans-eQTLs. Local h^2 is estimated with SNPs within 1 Mb of each gene. Known trans h^2 is estimated with SNPs that are trans-eQTLs in the Framingham Heart Study for each gene (FDR < 0.05).




Figure 3: Cross-tissue heritability (h^2) compared to tissue-wide h^2 . Cross-tissue local h^2 is estimated using the cross-tissue component (random effects) of the mixed effects model for gene expression and SNPs within 1 Mb of each gene. Tissue-wide local h^2 is estimated using the measured gene expression for each respective tissue and SNPs within 1 Mb of each gene.




Figure 4: Cross-tissue and tissue-wide comparison of heritability (h^2 , **A**) and standard error (SE, **B**). Cross-tissue local h^2 is estimated using the cross-tissue component (random effects) of the mixed effects model for gene expression and SNPs within 1 Mb of each gene. Tissue-wide local h^2 is estimated using the measured gene expression for each respective tissue and SNPs within 1 Mb of each gene.




Figure 5: Cross-tissue heritability (h^2) compared to tissue-specific h^2 . Cross-tissue local h^2 is estimated using the cross-tissue component (random effects) of the mixed effects model for gene expression and SNPs within 1 Mb of each gene. Tissue-specific local h^2 is estimated using the tissue-specific component (residuals) of the mixed effects model for gene expression for each respective tissue and SNPs within 1 Mb of each gene.




Figure 6: Cross-tissue and tissue-specific comparison of heritability (h^2 , **A**) and standard error (SE, **B**) estimation. Cross-tissue local h^2 is estimated using the cross-tissue component (random effects) of the mixed effects model for gene expression and SNPs within 1 Mb of each gene. Tissue-specific local h^2 is estimated using the tissue-specific component (residuals) of the mixed effects model for gene expression for each respective tissue and SNPs within 1 Mb of each gene.

3. R Core Team. R: A language and environment for statistical computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2014. Available: <http://www.R-project.org/>
4. Boettiger C. knitcitations: Citations for knitr markdown files [Internet]. 2014. Available: <http://CRAN.R-project.org/package=knitcitations>