

TOPICS IN TOPOLOGY: STABLE ∞ -CATEGORIES

1. 8/23/21

For the first few lectures we will try to motivate the theory of stable ∞ -categories from the point of view of algebra, in particular the theory of derived functors and categories. These in turn are motivated by the following general strategy in mathematics: to study a complicated object, try to build it out of simple parts and then reduce questions about the complicated object to questions about the simple parts.

Consider the following example in the setting of abelian groups. Given $n \in \mathbb{N}$, we have the cyclic group

$$\mathbb{Z}/n\mathbb{Z} := \text{cok}(\mathbb{Z} \xrightarrow{n} \mathbb{Z}),$$

i.e. the cokernel of the map from \mathbb{Z} to itself given by multiplication by n . We can think of this description as explaining how to build the “complicated” abelian group $\mathbb{Z}/n\mathbb{Z}$ out of two copies of the “simple” abelian group \mathbb{Z} . (Of course this is a toy example and $\mathbb{Z}/n\mathbb{Z}$ isn’t that complicated, but it’s more complicated than \mathbb{Z} in so far as you learn ordinary arithmetic before modular arithmetic).

Suppose we want to understand the tensor product $\mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z}/n\mathbb{Z}$ for some $m \in \mathbb{N}$. Naively, we could try to use the above description of $\mathbb{Z}/n\mathbb{Z}$ together with the trivial identity $\mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z} \cong \mathbb{Z}/m\mathbb{Z}$. That is, we could naively expect that

$$\begin{aligned} \mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z}/n\mathbb{Z} &\cong \mathbb{Z}/m\mathbb{Z} \otimes \text{cok}(\mathbb{Z} \xrightarrow{n} \mathbb{Z}) \\ &\cong \text{cok}(\mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z} \xrightarrow{n \otimes 1} \mathbb{Z}/m\mathbb{Z} \otimes \mathbb{Z}) \\ &\cong \text{cok}(\mathbb{Z}/m\mathbb{Z} \xrightarrow{n} \mathbb{Z}/m\mathbb{Z}), \end{aligned}$$

and then identify this last line as $\mathbb{Z}/\gcd(m, n)$ using Bezout’s identity. And in fact this is actually correct! The key fact that makes this work is that the operation $A \mapsto \mathbb{Z}/m\mathbb{Z} \otimes A$ preserves cokernels of abelian groups.

Now suppose we try to compute $\text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$, the abelian group of homomorphisms from $\mathbb{Z}/m\mathbb{Z}$ to $\mathbb{Z}/n\mathbb{Z}$, the same way. Our naive calculation would be

$$\begin{aligned} \text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}/n\mathbb{Z}) &\cong \text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, \text{cok}(\mathbb{Z} \xrightarrow{n} \mathbb{Z})) \\ &\cong \text{cok}(\text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z}) \xrightarrow{n} \text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, \mathbb{Z})) \\ &\cong \text{cok}(0 \xrightarrow{n} 0), \end{aligned}$$

which is of course just zero. But this is obviously wrong: say, when $m = n$ the identity morphism is a nonzero element of $\text{Hom}_{\text{Ab}}(\mathbb{Z}/n\mathbb{Z}, \mathbb{Z}/n\mathbb{Z})$. The key fact that makes this fail is that the operation $A \mapsto \text{Hom}_{\text{Ab}}(\mathbb{Z}/m\mathbb{Z}, A)$ does not preserve cokernels of abelian groups.

This distinction reflects a sense in which the operation $A \mapsto \mathrm{Hom}_{\mathrm{Ab}}(\mathbb{Z}/m\mathbb{Z}, A)$ is harder to study than $A \mapsto \mathbb{Z}/m\mathbb{Z} \otimes A$: there is at least one kind of argument available to us in studying the latter which is not available when studying the former.

Note that in the example the role of cokernels is a kind of “blueprint” for how to build a complicated thing out of two (potentially) simpler things. This is a special case of a colimit:

$$\mathrm{cok}(\mathbb{Z} \xrightarrow{n} \mathbb{Z}) \cong \mathrm{colim} \left(\begin{array}{ccc} \mathbb{Z} & \xrightarrow{n} & \mathbb{Z} \\ \downarrow & & \\ 0 & & \end{array} \right).$$

A colimit is a more general kind of blueprint for building a complicated thing out of a collection of (potentially) simpler things organized into a diagram (such as the one to the right above). We also have the related notion of limit, which again reads a diagram as a blueprint for building a new object but does so in a dual way. These satisfy dual universal properties, but for now these universal properties aren’t what we care about per se — just the fact that both notions give a way of building complicated objects out of simpler ones.

Returning to the example, the moral is that our ability to understand a given operation (e.g. a functor like $\mathbb{Z}/m\mathbb{Z} \otimes -$ or $\mathrm{Hom}_{\mathrm{Ab}}(\mathbb{Z}/m\mathbb{Z}, -)$) is constrained by the types of blueprints it preserves — the more it preserves, the easier it will be to study. With that in mind, the deep idea of derived categories and functors is the following: to study a functor which doesn’t preserve some class of blueprints, try to “upgrade” it so that it does, then study the “upgraded” functor. To discuss this more precisely, let’s introduce some terminology.

Definition 1.1. Let $F : \mathcal{C} \rightarrow \mathcal{C}'$ be a functor.

- (1) F is right exact if it preserves all finite colimits.
- (2) F is left exact if it preserves all finite limits.
- (3) F is exact if it is both right and left exact.

This terminology is in turn motivated by the following special case. Recall that a category is abelian if, roughly speaking, it behaves like the category of abelian groups.

Theorem 1.2. *A functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ between abelian categories is right exact (resp. left exact) if and only if it preserves finite direct sums and cokernels (resp. kernels).*

Now we can give a slightly more precise version of the “deep idea” above. Suppose $F : \mathcal{C} \rightarrow \mathcal{C}'$ is a functor between abelian categories which is left but not right exact (a dual version of the following discussion applies to right but not left exact functors). Then (under mild hypotheses) we can canonically “upgrade” F to an exact functor $R^\bullet F$. BUT we have to “upgrade” the source and target category as well, so that we have a functor

$$R^\bullet F : \mathcal{D}^-(\mathcal{C}) \rightarrow \mathcal{D}^-(\mathcal{C}').$$

Here $\mathcal{D}^-(\mathcal{C})$ and $\mathcal{D}^-(\mathcal{C}')$ are called the (bounded below) “derived categories” of \mathcal{C} and \mathcal{C}' , and $R^\bullet F$ is called the “total right derived functor” of F .

2. 8/27/21

2.1. **Review.**

2.1.1. *General problem.* Given a functor $F: \mathcal{C} \rightarrow \mathcal{C}'$ and an object $X \in \mathcal{C}$, how shall we compute $F(X)$?

2.1.2. *General strategy.* Find a way to write

$$X \cong \operatorname{colim}_{i \in I} X_i \text{ (or } X \simeq \lim_{i \in I} X_i \text{)}$$

so that each X_i is simpler than X and so that F preserves $\operatorname{colim}_{i \in I} X_i$ and $\lim_{i \in I} X_i$, i.e.

$$F(\operatorname{colim}_{i \in I} X_i) = \operatorname{colim}_{i \in I} F(X_i), \quad F(\lim_{i \in I} X_i) = \lim_{i \in I} F(X_i).$$

In general, functors which preserve many limits or colimits are easier to study.

2.2. **Extended general strategy.** If you can, upgrade F so that it preserves more limits or colimits, hence it becomes easier to study.

2.2.1. *Specific instance.* Given a left (but not right) exact functor, $F: \mathcal{C} \rightarrow \mathcal{C}'$ of Abelian categories we can upgrade F to an exact "total (right) derived functor" $R^\bullet F: D^-(\mathcal{C}) \rightarrow D^-(\mathcal{C}')$ between "(bounded below) derived categories".

2.2.2. *Question.* How is $D^-(\mathcal{C})$ related to \mathcal{C} and how is $R^\bullet F$ related to F ?

2.2.3. *Rough description of $D^-(\mathcal{C})$.*

- (1) For each $n \in \mathbb{Z}$ we have a fully faithful functor $i_n: \mathcal{C} \rightarrow D^-(\mathcal{C})$ and an essentially surjective functor ("cohomology") $H^n: D^-(\mathcal{C}) \rightarrow \mathcal{C}$ such that for all $m, n \in \mathbb{Z}$,

$$H^n \circ i_m \cong \begin{cases} \operatorname{id}_{\mathcal{C}} & m = n \\ 0 & m \neq n \end{cases}$$

- (2) There is an autoequivalence ("the shift functor") $[1]: D^-(\mathcal{C}) \rightarrow D^-(\mathcal{C})$ such that $H^n \circ ([1]) \simeq H^{n+1}$ and $[1] \circ i_n \simeq i_{n-1}$.

- (3) $D^-(\mathcal{C})$ has a zero object and $X \simeq 0$ if and only if $H^n(x) \simeq 0$ for all n .

2.2.4. *Analogy.* Passing from \mathcal{C} to $D^-(\mathcal{C})$ is like passing from \mathbb{R} to \mathbb{C} : the latter is more abstract but it's easier to study because it has better formal properties. The cohomologies $H^n(X) \subset \mathcal{C}$ are analogous to the real and imaginary parts of a complex number.

2.2.5. *Warnings.* Given $X \in D^-(\mathcal{C})$, for all $n \in \mathbb{Z}$ implies $X \simeq 0$. But given $X, Y \in D^-(\mathcal{C})$, $H^n(X) \simeq H^n(Y)$ for all $n \in \mathbb{Z}$ does not imply $X \simeq Y$.

Later we'll axiomatize the above data and their main properties as saying " $D^-(\mathcal{C})$ has a t -structure whose heart is \mathcal{C} ."

2.2.6. *Rough description of $R^\bullet F$.*(1) We can recover F from

$$\begin{array}{ccc} \mathcal{D}^-(\mathcal{C}) & \xrightarrow{R^\bullet F} & \mathcal{D}^-(\mathcal{C}') \\ i_0 \downarrow & & \downarrow H^0 \\ \mathcal{C} & \xrightarrow{F} & \mathcal{C}' \end{array}$$

(2) Given $n \in \mathbb{Z}$, we call the functor $R^n F$ defined by

$$\begin{array}{ccc} \mathcal{D}^-(\mathcal{C}) & \xrightarrow{R^\bullet F} & \mathcal{D}^-(\mathcal{C}') \\ i_0 \downarrow & & \downarrow H^n \\ \mathcal{C} & \xrightarrow{R^n F} & \mathcal{C}' \end{array}$$

the " n th (right) derived functors of F ".(3) We have $R^n F \simeq 0$ for $n < 0$ ($R^\bullet F$ is "left t -exact").(4) The right derived functors together take short exact sequences in \mathcal{C} to long exact sequences in \mathcal{C}' .

3. 8/30/21

Last time: Given a left exact functor $F: \mathcal{C} \rightarrow \mathcal{C}'$ between 2 abelian categories we described the "total right derived functor"

$$R^\bullet F: D^-(\mathcal{C}) \rightarrow D^-(\mathcal{C}').$$

- Ultimately $R^\bullet F$ is easier to study than F because it's exact. Aside: last time in our "rough description of $D^-(\mathcal{C})$ ", part (3) a morphism $f: X \rightarrow Y$ in $D^-(\mathcal{C})$ is an isomorphism if and only if $H^n(f): H^n(X) \rightarrow H^n(Y)$ is an isomorphism for all $n \in \mathbb{Z}$. Last thing we said: (4) The $\{R^n F\}_{n \in \mathbb{Z}}$ together take short exact sequences in \mathcal{C} to long exact sequences in \mathcal{C}' . (Historically, people discovered the $R^n F$'s before they discovered $R^\bullet F$ and $D^-(\mathcal{C})$).

Example (abelian groups)

- Fact (specific to Ab): given $A, B \in \text{Ab}$, $R^n \text{Hom}_{\text{Ab}}(A, B) \sim 0$ for $n > 1$ (and $n < 0$).

- convention: we write Ext^n for $R^n \text{Hom}$

Back to our example: (4) says we have a LES:

$$\begin{array}{ccccccc} & 0 & & 0 & & & \\ & | \simeq & & | \simeq & & & \\ 0 & \longrightarrow & \text{Hom}(\mathbb{Z}/m, \mathbb{Z}) & \xrightarrow{n} & \text{Hom}(\mathbb{Z}/m, \mathbb{Z}) & \longrightarrow & \text{Hom}(\mathbb{Z}/m, \mathbb{Z}/n) \\ & & & & & \searrow & \\ & & & & & \text{Ext}^1(\mathbb{Z}/m, \mathbb{Z}) & \longrightarrow & \text{Ext}^1(\mathbb{Z}/m, \mathbb{Z}) & \longrightarrow & \text{Ext}^1(\mathbb{Z}/m, \mathbb{Z}/n) & \longrightarrow & 0 \\ & & & & & | \simeq & & | \simeq & & & & \\ & & & & & \mathbb{Z}/m & \xrightarrow{n} & \mathbb{Z}/m & & & & \end{array}$$

$$\begin{aligned} \Rightarrow \operatorname{Hom}(\mathbb{Z}/m, \mathbb{Z}/n) &\simeq \mathbb{Z}/\gcd(m, n) \\ \mathbb{Z}/\gcd(m, n) &\simeq \mathbb{Z}/\frac{m}{d} \simeq \frac{m}{d}\mathbb{Z}/m. \end{aligned}$$

More generally, suppose $0 \rightarrow A \xrightarrow{f} B \rightarrow C \rightarrow 0$ is a SES in \mathcal{C} .

Then $\{R^n F(C)\}_{n \in \mathbb{Z}}$ is sometimes determined by the data of $\{R^n F(A) \xrightarrow{R^n F(f)} R^n F(B)\}_{n \in \mathbb{Z}}$.

BUT, $R^\bullet F(i_0 C)$, hence $\{R^n F(C)\}_{n \in \mathbb{Z}}$, is determined by $R^\bullet F(A) \xrightarrow{R^\bullet F(f)} R^\bullet F(B)$.

\Rightarrow we lose information by passing from $R^\bullet F$ to the $\{R^n F\}_{n \in \mathbb{Z}}$.

4. 9/01/21

Big picture so far: Derived categories "upgrade" abelian categories to have more exact functors between them, which makes them easier to study.

Given a left exact functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ between two abelian categories, we discussed

- The total right derived functor $R^\bullet F : D^-(\mathcal{C}) \rightarrow D^-(\mathcal{C}')$
- Component right derived functors $R^n F : \mathcal{C} \rightarrow \mathcal{C}'$

Theorem 4.1. *Suppose we have a short exact sequence $0 \rightarrow A \xrightarrow{\phi} B \rightarrow C \rightarrow 0$ in \mathcal{C} .*

- (1) $F(\phi) : F(A) \rightarrow F(B)$ is rarely enough to determine $F(C)$
- (2) $R^n F(\phi) : R^n F(A) \rightarrow R^n F(B)$ is sometimes enough to determine $R^n F(C)$.
- (3) $R^\bullet F(i_0 \phi) : R^\bullet F(i_0 A) \rightarrow R^\bullet F(i_0 B)$ is always enough to determine $R^\bullet F(i_0 C)$.

Recall from last time that we can compute $F(C)$ from $R^\bullet F(i_0 C)$, so $R^\bullet F(i_0 \phi) : R^\bullet F(i_0 A) \rightarrow R^\bullet F(i_0 B)$ also determines $F(C)$. A naive argument that (3) is true could go as follows:

$$\begin{aligned} R^\bullet F(i_0 C) &\cong R^\bullet F(i_0(\operatorname{cok}(A \xrightarrow{\phi} B))) \\ &\cong R^\bullet F(\operatorname{cok}(i_0 A \xrightarrow{i_0 \phi} i_0 B)) \\ &\cong \operatorname{cok}(R^\bullet F(i_0 A) \xrightarrow{R^\bullet F(i_0 \phi)} R^\bullet F(i_0 B)) \end{aligned}$$

The first isomorphism is immediate, and the third isomorphism holds because $R^\bullet F$ is an exact functor, so all we need to examine is the second isomorphism. In particular, we want to know if i_0 preserves cokernels. This "naive" argument actually works, but there is a subtlety that arises.

Important note

There are two versions of the derived category $D^-(\mathcal{C})$.

- (1) The derived 1-category $D_1^-(\mathcal{C})$
- (2) The derived ∞ -category $D_\infty^-(\mathcal{C})$

Rough idea of ∞ -categories

Definition 4.2. **Top** is the category whose objects are topological spaces and whose morphisms are continuous maps.

Definition 4.3. An ∞ -category \mathcal{C} is a category enriched in **Top**.

This means that an ∞ -category \mathcal{C} has

- A set of objects like a normal category
- For each pair of objects $X, Y \in \text{Ob}(\mathcal{C})$, a topological space of morphisms $\text{Hom}_{\mathcal{C}}(X, Y)$.
- For all $X, Y, Z \in \text{Ob}(\mathcal{C})$, a continuous composition map $\circ : \text{Hom}_{\mathcal{C}}(X, Y) \times \text{Hom}_{\mathcal{C}}(Y, Z) \rightarrow \text{Hom}_{\mathcal{C}}(X, Z)$.

Definition 4.4. A ∞ -functor $F : \mathcal{C} \rightarrow \mathcal{C}'$ is a functor enriched in **Top**. Namely, $F : \text{Hom}_{\mathcal{C}}(X, Y) \rightarrow \text{Hom}_{\mathcal{C}'}(F(X), F(Y))$ is a continuous map for all pairs of objects $X, Y \in \text{Ob}(\mathcal{C})$.

A normal 1-category is naturally an ∞ -category by giving the each hom-set the discrete topology. Conversely, if \mathcal{C} is an ∞ -category, we define a 1-category $h\mathcal{C}$ called the homotopy category. $h\mathcal{C}$ has the same objects as \mathcal{C} , and the hom-sets of $h\mathcal{C}$ are given by the connected components of the hom-sets of \mathcal{C} .

- $\text{Ob}(\mathcal{C}) = \text{Ob}(h\mathcal{C})$
- $\text{Hom}_{h\mathcal{C}}(X, Y) := \pi_0(\text{Hom}_{\mathcal{C}}(X, Y))$

Given a morphism in $\text{Hom}_{\mathcal{C}}(X, Y)$, we can get a morphism in $\text{Hom}_{h\mathcal{C}}(X, Y)$ by taking the connected component which it lies in.

There is an obvious ∞ -functor

$$h : \mathcal{C} \rightarrow h\mathcal{C}$$

. This functor serves as a bridge between the two versions of the derived category.

$$D_1^-(\mathcal{C}) \cong h(D_{\infty}^-(\mathcal{C}))$$

Everything we've mentioned so far in our rough description of $D^-(\mathcal{C})$ applies to both versions.

There are also two versions of the functors i_n and H^n , which are related by h as follows:

$$\begin{array}{ccccc} & & D_{\infty}^-(\mathcal{C}) & & \\ & i^n \nearrow & \downarrow h & \nwarrow H^n & \\ \mathcal{C} & & & & \mathcal{C} \\ & i^n \searrow & \downarrow h & \swarrow H^n & \\ & & D_1^-(\mathcal{C}) & & \end{array}$$

5. 9/3/2021

The above diagram implies that, given two objects $X, Y \in D_{\infty}^-(\mathcal{C})$, we have

$$\text{Hom}_{D_1^-(\mathcal{C})}(X, Y) \simeq \pi_0 \text{Hom}_{D_{\infty}^-(\mathcal{C})}(X, Y).$$

So $D_{\infty}^-(\mathcal{C})$ contains more information than $D_1^-(\mathcal{C})$.

Recall: Given $F : \mathcal{C} \rightarrow \mathcal{C}'$ a left exact functor between abelian categories and $0 \rightarrow A \rightarrow B \rightarrow C \rightarrow 0$ a short exact sequence in \mathcal{C} , then $R^{\bullet}F(i_0C)$ is determined by $R^{\bullet}F(i_0A) \xrightarrow{R^{\bullet}F(i_0\phi)} R^{\bullet}F(i_0B)$ (whereas $F(C)$ is not determined by $F(A) \xrightarrow{F(\phi)} F(B)$ in general). This is true as

stated for derived ∞ -categories: $i_0: \mathcal{C} \rightarrow D_{\infty}^{-}(\mathcal{C})$ preserves cokernels of monomorphisms (e.g. $A \xrightarrow{\phi} B$) and $R^{\bullet}F: D_{\infty}^{-}(\mathcal{C}) \rightarrow D_{\infty}^{-}(\mathcal{C}')$ preserves cokernels.

$$\left(\begin{array}{ccc} \text{To be explicit, we can define } R^{\bullet}F \text{ either for } D_{\infty} \text{ or } D_1, \text{ and these are related by } h: & & \\ D_{\infty}^{-}(\mathcal{C}) & \xrightarrow{R^{\bullet}F} & D_{\infty}^{-}(\mathcal{C}') \\ \downarrow h & & \downarrow h \\ D_1^{-}(\mathcal{C}) & \xrightarrow{R^{\bullet}F} & D_1^{-}(\mathcal{C}') \end{array} \right)$$

But cokernels (and colimits/limits in general) don't behave well in $D_1^{-}(\mathcal{C})$, and $i_0: \mathcal{C} \rightarrow D_1^{-}(\mathcal{C})$ does not preserve cokernels of monomorphisms in general (in particular of $A \xrightarrow{\phi} B$).

Workaround: Equip $D_1^{-}(\mathcal{C})$ with an extra structure to remember what morphisms were cokernels in $D_{\infty}^{-}(\mathcal{C})$. This is the structure of a **triangulated category**.

Rough Idea: A triangulated category is a (1-)category \mathcal{C} equipped with a distinguished autoequivalence $[1]: \mathcal{C} \rightarrow \mathcal{C}$ and a class of distinguished compositions $X \xrightarrow{f} Y \rightarrow Z$ called **exact triangles**.

These would satisfy some axioms, in particular that Z is determined up to isomorphism by f , and that any morphism $f: X \rightarrow Y$ can be completed to an exact triangle.

We call Z a "mapping cone" of f and sometimes write it $\text{cone}(f)$, but it does not satisfy any universal property.

We can define a triangulated structure on $D_1^{-}(\mathcal{C})$ by declaring mapping cones to be the images of cokernels in $D_{\infty}^{-}(\mathcal{C})$.

This gives a way of fixing our naive argument:

$$\begin{aligned} R^{\bullet}F(i_0 C) &\simeq R^{\bullet}F(i_0 \text{cok}(A \xrightarrow{\phi} B)) && \text{cokernel in } \mathcal{C} \\ &\simeq R^{\bullet}F(\text{cone}(i_0 A \xrightarrow{i_0 \phi} i_0 B)) && \text{cokernel in } D_1^{-}(\mathcal{C}) \\ &\simeq \text{cone}(R^{\bullet}F(i_0 A) \xrightarrow{R^{\bullet}F(i_0 \phi)} R^{\bullet}F(i_0 B)). && \text{cokernel in } D_1^{-}(\mathcal{C}') \end{aligned}$$

This works: $i_0: \mathcal{C} \rightarrow D_1^{-}(\mathcal{C})$ take cokernels of monomorphisms to mapping cones and $R^{\bullet}F: D_1^{-}(\mathcal{C}) \rightarrow D_1^{-}(\mathcal{C}')$ preserves mapping cones (whenever we say a functor between triangulated categories is exact we also mean that this preserves mapping cones).

6. 9/8/2021

In the previous lectures, we've seen there is a workaround of the poor behaviors of certain functors by working in the derived category. In particular, we claimed that $R^{\bullet}F(i_0 \text{cok}(\phi))$ is determined by $R^{\bullet}F(i_0 \phi)$ while $R^{\bullet}(\text{cok}(\phi))$ is not determined by $R^{\bullet}(\phi)$. However, even though we know derived functors $R^{\bullet}F$ behaves nicely, the derived category itself does not have "well-behaved" cokernels. In particular, i_0 does not preserve the cokernels of monomorphisms. So, we work with the derived ∞ -category: $D_{\infty}^{-}(\mathcal{C})$ instead of the derived 1-category: $D_1^{-}(\mathcal{C})$.

Note that we can also salvage the argument in the derived 1-category $D_1^-(\mathcal{C})$ if we consider as a triangulated category. The derived ∞ -category $D_\infty^-(\mathcal{C})$ is a basic example of a stable ∞ -category.

Definition 6.1. An ∞ -category is stable if:

- (1) It has a zero object.
- (2) Every morphism has a kernel (fiber) and cokernel (cofiber), and
- (3) the kernels (fibers) and cokernels (cofibers) behave especially well: every morphism is the kernel (fiber) of its cokernel (cofiber), and the cokernel (cofiber) of its kernel (fiber).

Remarks: (1) By convention, we will say "fiber" and "cofiber" instead of "kernels" and "cokernels" unless specifically working in an abelian category. But the definition will be the same:

Definition 6.2. A cofiber of a morphism $X \xrightarrow{f} Y$ in a stable ∞ -category $\text{cof}(X \xrightarrow{f} Y)$ is the colimit of the following diagram:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \\ 0 & & \end{array}$$

- (2) It turns out if an ∞ -category \mathcal{C} is stable, then its homotopy category $h\mathcal{C}$ is canonically triangulated: given $X \xrightarrow{f} Y$ in \mathcal{C} , set

$$\text{Cone}(hX \xrightarrow{hf} hY) := h(\text{cof}(X \xrightarrow{f} Y))$$

and

$$hX[1] := h(\text{cof}(X \xrightarrow{f} Y))$$

- (3) The definition of stable ∞ -categories is closer to the definition of abelian categories than the definition of triangulated categories. For example, the abelian category axioms include: "every monomorphism is a kernel of its cokernel, and every epimorphism is a cokernel of its kernel".

Related to (3), stable ∞ -categories are in some ways "easier" to work with than triangulated categories. Now we will discuss an example, the Riemann sphere \mathbb{CP}^1 .

We can obtain \mathbb{CP}^1 by gluing two copies of \mathbb{C} along \mathbb{C}^* , i.e. a colimit in the category of complex manifolds or the category of algebraic varieties. Specifically, \mathbb{CP}^1 is the colimit of the following diagram.

$$\begin{array}{ccc} \mathbb{C}^* & \longrightarrow & \mathbb{C} \\ \downarrow & & \\ \mathbb{C} & & \end{array}$$

. In algebraic geometry, a module over $\mathbb{C}[x]$ is a quasi-coherent sheaf over $\mathbb{C} \cong \text{Spec} \mathbb{C}[x]$. And a quasi-coherent sheaf on \mathbb{C} can be restricted to a quasi-coherent sheaf on \mathbb{C}^* . (Algebraically, this is a localization:

$$- \otimes_{\mathbb{C}[x]} \mathbb{C}[x^{\pm 1}] : \mathbb{C}[x]\text{-Mod} \rightarrow \mathbb{C}[x^{\pm 1}]\text{-Mod}$$

. A quasi-coherent sheaf on \mathbb{CP}^k is defined as a pair of quasi-coherent sheaves, each on one copy of \mathbb{C} , together with isomorphisms between their restrictions to $QCoh(\mathbb{C}^*)$. In particular $QCoh(\mathbb{CP}^1)$ can be defined as the limit (in the category of abelian categories) of the following diagram:

$$\begin{array}{ccc} QCoh(\mathbb{C}^*) & \longleftarrow & QCoh(\mathbb{C}) \\ \uparrow & & \\ QCoh(\mathbb{C}) & & \end{array}$$

. This remains true if we pass to derived infinity categories, but not derived one-categories! $D_{\infty}^{-} QCoh(\mathbb{CP}^1)$ is just the limit of the diagram:

$$\begin{array}{ccc} D_{\infty}^{-} QCoh(\mathbb{C}^*) & \longleftarrow & D_{\infty}^{-} QCoh(\mathbb{C}) \\ \uparrow & & \\ D_{\infty}^{-} QCoh(\mathbb{C}) & & \end{array}$$

.

7. 9/10/2021

Next up: ∞ -categories + simplicial set (ref: Lurie HTT Ch1)

- We said that topological categories capture the basic intuition of ∞ -categories. but really we will formalize the latter differently using simplicial sets.

- Later we'll discuss a sense in which the two notions are equivalent.

- For $n \in \mathbb{N}$, we write $[n] := \{0, 1, \dots, n\}$

Definition 7.1. (1) The simplex category Δ has objects $\{[n]\}_{n \in \mathbb{N}}$ and morphisms (non-strictly) order-preserving maps.

(2) If \mathcal{C} is a category, a simplicial object of \mathcal{C} is a functor $\Delta^{op} \rightarrow \mathcal{C}$. We write \mathcal{C}_{Δ} for the category of simplicial objects of \mathcal{C} .

(3) A cosimplicial object of \mathcal{C} is a functor $\Delta \rightarrow \mathcal{C}$

(4) A simplicial set is a simplicial object of **Set**


- Explicitly, a simplicial set S_{\bullet} is the data of

- a set S_n for each $n \in \mathbb{N}$
- a function $p^*: S_n \rightarrow S_m$ for each order-preserving function $p: [m] \rightarrow [n]$. Such that the p^* are compatible with composition.

- Some standard terminology/notation:


- p^* is the "pullback along p "
- for $j \in [n]$ the face map $d_j: S_n \rightarrow S_{n-1}$ is the pullback along

$$i \rightarrow \begin{cases} i & i \subset j \\ i+1 & i \supseteq j \end{cases}$$



- for $j \in [n]$ the degeneracy map $f_j: S_n \rightarrow S_{n+1}$ is the pullback along

$$i \rightarrow \begin{cases} i & i \subseteq j \\ i-1 & i \supset j \end{cases}$$



- Every order-preserving map is a composition of these, hence a simplicial set S_\bullet is determined by the sets S_n together with its face and degeneracy maps.

-Idea behind the terminology

- Given $n \in \mathbb{N}$, we have the standard (topological) n -simplex

$$|\Delta^n| = \{(x_0, \dots, x_n) \in [0, 1]^{n+1} \subset \mathbb{R}^{n+1} \mid \sum_{i=0}^n x_i = 1\}$$

- for each $j \in [n]$ the inclusion of the j th face $|\Delta^{n-1}| \hookrightarrow |\Delta^n|$ is given by

$$(x_0, \dots, x_{n-1}) \mapsto (x_0, \dots, x_{j-1}, 0, x_j, \dots, x_{n-1})$$

- we also have the "degeneration" map $|\Delta^{n+1}| \rightarrow |\Delta^n|$ given by

$$(x_0, \dots, x_{n+1}) \mapsto (x_0, \dots, x_j + x_{j+1}, \dots, x_{n+1})$$

- one can define a similar map $|\Delta^m| \xrightarrow{p^*} |\Delta^n|$ for any order-preserving map $[m] \xrightarrow{p} [n]$ compatibly with composition. Thus the $|\Delta^n|$ form a cosimplicial object of **Top**.

Key Example #1 :

Given a topological space X , we define the fundamental ∞ -groupoid of X (or the singular complex of X) is the simplicial set defined $\Pi(X)_\bullet$ defined by:

- $\Pi(X)_n = \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X)$
- given $p: [m] \rightarrow [n]$, we defined $p^*: \text{Hom}_{\mathbf{Top}}(|\Delta^n|, X) \rightarrow \text{Hom}_{\mathbf{Top}}(|\Delta^m|, X)$ by taking $f: |\Delta^n| \rightarrow X$ to $f \circ p_*: |\Delta^m| \rightarrow X$.

8. 9/13/21

Recall: Last time we introduced the simplex category by considering finite sets $[n]$ and order preserving maps.

Key example #0 The standard cosimplicial space $|\Delta^\bullet|$, i.e., $\Delta \xrightarrow{|\Delta^\bullet|} \mathbf{Top}$, $[n] \mapsto |\Delta^n|$

Key example #1. The fundamental ∞ -groupoid $\Pi(X)_\bullet$ of a space X .

Quick definition: $\Pi(X)$ is defined as the composition

$$\Delta^{op} \xrightarrow{|\Delta^\bullet|^{op}} \mathbf{Top}^{op} \xrightarrow{\mathbf{Hom}_{\mathbf{Top}}(-, X)} \mathbf{Set}$$

Remark. Given a base point $x_0 \in X$, $\Pi(X)$ contains all the data needed to define $\pi_1(X, x_0)$.

• The point x_0 defines a 0-simplex $|\Delta^0| \xrightarrow{x_0} X$, where x_0 denotes the constant map in $\Pi(X)_0 = \mathbf{Hom}_{\mathbf{Top}}(|\Delta^0|, X)$.

- A path $\gamma: x_0 \rightarrow x_0$ defines a 1-simplex

$$|\Delta^1| \xrightarrow{\gamma} X$$


such that $d_0(\gamma) = d_1(\gamma) = x_0$.

- A homotopy h from γ_1 to γ_2 defines a 2-simplex

$$|\Delta^2| \xrightarrow{h} X$$

such that $d_1(h) = \gamma_1$, $d_2(h) = \gamma_2$, $d_3(h) = Id_{x_0}$.

Here is the picture of $|\Delta^2|$:



More generally, any 2-simplex $|\Delta^2| \xrightarrow{h} X$ defines a homotopy between $d_1(h)$ and $d_0(h) \circ d_2(h)$. Therefore, $\Pi(X)$ encodes the composition law for homotopy classes of paths.

Preview: If X is sufficiently nice, e.g., a C.W. complex, then we can recover the homotopy type of X from $\Pi(X)$.

Summary $\Pi(X)_\bullet$ extends $\pi_1(X, x_0)$ by allowing x_0 to vary and by remembering actual homotopies between paths rather than just the relation of homotopy equivalence.

Key example #2 Given a category \mathcal{C} , we define the **nerve** $\mathcal{N}(\mathcal{C}) \in \mathbf{Set}_\Delta$ as follows:

Let $\mathcal{N}(\mathcal{C})_n$ be the set of composable sequences of morphisms in \mathcal{C} .

$$\mathcal{N}(\mathcal{C})_n := \{A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_{n-1} \xrightarrow{f_n} A_n\}$$

The face map $d_j : \mathcal{N}(\mathcal{C})_n \rightarrow \mathcal{N}(\mathcal{C})_{n-1}$ is defined by the composition

$$d_j(A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_{n-1} \xrightarrow{f_n} A_n) = A_0 \xrightarrow{f_1} \cdots \rightarrow A_{j-1} \xrightarrow{f_{j+1} \circ f_j} A_{j+1} \rightarrow \cdots \xrightarrow{f_n} A_n$$

9. 9/15/2021

Last time: more on the fundamental co-groupoid/singular complex, started discussing nerves.

Key example #2: Given a category \mathcal{C} , we define $\mathcal{N}(\mathcal{C}) \in \mathbf{Set}_\Delta$ as follows:

$\mathcal{N}(\mathcal{C})_n$ is the set of composable sequences of n morphisms:

$$\mathcal{N}(\mathcal{C})_n = \{A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_{n-1} \xrightarrow{f_n} A_n\}$$

-The face map $d_j : \mathcal{N}(\mathcal{C})_n \rightarrow \mathcal{N}(\mathcal{C})_{n-1}$ is defined by composition:

$$d_j(A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_{n-1} \xrightarrow{f_n} A_n) = A_0 \xrightarrow{f_1} A_1 \rightarrow \cdots \rightarrow A_j \xrightarrow{f_{j+1} \circ f_j} A_{j+1} \rightarrow \cdots \xrightarrow{f_n} A_n$$

-The degeneracy map $s_j : \mathcal{N}(\mathcal{C})_n \rightarrow \mathcal{N}(\mathcal{C})_{n+1}$ is defined using identity morphisms:

$$s_j(A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} A_{n-1} \xrightarrow{f_n} A_n) = A_0 \xrightarrow{f_1} \cdots \xrightarrow{f_j} A_j \xrightarrow{id_{A_j}} A_j \xrightarrow{f_{j+1}} \cdots \xrightarrow{f_n} A_n$$

-compatibility with composition in $\Delta \Leftrightarrow$ associativity and identity axioms for categories.

Quick definition: note that a poset (like $[n]$) is the same data as a category with at most one morphism between any two objects (i.e. $i \leq j \Leftrightarrow \text{Hom}(i, j)$ nonempty), and order-preserving maps are the same as functors.

\Rightarrow the assignment $n \mapsto [n] \in \mathbf{Cat}$ (This means category of category) defines a cosimplicial category $\Delta \xrightarrow{[\bullet]} \mathbf{Cat}$.

-Then $\mathcal{N}(\mathcal{C})$ is just the composition

$$\Delta^{op} \xrightarrow{[\bullet]^{op}} \mathbf{Cat}^{op} \xrightarrow{\mathbf{Hom}_{\mathbf{Cat}}(-, \mathcal{C})} \mathbf{Set}$$

-The data of \mathcal{C} is equivalent to the data of $\mathcal{N}(\mathcal{C})$:

$$Ob(\mathcal{C}) = \mathcal{N}(\mathcal{C})_0$$

$$Mor(\mathcal{C}) = \mathcal{N}(\mathcal{C})_1$$

$$composition \leftrightarrow d_1 : \mathcal{N}(\mathcal{C})_2 \rightarrow \mathcal{N}(\mathcal{C})_1$$

-The simplicial sets $\Pi(X)$, and $\mathcal{N}(\mathcal{C})$ suggest an analogy between spaces and categories:

$$points \leftrightarrow objects$$

$$paths \leftrightarrow morphisms$$

$$homotopies \text{ between } paths \leftrightarrow ???$$

-Topological categories are one way of formalizing “homotopies between morphisms” as “paths between points in a mapping space”

-There are other ways of formalizing the same idea.

-The one that we will focus on is that of ∞ -**category** or **quasicategories**.

-The idea: identify a class of simplicial sets which are close enough to being of the form $\mathcal{N}(\mathcal{C})$ to “do category theory with them” but which also include simplicial sets of the form $\Pi(X)$.

Q: Given $S. \in \mathbf{Set}_\Delta$, 1) how can we tell if $S. \cong \mathcal{N}(\mathcal{C})$ for some category \mathcal{C} ? and 2) how can we tell if $S. \cong \Pi(X)$ for some space X ?

Back to $\mathcal{N}(\mathcal{C})$: explicitly $\mathcal{N}(\mathcal{C})_2 = \{A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} A_2\}$, $\mathcal{N}(\mathcal{C})_1 = \{A_0 \xrightarrow{f_1} A_1\}$, and $\mathcal{N}(\mathcal{C})_0 = \{A_0\}$.

$$d_0 : \mathcal{N}(\mathcal{C})_1 \rightarrow \mathcal{N}(\mathcal{C})_0, (A_0 \xrightarrow{f_1} A_1) \mapsto A_1$$

$$d_1 : \mathcal{N}(\mathcal{C})_1 \rightarrow \mathcal{N}(\mathcal{C})_0, (A_0 \xrightarrow{f_1} A_1) \mapsto A_0$$

$$d_0 : \mathcal{N}(\mathcal{C})_2 \rightarrow \mathcal{N}(\mathcal{C})_1, (A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} A_2) \mapsto (A_1 \xrightarrow{f_2} A_2)$$

$$d_2 : \mathcal{N}(\mathcal{C})_2 \rightarrow \mathcal{N}(\mathcal{C})_1, (A_0 \xrightarrow{f_1} A_1 \xrightarrow{f_2} A_2) \mapsto (A_0 \xrightarrow{f_1} A_1)$$

Observation: the diagram

$$\begin{array}{ccc} \mathcal{N}(\mathcal{C})_2 & \xrightarrow{d_2} & \mathcal{N}(\mathcal{C})_1 \\ \downarrow d_0 & & \downarrow d_0 \\ \mathcal{N}(\mathcal{C})_1 & \xrightarrow{d_1} & \mathcal{N}(\mathcal{C})_0 \end{array}$$

is Cartesian (i.e. $\mathcal{N}(\mathcal{C})_2 \simeq \mathcal{N}(\mathcal{C})_1 \times_{\mathcal{N}(\mathcal{C})_0} \mathcal{N}(\mathcal{C})_1$)

Note: for any $S. \in \mathbf{Set}_\Delta$ we have a commutative diagram

$$\begin{array}{ccc} S_2 & \xrightarrow{d_2} & S_1 \\ \downarrow d_0 & & \downarrow d_0 \\ S_1 & \xrightarrow{d_1} & S_0 \end{array}$$

but it’s usually not Cartesian.

10. 9/17/2021

Question: How do we characterize simplicial sets of the form $N(\mathcal{C})$? Furthermore, how do we characterize of the form $\Pi(X)$?

To answer these two questions, we need to introduce the notion of a *horn*.

Definition 10.1. (1) For $n \in \mathbb{N}$, define

$$\Delta^n = \text{Hom}_{\Delta}(-, [n]) \in \text{Set}_{\Delta} := \text{Fun}(\Delta^{op}, \text{Set})$$

(2) For $j \in [n]$, define $\Lambda_j^n \in \text{Set}_{\Delta}$ by

$$\begin{aligned} (\Lambda_j^n)_m &:= \{\text{order-perserving } p : [m] \rightarrow [n] \text{ such that } \{j\} \bigcup p([m]) \neq [n]\} \\ (10.2) \quad &\subset \{\text{order-perserving } p : [m] \rightarrow [n]\} \\ &= (\Delta^n)_m \end{aligned}$$


with face and degeneracy maps obtained by the restriction from Δ^n .

(3) In particular, we have a canonical monomorphism $\Lambda_j^n \hookrightarrow \Delta^n$ in Set_{Δ} . We call Λ_j^n the j^{th} *horn* of Δ^n .


(4) If $0 < j < n$, then we call Λ_j^n an *inner horn*.

Remark. For any $K \in \text{Set}_{\Delta}$, we have $K_n := \text{Hom}_{\text{Set}_{\Delta}}(\Delta^n, K)$ by the *Yoneda lemma*.


Example. Let's picture the elements of $[2]$ as the vertices of $|\Delta^2|$:




Then, we can picture the 1-simplices of Δ^2 and its horns. $(\Delta^2)_1 = \{p_2, p_1, p_0, q_0, q_1, q_2\}$ is a six-element set containing the following order-preserving maps:




We call the three elements in the bottom row “degenerate simplices” because they are in the image of a degeneracy map. By definition of $(\Lambda_0^2)_1$, one can check that it is the following five maps:



Similarly, $(\Lambda_1^2)_1$ is:



Finally, $(\Lambda_2^2)_2$ is:



Proposition. (HTT 1.1.2.2) $K \in \text{Set}_\Delta$ is of the form $N(\mathcal{C})$ for some category \mathcal{C} if and only if whenever we have a morphism $\Lambda_j^n \rightarrow K$ with $0 < j < n$, there exists a *unique* morphism $\Delta^n \rightarrow K$ such that the following diagram commutes in Set_Δ :

$$\begin{array}{ccc} \Lambda_j^n & \hookrightarrow & \Delta^n \\ & \searrow & \downarrow \exists! \\ & & K \end{array}$$

Example. Let's see why a morphism $\Lambda_1^2 \xrightarrow{\phi} N(\mathcal{C})$ satisfies this condition. On 1-simplices, ϕ is a function of the form:

$$p_2 \xrightarrow{\phi} (A_0 \xrightarrow{f_1} A_1)$$

$$p_0 \xrightarrow{\phi} (A_1 \xrightarrow{f_2} A_2)$$

$$q_0 \xrightarrow{\phi} (A_0 \xrightarrow{id_{A_0}} A_0)$$

\vdots

The fact that ϕ is compatible with composition in Δ implies that the target of f_1 is equal to the source of f_2 . Furthermore, it also means that degenerate simplices go to identity morphisms.

A morphism $\Delta^2 \xrightarrow{\bar{\phi}} N(\mathcal{C})$ is determined by $(\Delta^2)_2 \ni |\Delta^2| \mapsto (B_0 \xrightarrow{g_1} B_1 \xrightarrow{g_2} B_2) \in N(\mathcal{C})_2$. Since $\bar{\phi}$ is compatible with face maps, we have

$$p_2 \xrightarrow{\bar{\phi}} (B_0 \xrightarrow{g_1} B_1)$$

$$p_0 \xrightarrow{\bar{\phi}} (B_1 \xrightarrow{g_2} B_2)$$

$$p_1 \xrightarrow{\bar{\phi}} (B_0 \xrightarrow{g_2 \circ g_1} B_2)$$

If $\bar{\phi}$ extends ϕ , this forces $A_0 = B_0$, $A_1 = B_1$, $A_2 = B_2$, $f_1 = g_1$, $f_2 = g_2$. So, $\bar{\phi}$ is unique if it exists.

11. 9/20/2021

We continue the discussion of characterizing nerves by horn-filling property. That is,

Proposition 11.1. (*HTT 1.1.2.2*) *Let $K \in \text{Set}_\Delta$ is of the form $N(\mathcal{C}) \Leftrightarrow$ Every j -th n -horn $\Lambda_j^n \rightarrow K$ with $0 < j < n$ extends uniquely to an n -simplex $\Delta^n \rightarrow K$*

Proof. (Idea) "Only If" direction: Define $Ob(\mathcal{C}) = K_0$, $Mor(\mathcal{C}) = K_1$, and source/target map $Mor(\mathcal{C}) \xrightarrow[t]{s} Ob(\mathcal{C})$ as degeneracy maps $K_1 \xrightarrow[t]{s} K_0$ and their composition from unique morphism extension property of the horn. That is, we have two different j -th n -horns and an unique extension to n -simplex defines a composition.

Q: We have witnessed characterization of nerve, which about the others? How to characterize when K is of the form $\Pi(X)$?

A: To address the question, it would be natural to construct a functor from topological space to simplicial sets. the following construction will be useful:

Proposition 11.2. *The functor $Top \xrightarrow{\Pi(\cdot)} \text{Set}$ denoted by $X \mapsto \Pi(X)$ has a left adjoint $\text{Set}_\Delta \xrightarrow{|\cdot|} Top$ denoted by $K \mapsto |K|$, we call $|K|$ the geometric realization of K*

Recall: this means for all $K \in \text{Set}_\Delta$ and $X \in Top$ we have a bijection between

$$Hom_{\text{Set}_\Delta}(K, \Pi(X)) \cong Hom_{Top}(|K|, X)$$

and these are compatible with composition.

Proof. (Idea) (Lurie, Goerss and Jardine, *Simplicial Homotopy Theory* Ch.1)

Define $\Delta \downarrow K$ a new category, the category of simplicies in K , by settings the objects to be

$$\sqcup_{n \in \mathbb{N}} K_n \text{ and } Hom_{\Delta \downarrow K}(\sigma, \tau) := \left\{ \begin{array}{ccc} \Delta^n & \xrightarrow{\sigma} & K \\ \downarrow \Theta & \nearrow \tau & \\ \Delta^m & & \end{array} \right\} \text{ for all } \Theta, \text{ where } \sigma, \tau \in K_n, K_m.$$

Remark 11.3. This set $Hom_{\Delta \downarrow K}(-, -)$ is a subset of morphism of $Hom_{Set_{\Delta}}(\Delta^n, \Delta^m) \cong Hom_{poset}([m], [n])$

Remark 11.4. This is really a general construction that takes any functor $F : \mathcal{C} \rightarrow Set$ to a category $\mathcal{C}^{op} \downarrow F$

Note there is a forgetful functor $\Delta \downarrow K \rightarrow Set_{\Delta}$ that takes $\Delta^n \xrightarrow{\sigma} K$ to Δ^n

Now we want to take the colimit. The colimit of the diagram recover the original simplicial sets K , we give a general categorical fact (construction), in the spirit of Yoneda Lemma:

$$K \cong colim_{\Delta^n \xrightarrow{\sigma} K \text{ in } \Delta \downarrow K} \Delta^n$$

Now let's define the geometric realization by:

$$|K| \cong colim_{\Delta^n \xrightarrow{\sigma} K \text{ in } \Delta \downarrow K} |\Delta^n|$$

We can obtain what we want, the idea is to "glue" the simplicies together encoded in the structure in K , then we get some space.

For any space $X \in Top$ we have

$$\begin{aligned} Hom_{Top}(|K|, X) &\cong Hom_{Top}(colim_{\Delta^n \downarrow K} \Delta^n, X) \\ &\cong lim_{\Delta^n \downarrow K} Hom_{Top}(|\Delta^n|, X) \\ &\cong lim_{\Delta^n \downarrow K} Hom_{Set_{\Delta}}(|\Delta^n|, \Pi(X)) \\ &\cong Hom_{Set_{\Delta}}(colim_{\Delta^n \downarrow K} \Delta^n, \Pi(X)) \\ &\cong Hom_{Top}(|K|, \Pi(X)) \end{aligned}$$

Thus we are done.

12. 9/22/2021

Last time we talked about geometric realization. Roughly, given a simplicial set $K \in Set_{\Delta}$, we can get $|K| = \sqcup K_n \times |\Delta^n| / \text{gluing}$.

Proposition 12.1. $|-|$ is left adjoint to $\Pi(-)$.

- Note that $|\Delta^n|$ is the geometric realization of Δ^n , as $\Delta \downarrow \Delta^n$ has a final object $\Delta^n \xrightarrow{id} \Delta^n$. Then we have $|\Delta^n| = colim_{\Delta^k \rightarrow \Delta^n} |\Delta^k| = |\Delta^n|$.

- Similarly $|\Lambda_j^n|$ is isomorphic to the colimits over just its non-degenerate $(n-1)$ -simplices and their facets.

Example 12.2. $|\Lambda_0^2|$

- Note that $|\Lambda_0^2| \rightarrow |\Delta^2|$ is a retract: there is a continuous map $|\Delta^2| \rightarrow |\Lambda_0^2|$ s.t. $|\Lambda_0^2| \rightarrow |\Delta^2| \rightarrow |\Lambda_0^2|$ is the identity.

- Fact: The same is true of $|\Lambda_j^n| \rightarrow |\Delta^n|$ for any $0 \leq j \leq n$.

Corollary 12.3. *For any space X , any morphism $|\Lambda_j^n| \rightarrow X$ ($0 \leq j \leq n$) extends to a morphism $|\Delta^n| \rightarrow X$.*

Corollary 12.4. *For any space X , any morphism $\Lambda_j^n \rightarrow \Pi(X)$ ($0 \leq j \leq n$) extends to a morphism $\Delta^n \rightarrow \Pi(X)$.*

The proof is by adjunction.

Definition 12.5. $K \in \text{Set}_\Delta$ is a Kan complex if it satisfies the extension condition in the above corollary.

- It is not true that every Kan complex is of the form $\Pi(X)$, but this is true up to homotopy.

- Recall that continuous maps $f, g : X \rightarrow Y$ are homotopic if there exists $h : X \times [0, 1] \rightarrow Y$

s.t. the diagram

$$\begin{array}{ccc}
 X \times \{0\} & & \\
 \downarrow & \searrow f & \\
 X \times [0, 1] & \xrightarrow{h} & Y \\
 \uparrow & \nearrow g & \\
 X \times \{1\} & &
 \end{array}$$

commutes.

Exercise: All limits and colimits exists in Set_Δ and are computed objectwise.

Definition 12.6. Two morphisms $f, g : J \rightarrow K$ in Set_Δ are homotopic if $\exists h : J \rightarrow \Delta^1 \rightarrow K$

such that

$$\begin{array}{ccc}
 J \times \Delta^0 & & \\
 \downarrow id \times s_0 & \searrow f & \\
 J \times \Delta^1 & \xrightarrow{h} & K \\
 \uparrow id \times s_1 & \nearrow g & \\
 J \times \Delta^0 & &
 \end{array}$$

commutes.

Fact: $\Pi(-)$ and $|-|$ take homotopic maps to homotopic maps.

13. 9/24/2021

Last time: We discussed Kan complex, homotopies.

Let $\mathbf{Kan} \subset \mathbf{Set}_\Delta$ be the full subcategory of Kan complexes and $H_0(\mathbf{Kan})$ the category with the same objects but

$$\text{Hom}_{H_0(\mathbf{Kan})}(J, K) := \text{Hom}_{\mathbf{Kan}}(J, K) / \sim, \text{ where } f \sim g \text{ if } f \text{ and } g \text{ are homotopic.}$$

Remark 13.1. The implicit proposition behind the definition is that simplicial homotopy is an equivalent relation and compatible with composition.

Similarly, let $CW \subset \mathbf{Set}$ be the full subcategory of CW -complexes, (for example, manifolds, $|K|$ for any $K \in \mathbf{Set}_\Delta$) and define $H_0(CW)$ similarly. (so for example any contractible space is isomorphic to a point in $H_0(CW)$.)

Theorem 13.2. *The adjoint functors $|-| : \mathbf{Set}_\Delta \rightarrow \mathbf{Top}$ and $\Pi(-) : \mathbf{Top} \rightarrow \mathbf{Set}_\Delta$ respect homotopy equivalence of morphisms and their restrictions induce inverse equivalences between $H_0(\mathbf{Kan})$ and $H_0(CW)$.*

Corollary 13.3. *For any $K \in \mathbf{Kan}$, the canonical morphism $K \rightarrow \Pi(|K|)$ is invertible up to homotopy.*

Terminology: We often just call $H_0(CW) \simeq H_0(\mathbf{Kan})$ the homotopy category of spaces.

Now we state a key definition:


Definition 13.4. $K \in \mathbf{Set}_\Delta$ is an ∞ -category if every morphism $\Lambda_j^n \rightarrow K$ with $0 < j < n$ extends to a morphism $\Delta^n \rightarrow K$.

Example 13.5. (1) $\mathcal{N}(\mathcal{C})$ is an ∞ -category for any category \mathcal{C} .

(2) Any Kan complex (e.g. $\Pi(X)$) is an ∞ -category.

If K is an ∞ -category, we refer to elements of K_0 as objects and elements of K_1 as morphisms (or 1-morphisms). Given a morphism $F \in K_1$ we call $d_1(f)$ its source and $d_0(f)$ its target.

Informally: In an ordinary category we can say " $\phi : A \rightarrow C$ is **the**(unique) composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ ", but in an ∞ -category we can only say " $\phi : A \rightarrow C$ is **a** composition of $f : A \rightarrow B$ and $g : B \rightarrow C$ ", and by this we mean there exists $h \in K_2$ such that $d_2(h), d_0(h) = g_1$ and $d_1(h) = \phi$. In pictures,



Now that we've defined ∞ -categories, our next task is to extend the key notions of ordinary (and topological) category theory.

Opposites: If \mathcal{C} is a category, \mathcal{C}^{op} has the same objects but

$$\mathrm{Hom}_{\mathcal{C}^{op}}(X, Y) := \mathrm{Hom}_{\mathcal{C}}(Y, X)$$

Definition 13.6. If \mathcal{C} is an ∞ -category, we define \mathcal{C}^{op} by setting $\mathcal{C}_n^{op} := \mathcal{C}_n$ for all $n \in \mathbb{N}$, and setting

$$\begin{aligned} (d_j : \mathcal{C}_n^{op} &\rightarrow \mathcal{C}_{n-1}^{op} := d_{n-j} : \mathcal{C}_n \rightarrow \mathcal{C}_{n-1}) \\ (s_j : \mathcal{C}_n^{op} &\rightarrow \mathcal{C}_{n+1}^{op} := s_{n-j} : \mathcal{C}_n \rightarrow \mathcal{C}_{n+1}) \end{aligned}$$

14. 9/27/2021

Last time: We defined ∞ -category.

Homotopy Categories (HTT 1.2.3)

Let $K \in \mathrm{Set}_{\Delta}$ and $f, g \in K$ be 1-simplices with the same faces i.e. $d_0(f) = d_0(g)$ and $d_1(f) = d_1(g)$.

Definition 14.1. f and g are homotopic if there exists $h \in K_2$ such that $d_2(h) = f$, $d_1(h) = g$, and $d_0(h) = s_0(d_0(f)) = \mathrm{Id}_{d_0(f)}$ if K is an ∞ -category.

$$\begin{array}{ccc} & d_0(f) & \\ & \bullet & \\ f \nearrow & & \searrow \mathrm{Id}_{d_0(f)} \\ \bullet & h & \bullet \\ g \rightarrow & & \end{array}$$

$d_1(f) = d_1(g)$ $d_0(g) = d_0(f)$

Theorem 14.2. If K is an ∞ -category, homotopy defines an equivalence relation on K_1 . Moreover, there exists a unique category hK (the homotopy category of K) such that

$$\begin{aligned} \mathrm{Ob}(hK) &:= K_0 \text{ and} \\ \mathrm{Mor}(hK) &:= K_1 / \text{homotopy} \end{aligned}$$

and such that the natural functions

$$\begin{aligned} K_0 &\simeq N(hK)_0 \text{ and} \\ K_1 &\mapsto N(hK)_1 \text{ i.e. the natural quotient map,} \end{aligned}$$

extend to a morphism $K \rightarrow N(hK)$ in Set_{Δ} (which is unique since the nerve of a category is determined by 0 and 1 simplices.)

Remark 14.3. The "functor" $h(-) : \mathrm{Cat}_{\infty} \rightarrow \mathrm{Cat}_1$ is a right adjoint of $N(-) : \mathrm{Cat}_1 \rightarrow \mathrm{Cat}_{\infty}$.

Informally, compositions in an ∞ -category are not unique, but they are unique up to homotopy equivalence.

Definition 14.4. A morphism in an ∞ -category K is an isomorphism (or equivalence) if its image in hK is an isomorphism in the usual sense.

Equivalently a morphism $f : X \rightarrow Y$ in K is an isomorphism if there exists a morphism $f^{-1} : Y \rightarrow X$ and a 2-simplex $h \in K_2$ such that

$$\begin{array}{ccc}
 & Y & \\
 f \nearrow & & \searrow f^{-1} \\
 X & \xrightarrow{Id_X := S_0(X)} & X
 \end{array}$$

Proposition 14.5. An ∞ -category K is a Kan complex if and only if every morphism in K is an isomorphism.

Mapping Spaces (HTT 1.2.2)

Part of the analogy comparing ∞ -categories to topological categories is that we can still define a mapping space between objects in an ∞ -category, but it's only canonical up to homotopy equivalence.

Notation: given a subset of the interval $I \subset [n]$, let $\Delta^I \subset \Delta^n$ be the largest simplicial subset with $\Delta_0^I = I$.

$$\begin{array}{ccc}
 \begin{array}{ccc}
 & 2 & \\
 \nearrow & & \searrow \\
 1 & \xrightarrow{\quad} & 3
 \end{array} & \subset & \begin{array}{ccc}
 & 1 & 3 \\
 \nearrow & \xrightarrow{\quad} & \searrow \\
 0 & \xrightarrow{\quad} & 2
 \end{array}
 \end{array}$$

15. 9/29/2021

Last time: homotopy category

Next up: mapping space

Notation: given $I \subset [n]$, let $\Delta^I \subset \Delta^n$ be the largest simplicial subset with $\Delta_0^I = I \subset [n] = \Delta_0^n$ (abstractly $\Delta^I \simeq \Delta^{|I|}$).

Definition 15.1. Let $K \subset Set_\Delta$ and $X, Y \in K_0$. Define $Hom_K^L \in Set_\Delta$ by

$$Hom_K^L(X, Y)_n = \text{maps } \phi : \Delta^{n+1} \rightarrow K \text{ such that the following diagrams commute:}$$

$$\begin{array}{ccc}
\Delta^{\{0\}} & \xrightarrow{\text{surj}} & \Delta^0 \\
\downarrow \text{incl} & & \downarrow \text{incl } X \\
\Delta^{n+1} & \xrightarrow{\phi} & K
\end{array}
\quad
\begin{array}{ccc}
\Delta^{\{1, \dots, n+1\}} & \xrightarrow{\text{surj}} & \Delta^0 \\
\downarrow \text{incl} & & \downarrow \text{incl } Y \\
\Delta^{n+1} & \xrightarrow{\phi} & K
\end{array}$$

Example 15.2. An element of $\text{Hom}_K^L(X, Y)_1$ is a 2-simplex $h : \Delta^2 \longrightarrow K$ of the following form

$$\begin{array}{ccc}
& & Y \ 1 \\
& \bullet & \\
& \nearrow & \searrow \text{id}_Y (= s_0(Y)) \\
X \ 0 & \xrightarrow{\phi} & 2 \ Y \\
& \bullet &
\end{array}$$

i.e. ϕ is a homotopy between a pair of 1-simplices ($d_2(\phi)$ and $d_1(\phi)$) with vertices X and Y .

Exercise. Define the face/degeneracy maps of $\text{Hom}_K^L(X, Y)$.

Proposition 15.3. *If K is an ∞ -category, then $\text{Hom}_K^L(X, Y)$ is a Kan complex. We call it the space of left morphisms from X to Y , and its image in $H_0(\text{Kan}) \simeq H_0(\text{CW})$ the mapping space from X to Y .*

Remark 15.4. The definition of $\text{Hom}_K^L(X, Y)$ was not canonical (i.e. it involved arbitrary choices), but it turns out to be canonical up to homotopy.

Functors (HTT Sec. 1.2.7)

Exercise. If C and D are ordinary categories, there is a canonical bijections between functors $C \longrightarrow D$ and morphisms $\mathcal{N}(C) \longrightarrow \mathcal{N}(D)$ in Set_Δ .

Definition 15.5. If J and K are ∞ -categories, we will call a morphism $J \longrightarrow K$ in Set_Δ a functor from J to K .

- In ordinary category theory, we define a category $\text{Fun}(C, D)$ whose objects are functors and whose morphisms are natural transformations.
- We generalize this construction to ∞ -categories as follows:

Definition 15.6. Given $J, K \in \text{Set}_\Delta$, let $\text{Map}_{\text{Set}_\Delta}(J, K) \in \text{Set}_\Delta$ be the functor $\Delta^{op} \longrightarrow \text{Set}$ that takes $[n]$ to $\text{Hom}_{\text{Set}_\Delta}(J \times \Delta^n, K)$, with structure maps defined by the Yoneda embedding $[n] \longrightarrow \Delta^n$.

Proposition 15.7. *If J and K are ∞ -categories, so is $\text{Map}_{\text{Set}_\Delta}(J, K)$. In this case we also write it as $\text{Fun}(J, K)$ and call it the ∞ -category of functors from J to K .*

Exercise. $\text{Fun}(J, K)_0 := \text{Hom}_{\text{Set}_\Delta}(J \times \Delta^0, K)$. But $J \times \Delta^0 \simeq J$ for any J , so this is just $\text{Hom}_{\text{Set}_\Delta}(J, K)$.

16. 10/1/2021

Last time: Mapping spaces, functors**Definition 16.1. (Proposition)** If J and K are ∞ -categories, the ∞ -category $\text{Fun}(J, K)$ of functors from J to K is defined by

$$\text{Fun}(J, K)_n := \text{Hom}_{\text{Set}_\Delta}(J \times \Delta^n, K).$$

Example 16.2. What is a 1-simplex $\phi \in \text{Fun}(J, K)$, explicitly?It is a morphism $J \times \Delta^1 \rightarrow K$ in Set_Δ . It defines functors $F, G: J \rightarrow K$ via

$$\begin{array}{ccc}
 J \simeq J \times \Delta^0 & & \\
 \downarrow id_J \times d_1 & \searrow F & \\
 J \times \Delta^1 & \xrightarrow{\phi} & K \\
 \uparrow id_J \times d_0 & \nearrow G & \\
 J \simeq J \times \Delta^0 & &
 \end{array}$$

Note that for any n we have (i) $(J \times \Delta^1)_n \cong J_n \times \Delta_n^1$ and (ii) $\phi_n: (J \times \Delta^1)_n \rightarrow K_n$. Recall that $K_1 = \text{Hom}_{\text{Set}_\Delta}(\Delta^1, K) \Rightarrow (\Delta^1)_1 = \text{Hom}(\Delta^1, \Delta^1)$, which contains id_{Δ^1} . For each $X \in J_0$, ϕ also defines a morphism $F(X) \xrightarrow{\phi_X} G(X)$ in K by the formula $\phi_X := \phi_1(s_0(X), id_{\Delta^1}) \in K_1$, where $id_X := s_0(X)$. We see that $F(x), G(x) \in K_0$, $\phi_X \in K_1$, and $d_1(\phi_X) = F(X)$, $d_0(\phi_X) = G(X)$.

Exercise. when $J \simeq N(\mathcal{C})$ and $K \simeq N(\mathcal{D})$, the morphisms ϕ_X define a natural transformation from F to G . This construction defines a bijection between $\text{Fun}(N(\mathcal{C}), N(\mathcal{C}))_1$ and natural transformations of functors from \mathcal{C} to \mathcal{D} .

16.1. Initial and Final Objects.

Definition 16.3. An object X in a category \mathcal{C} is **initial** (resp. **final**) if for every object Y , $\text{Hom}_{\mathcal{C}}(X, Y)$ (resp. $\text{Hom}_{\mathcal{C}}(Y, X)$) has a single element.

- I.e. “ $\text{Hom}_{\mathcal{C}}(X, Y)$ is a point”
- To extend this to the setting of ∞ -categories, we just reinterpret it as a statement about mapping spaces.

Definition 16.4. An object X of an ∞ -category K is **initial** (resp. **final**) if for every object Y , the mapping space $\text{Map}_K(X, Y)$ (resp. $\text{Map}_K(Y, X)$) is contractible.

Limits and Colimits

- Let $p: I \rightarrow \mathcal{C}$ be a functor between ordinary categories.
- The **overcategory** of p is the category $\mathcal{C}/_p$ with objects $\text{Ob}(\mathcal{C}/_p) = \{(x, \{f_i: X \rightarrow x_i := p(i)\}_{i \in I})\}$ such that

$$\begin{array}{ccc} X & \longrightarrow & X_j \\ & \nearrow p(g) & \\ & X_i & \end{array}$$

commutes for every $g \in \text{Hom}_I(i, j)$ and all $i, j \in I$, for $x \in \text{Ob}(\mathcal{C})$ and $f_i \in \text{Mor}(\mathcal{C})$. And $\mathcal{C}/_p$ has morphisms $\text{Hom}_{\mathcal{C}/_p}((x, \{f_i\}), (x', \{f'_i\})) = \{\phi \in \text{Hom}_{\mathcal{C}}(X, X') \text{ such that}$

$$\begin{array}{ccc} X & \xrightarrow{\phi} & X' \\ r_i \searrow & & \nearrow r'_i \\ & X_i & \end{array}$$

commutes for all $i \in I$.

Definition 16.5. A **limit** of p is a final object of the overcategory $\mathcal{C}/_p$.

Remark 16.6. if $(X, \{f_i\})$ is a limit of p_i we often just say “ X is a limit of p ” and leave the $\{f_i\}$ implicit.

17. 10/4/2021

Last time: ∞ -category $\text{Fun}(J, K)$, initial and final objects, limits and colimits

Now we need to figure out suitable definitions for initial and final objects as well as limits and colimits for ∞ -categories. We cannot directly duplicate these definition from ordinary categories because the definition of morphisms in the overcategory requires us to refer to the composition of two morphisms and check that it agrees with another. However, in an ∞ -category the composition of two morphisms is not defined uniquely but only up to homotopy.

The limit of an ∞ -category \mathcal{C} as an ∞ -category is generally more interesting than the ordinary limit of the its homotopy category $h(\mathcal{C})$. The limit will incorporate information from all the simplices of \mathcal{C} rather than just the 0-simplices and 1-simplices used in the limit of $h\mathcal{C}$. We give an explicit example of a computation of a limit in the usual category sense using the overcategory $\mathcal{C}/_p$ to help motivate the definition we will see next time for ∞ -categories.

We will use the following example throughout the next few lectures. Let the category I be

$$I = \begin{array}{ccc} & \bullet & b \\ & \downarrow & bc \\ a \bullet & \xrightarrow{ac} & \bullet c \end{array}$$

Explicitly the objects of I are $\text{Ob}(I) = \{a, b, c\}$ and the morphisms of I are $\text{Mor}(I) = \{ac, bc, aa, bb, cc\}$. Let $p : I \rightarrow \mathcal{C}$. Then by definition the set of objects of the overcategory \mathcal{C}/p is

$$\text{Ob}(\mathcal{C}/p) = \left\{ (X, f_a, f_b) \text{ such that } \begin{array}{ccc} X \bullet & \xrightarrow{f_b} & \bullet p(b) \\ f_a \downarrow & & \downarrow p(bc) \\ p(a) \bullet & \xrightarrow{p(ac)} & \bullet p(c) \end{array} \text{ commutes} \right\}.$$

Note an object of \mathcal{C}/p also has a map $f_c : X \rightarrow p(c)$ but f_c is determined by f_a and f_b , so we omit it to simplify the diagrams. We now also can explicitly write the morphisms in \mathcal{C}_p as

$$\text{Mor}(\mathcal{C}/p) = \left\{ (X', f'_a, f'_b, \phi) \text{ such that } \begin{array}{ccc} X' \bullet & \xrightarrow{f'_b} & \bullet p(b) \\ \phi \swarrow & \searrow f_b & \\ X \bullet & \xrightarrow{f_b} & \bullet p(b) \\ f'_a \downarrow & & \downarrow p(bc) \\ p(a) \bullet & \xrightarrow{p(ac)} & \bullet p(c) \end{array} \text{ commutes} \right\}.$$


Then by definition a final object of \mathcal{C}/p is a commutative diagram

$$\begin{array}{ccc} X \bullet & \xrightarrow{f_b} & \bullet X_b \\ f_a \downarrow & & \downarrow \\ X_a \bullet & \xrightarrow{\quad} & \bullet X_c \end{array}$$

such that given any other commutative diagram


$$\begin{array}{ccc} X' \bullet & \xrightarrow{f'_b} & \bullet X'_b \\ f'_a \downarrow & & \downarrow \\ X'_a \bullet & \xrightarrow{\quad} & \bullet X'_c \end{array}$$

there exists a unique morphism $\phi \in \text{Mor}(\mathcal{C}_{/p})$ with $\phi : X' \rightarrow X$. This means that the diagram



commutes. Then the final object in $\mathcal{C}_{/p}$ is exactly the usual pullback along the maps $p(ac)$ and $p(bc)$. Thus we have realized the pullback as a limit over a diagram of shape I .

We cannot just replace "category" by " ∞ -category" in the definition of the overcategory. The main problem is that composition is not defined uniquely in ∞ -categories. Instead, we should replace the condition that $p(g)' \circ f_i = f_j$ in the condition on objects in the overcategory with the condition that there exists a 2-simplex whose boundary one direction is $p(g) \circ f_i$ and the boundary the other direction is f_j for every $g : i \rightarrow j$. As a diagram this is



where the shaded region in the middle is the 2-simplex whose existence is a condition for an object (X, f_i, f_j) to be in the overcategory. This idea will be made more rigorous in the next few sections using the idea of a join.

18. 10/6/2021

Last time we defined limits and colimits in ordinary categories. In order to define limits and colimits in ∞ -categories, like before, we want to first write down a suitable definition of overcategories; that requires the notion of a *join*. By convention, for all $K \in \text{Set}_\Delta$, define $K_{-1} = \text{pt}$.

Definition 18.1. Given $J, K \in \text{Set}_\Delta$, the join $J * K \in \text{Set}_\Delta$ is defined by setting

$$\begin{aligned}
(J * K)_n &= \coprod_{m=-1}^n J_m \times K_{n-m-1} = \coprod_{\substack{m+l=n-1 \\ -1 \leq m, l \leq n}} J_m \times K_l \\
&= (J_{-1} \times K_n) \amalg (J_0 \times K_{n-1}) \amalg \cdots \amalg (J_n \times K_{-1}) \\
&= K_n \amalg (J_0 \times K_{n-1}) \amalg \cdots \amalg J_n,
\end{aligned}$$

and for $(j, k) \in J_m \times K_{n-m-1}$ setting the degenercy and face maps to be

$$d_i(j, k) = \begin{cases} (d_i(j), k) \in J_{m-1} \times K_{n-m-1} & i \leq m \\ (j, d_{i-m-1}(k)) \in J_m \times K_{n-m-2} & i \geq m+1 \end{cases}$$

Similarly for $s_i(j, k)$.

Note that J and K are naturally simplicial subsets of $J * K$.

Definition 18.2. Given categories \mathcal{C} and \mathcal{D} , the join $\mathcal{C} * \mathcal{D}$ is the category defined by

$$\begin{aligned}
\text{Ob}(\mathcal{C} * \mathcal{D}) &:= \mathcal{C} \amalg \mathcal{D} \\
\text{Hom}_{\mathcal{C} * \mathcal{D}}(X, Y) &:= \begin{cases} \text{Hom}_{\mathcal{C}}(X, Y) & X, Y \in \mathcal{C}; \\ \text{Hom}_{\mathcal{D}}(X, Y) & X, Y \in \mathcal{D}; \\ \text{pt} & X \in \mathcal{C}, Y \in \mathcal{D}; \\ \emptyset & X \in \mathcal{D}, Y \in \mathcal{C}. \end{cases}
\end{aligned}$$

Then $\mathcal{N}(\mathcal{C} * \mathcal{D}) = \mathcal{N}(\mathcal{C}) * \mathcal{N}(\mathcal{D})$.

Proposition 18.3. If J and K are ∞ -categoriues, so is $J * K$.

Remark 18.4. We can rephrase our definition of overcategory $\mathcal{C}_{/p}$ in terms of joins:

- Regard $[n]$ as a certain category with at most one morphism between any two objects,
- The objects of $\mathcal{C}_{/p}$ are functors $[0] * I \rightarrow \mathcal{C}$ whose restriction to I coincides with p .
- For example, if

$$\begin{array}{ccc}
I = & \begin{array}{ccc} & \bullet & b \\ & \downarrow & bc \\ a & \bullet & \xrightarrow{ac} \bullet & c \end{array} & [0] * I = \begin{array}{ccccc} & 0 & \bullet & \xrightarrow{\quad} & \bullet & b \\ & \downarrow & \searrow & & \downarrow & bc \\ a & \bullet & \xrightarrow{ac} & \bullet & c \end{array}
\end{array}$$

Then a diargam in \mathcal{C} of the form

$$\begin{array}{ccc}
X & \longrightarrow & p(b) \\
\downarrow & \searrow & \downarrow p(bc) \\
p(a) & \xrightarrow{p(ac)} & p(c)
\end{array}$$

is really a picture of a functor $[0] * I \rightarrow \mathcal{C}$ whose restriction to I is given by p .

- Similarly, a morphism in $\mathcal{C}_{/p}$ is a functor $[1] * I \rightarrow \mathcal{C}$ whose restriction to I coincides with p . In above example,

$$\begin{array}{ccccc}
& 0 & & & \\
& \bullet & \xrightarrow{\quad} & \bullet & \\
& \downarrow \scriptstyle 01 & \searrow & \downarrow & \\
[1] * I = & \bullet & \xrightarrow{\quad} & \bullet & b \\
& \downarrow & \searrow & \downarrow & bc \\
& \bullet & \xrightarrow{\quad} & \bullet & c \\
& a & \xrightarrow{ac} & &
\end{array}$$

So a diagram

$$\begin{array}{ccccc}
X' & \xrightarrow{\quad} & X & \longrightarrow & p(b) \\
\downarrow \scriptstyle \phi & \searrow & \downarrow & \searrow & \downarrow p(bc) \\
& & p(a) & \xrightarrow{p(ac)} & p(c)
\end{array}$$

is a picture of a functor $[1] * I \rightarrow \mathcal{C}$.

Definition 18.5. Let $p : J \rightarrow K$ be a morphism in \mathbf{Set}_Δ . Then $K_{/p} \in \mathbf{Set}_\Delta$ is defined by

$$(K_{/p})_n = \left\{ \phi \in \mathbf{Hom}_{\mathbf{Set}_\Delta}(\Delta^n * J, K) \text{ such that } \phi|_J \equiv p \right\}$$

with structure maps defined by the Yoneda embedding $\Delta^{\text{op}} \rightarrow \mathbf{Set}_\Delta$, $[n] \mapsto \Delta^n$.

Exercise.

Let $\mathcal{N}(p) : \mathcal{N}(I) \rightarrow \mathcal{N}(\mathcal{C})$ be the morphism associated to a functor $p : I \rightarrow \mathcal{C}$. Then

$$\mathcal{N}(\mathcal{C}_{/p}) \cong \mathcal{N}(K)_{/\mathcal{N}(p)}.$$

Proposition 18.6. If K is an ∞ -category, then so is $K_{/p}$.

Definition 18.7. If $p : J \rightarrow K$ is a morphism in \mathbf{Set}_Δ and K is an ∞ -category, we call a final object of $K_{/p}$ a **limit of p** .

19. 10/11/2021

Last time : We defined limits in ∞ -categories.

Definition 19.1. Let K and K' be ∞ -categories and $p : J \rightarrow K$ and $F : K \rightarrow K'$ be morphisms in \mathbf{Set}_Δ . Note F induces a functor $K_{/p} \xrightarrow{F} K'_{/F \circ p}$.

If $X \in \mathbf{Ob}(K_{/p})$ is a limit of p , then F preserves the limit of p if $F(X) \in \mathbf{Ob}(K'_{/F \circ p})$ is a limit of $F \circ p$.

Fundamental Case :

Recall that any ∞ -category has a canonical functor

$$\mathbf{Ho} : K \rightarrow \mathcal{N}(\mathbf{Ho}(K))$$

to the nerve of the homotopy category of K . Recall

$$\mathbf{Ob}(\mathbf{Ho}(K)) = \mathbf{Ob}(K), \mathbf{Mor}(\mathbf{Ho}(K)) = \mathbf{Mor}(K)/\text{homotopy equivalence}$$

Principle : \mathbf{Ho} often fails to preserve limits and colimits. When it fails to preserve the limit of some diagram $p : J \rightarrow K$, the limit of p is usually the more interesting object than the limit of $\mathbf{Ho} \circ p$.

Definition 19.2. (Partial) The category **Top** of topological spaces can be enhanced to an ∞ -category **Top** $_\infty$ with the following properties :

- $(\mathbf{Top}_\infty)_0 = \mathbf{Ob}(\mathbf{Top})$, i.e. spaces.
- $(\mathbf{Top}_\infty)_1 = \mathbf{Mor}(\mathbf{Top})$, i.e. continuous maps between spaces.
- 2-simplices of **Top** $_\infty$ with the boundary given as the diagram below is given by a continuous map $h : X \times [0, 1] \rightarrow Z$ such that $h|_{X \times \{0\}} = f_{YZ} \circ f_{XY}$ and $h|_{X \times \{1\}} = f_{XZ}$, i.e. a homotopy between continuous maps.

$$\begin{array}{ccc} X & \xrightarrow{f_{XY}} & Y \\ & \searrow f_{XZ} & \swarrow f_{YZ} \\ & Z & \end{array}$$

Corollary :

$$\mathbf{Ho}(\mathbf{Top}_\infty) \simeq \mathbf{Ho}(\mathbf{Top})$$

Example 19.3. Let $X \in \mathbf{Top}$ and $x_0 \in X$ be a base point. Let

$$I = \begin{array}{ccc} & \bullet & b \\ & \downarrow & bc \\ a & \bullet & \xrightarrow{ac} \bullet & c \end{array}$$

Consider the diagram $p : I \rightarrow \mathbf{Top}_\infty$ pictured as

$$\begin{array}{ccc} & \bullet & pt \\ & \downarrow & x_0 \\ pt & \bullet & \xrightarrow{x_0} \bullet & X \end{array}$$

What is the limit of $p : I \rightarrow \mathbf{Top}_\infty$ and $\tilde{p} : I \rightarrow \mathbf{Ho}(\mathbf{Top})$? Here \tilde{p} is the composition of p with the natural morphism $\mathbf{Top}_\infty \rightarrow \mathbf{Ho}(\mathbf{Top}_\infty) \simeq \mathbf{Ho}(\mathbf{Top})$

$$\mathrm{Ob}(\mathbf{Ho}(\mathbf{Top})_{/\tilde{p}}) = \left\{ \text{commutative diagrams } \begin{array}{ccc} Y & \xrightarrow{\quad} & pt \\ \downarrow & \searrow f & \downarrow x_0 \\ pt & \xrightarrow{x_0} & X \end{array} \text{ in } \mathbf{Ho}(\mathbf{Top}) \right\}$$

Note that the upper right triangle in the diagram is redundant so we can rewrite the objects of the overcategory as

$$\mathrm{Ob}(\mathbf{Ho}(\mathbf{Top})_{/\tilde{p}}) = \left\{ \text{commutative diagrams } \begin{array}{ccc} Y & & \\ \downarrow & \searrow f & \\ pt & \xrightarrow{x_0} & X \end{array} \text{ in } \mathbf{Ho}(\mathbf{Top}) \right\}$$

Notice the morphism $Y \rightarrow pt$ is uniquely defined and we have also fixed the morphism $pt \rightarrow X$ as x_0 , so every object is determined by the map f . Now the morphisms of this overcategory can be described as

$$\mathrm{Mor}(\mathbf{Ho}(\mathbf{Top})_{/\tilde{p}}) = \left\{ \text{commutative diagrams } \begin{array}{ccc} Y & & \\ \downarrow g & \searrow f & \\ Y' & \xrightarrow{f'} & X \end{array} \text{ in } \mathbf{Ho}(\mathbf{Top}) \right\}$$

Here the maps f and f' must be homotopic to the constant map x_0 . From the above information we see the final object in $\mathrm{Ho}(\mathbf{Top})_{/\bar{p}}$ is given by

$$\begin{array}{ccc} pt & & \\ \downarrow & \searrow^{x_0} & \\ pt & \xrightarrow{x_0} & X \end{array}$$

Now let us consider $p : I \rightarrow \mathbf{Top}_\infty$ then

$$\mathrm{Ob}(\mathbf{Top}_{\infty/p}) = \{ \phi \in \mathrm{Hom}_{\mathrm{Set}_\Delta}(\Delta^0 * \mathcal{N}(I), \mathbf{Top}_\infty), \text{ such that } \phi|_{\mathcal{N}(I)} = \mathcal{N}(p) \}$$

Note : Such a morphism ϕ is determined by where it sends non-degenerate simplices of $\Delta^0 * \mathcal{N}(I) = \mathcal{N}([0] * I)$ where $[0]$ is the category with only one object and the identity morphism. We can picture this as follows :

$$\begin{array}{ccc} 0 & \xrightarrow{0b} & b \\ \downarrow^{0a} & \searrow^{0c} & \downarrow^{bc} \\ a & \xrightarrow{ac} & c \end{array}$$

There are two 2-simplices $0bc$ and $0ac$.

20. 10/13/2021

Last time: extended example

-Given a space X and a point $x_0 \in X$, we can construct the limit of

$$\begin{array}{ccc} & pt & \\ & \downarrow^{x_0} & \\ pt & \xrightarrow{x_0} & X \end{array}$$

in $\mathrm{Ho}(\mathbf{Top})$ or \mathbf{Top}_∞ .

-Recall that a 2-simplex \mathbf{Top}_∞ is pictured as

$$\begin{array}{ccccc} & & \bullet & & Y \\ & \nearrow^{f_{XY}} & & \searrow^{f_{YZ}} & \\ X & \bullet & \xrightarrow{f_{XZ}} & \bullet & Z \end{array}$$

h

where X, Y, Z are spaces, f_{XY}, f_{YZ}, f_{XZ} are continuous maps and $h : X \times [0, 1] \rightarrow Z$ such that $h|_{X \times \{0\}} = f_{YZ} \circ f_{XY}$ and $h|_{X \times \{1\}} = f_{XZ}$.

-Let $p : \mathcal{N}(I) \rightarrow \mathbf{Top}_\infty$ be as above. We have

$$Ob(\mathbf{Top}_{\infty/p}) = \left\{ \phi \in \text{Hom}_{\text{Set}_\Delta}(\mathcal{N}([0] * I) \simeq \Delta^0 * \mathcal{N}(I), \mathbf{Top}_\infty) \text{ such that } \phi|_{\mathcal{N}(I)} \equiv p. \right\}$$

-Let's unpack $\mathcal{N}([0] * I)$: $I =$


$$\begin{array}{ccc} & \bullet & b \\ & \downarrow & bc \cdot [0] = \cdot^0 \\ a \bullet & \xrightarrow{ac} & \bullet c \end{array}$$

$$(\Delta^0 * \mathcal{N}(I))_0 = \{0, a, b, c\}$$

$$\begin{aligned} (\Delta^0 * \mathcal{N}(I))_1 &= \Delta_1^0 \cup (\Delta_0^0 \times \mathcal{N}(I)_0) \cup \mathcal{N}(I)_1 \\ &= \{0a, 0b, 0c, ac, bc\} + \text{degenerate 1-simplices} \end{aligned}$$

$$\begin{aligned} (\Delta^0 * \mathcal{N}(I))_2 &= \Delta_2^0 \cup (\Delta_1^0 \times \mathcal{N}(I)_0) \cup (\Delta_0^0 \times \mathcal{N}(I)_1) \cup \mathcal{N}(I)_2 \\ &= \{0ac, 0bc\} + \text{degenerate 1-simplices} \end{aligned}$$

-Thus we can picture $\Delta^0 * \mathcal{N}(I)$ as



-Thus let's see that

$$Ob(\mathbf{Top}_{\infty/p}) = \left\{ \text{diagrams } \begin{array}{ccc} Y & \xrightarrow{\quad} & pt \\ \downarrow h_2 & \searrow f & \downarrow x_0 \\ pt & \xrightarrow{x_0} & X \end{array} \text{ in } \mathbf{Top}_\infty \right\}$$


$$= \{(Y, f, h_1, h_2) \text{ where } f : Y \rightarrow X \text{ is a continuous}$$

$$\text{map and } h_1, h_2 : Y \times [0, 1] \rightarrow X \text{ such that}$$

$$h_1|_{Y \times \{0\}} = h_2|_{Y \times \{0\}} \equiv x_0 \text{ and } h_1|_{Y \times \{1\}} = h_2|_{Y \times \{1\}} = f\}$$

-The natural functor $\mathbf{Top}_{\infty/p} \rightarrow Ho(\mathbf{Top})_p$ forgets the data of h_1 and h_2 , and just remember the property that f was nullhomotopic.

-We can “glue h_1 and h_2 together” to get a continuous map $h : Y \times [0, 1] \rightarrow X$ such that $h|_{Y \times \{0\}} = h|_{Y \times \{1\}} \equiv x_0$. Here ’s the picture:



-Thus loses no information, so

$$Ob(\mathbf{Top}_{\infty/p}) = \{(Y, h) \text{ where } h : Y \times [0, 1] \rightarrow X \text{ is a map} \\ \text{such that } h|_{Y \times \{0\}} = h|_{Y \times \{1\}} \equiv x_0\}$$

-Aside: gives spaces X, Y , the space $Maps(X, Y)$ is characterized by there being (functorial) bijection

$$Hom_{Top}(Z, Maps(X, Y)) \simeq Hom_{Top}(Z \times X, Y)$$

for all spaces Z .

-Thus we also have

$$Ob(\mathbf{Top}_{\infty/p}) = \left\{ (Y, \bar{h}) \text{ where } \bar{h} : Y \rightarrow \Omega_{x_0} X \right\}$$

where the based loop space $\Omega_{x_0} X$ is the subspace of $Maps([0, 1], X)$ consisting of paths beginning and ending at x_0 .

(Ex: the fundamental group $\pi_1(X, x_0)$ is the set of connected components of $\Omega_{x_0} X$)

-If we follow the definition of morphisms $\mathbf{Top}_{\infty/p}$ through this, we find that

$$Mor(\mathbf{Top}_{\infty/p}) = \left\{ \text{commutative diagrams } \begin{array}{ccc} Y' & \xrightarrow{\bar{h}'} & \Omega_{x_0} X \\ \phi \downarrow & & \uparrow \bar{h} \\ Y & & \end{array} \right\}$$

-Thus the final object of $\mathbf{Top}_{\infty/p}$ is $\Omega_{x_0} X \xrightarrow{id} \Omega_{x_0} X$, because any map $Y \xrightarrow{\bar{h}} \Omega_{x_0} X$ extends uniquely to a commutative diagram

$$\begin{array}{ccc} Y & \xrightarrow{\bar{h}} & \Omega_{x_0} X \\ \bar{h} \downarrow & & \uparrow id \\ \Omega_{x_0} X & & \end{array}$$

\Rightarrow the limit of p in \mathbf{Top}_{∞} is $\Omega_{x_0} X$, which is much more interesting than just a point!

21. 10/18/2021

Last time we talked about limits and loop spaces.

Definition 21.1. Given a morphism $p : J \rightarrow K$ in \mathbf{Set}_{Δ} . The **undercategory** $K_{p/}$ of p is defined by $(K_{p/})_n = \{\phi \in Hom_{\mathbf{Set}_{\Delta}}(J * \Delta^n, K) \text{ such that } \phi|_J = p\}$

A **colimit** of p is an initial object of $K_{p/}$. Or sometimes, it is just the image of $O \in \Delta_0^0 \subset (J * \Delta^0)_0$ in K_0 under an initial object $\phi : J * \Delta^0 \rightarrow K$.

Exercise:

$$\Delta^0 * N \left(\begin{array}{ccc} & \bullet & \\ & \downarrow & \\ \bullet & \longrightarrow & \bullet \end{array} \right) \simeq \Delta^1 \times \Delta^1 \simeq N \left(\begin{array}{ccc} \bullet & \longrightarrow & \bullet \\ \downarrow & & \\ \bullet & & \end{array} \right) * \Delta^0$$

Terminology A diagram $\Delta^1 \times \Delta^1 \rightarrow K$ in an ∞ -category K is **Cartesian** or a **pullback square** if it is the limit of its restriction to $N \left(\begin{array}{ccc} & \bullet & \\ & \downarrow & \\ \bullet & \longrightarrow & \bullet \end{array} \right)$ and **coCartesian** or a **pushout square** if it is the colimit of its restriction to $N \left(\begin{array}{ccc} \bullet & \longrightarrow & \bullet \\ \downarrow & & \\ \bullet & & \end{array} \right)$

Example 21.2. Given a space X , let's compute the colimit of

$$\begin{array}{ccc} X & \longrightarrow & pt \\ \downarrow & & \\ pt & & \end{array} \quad \text{in } \mathbf{Top}_\infty$$


Following the analysis of last time, we have

$$\begin{aligned} Ob(\mathbf{Top}_\infty)_{p/} &= \{ \text{diagrams } \begin{array}{ccc} X & \longrightarrow & pt \\ \downarrow & \searrow^{h_2} & \downarrow \\ pt & \xrightarrow{h_1} & Y \end{array} \text{ in } \mathbf{Top}_\infty \} \\ &= \{ (Y, f, h_1, h_2) \text{ such that } f : X \rightarrow Y \text{ and } h_1, h_2 : X \times [0, 1] \rightarrow Y \\ &\quad \text{are continuous maps with } h_1|_{X \times \{0\}} = h_2|_{X \times \{0\}} = f \\ &\quad \text{and } h_1|_{X \times \{1\}}, h_2|_{X \times \{1\}} \text{ are constant maps} \} \\ &= \{ (Y, h) \text{ such that } h : X \times [0, 1] \rightarrow Y \text{ is continuous and such that} \\ &\quad h|_{X \times \{0\}} \text{ and } h|_{X \times \{1\}} \text{ are constant} \} \end{aligned}$$

$$Mor(Top_\infty)_{p/} = \left\{ \text{commutative diagrams } \begin{array}{ccc} & & Y \\ & \nearrow & \downarrow \\ X \times [0, 1] & & Y' \\ & \searrow & \end{array} \right\}$$

This implies that the initial object of $(\mathbf{Top}_\infty)_{p/}$ is the projection $X \times [0, 1] \rightarrow X \times [0, 1]/(p, 0) \sim (q, 0), (p, 1) \sim (q, 1)$ for all $p, q \in X$. This is called the **suspension** of X and is denoted ΣX .

Exercise $\Sigma S^n \simeq S^{n+1}$ for any $n \in \mathbb{N}$.



A similar computation to last time shows that the colimit of $\begin{array}{ccc} X & \longrightarrow & pt \\ \downarrow & & \\ pt & & \end{array}$ in $\mathbf{Ho}(\mathbf{Top})$ is

again just a point.

Fact: $X \simeq \Omega_{x_0} \Sigma X$ for any space X . This is a weak homotopy equivalence.

22. 10/20/2021

Last time: colimits and suspensions


Next up: loops and suspensions

To discuss the relationship between loop spaces and suspensions.

It is convenient to work with pointed spaces.

Definition 22.1. The ∞ -category Top_{∞}^+ of pointed spaces has
objects: $X_{\bullet} := (X, x_0)$ where X is a space and $x_0 \in X$ a point
morphisms: continuous maps $f : X \rightarrow Y$ such that $f(x_0) = y_0$.

2-simplices: pictured as



where $h : X \times [0, 1] \rightarrow Z$ is a continuous map such that

$$h|_{X \times \{0\}} = g \circ f, \quad h|_{X \times \{1\}} = \phi, \quad \text{and} \quad h|_{\{x_0\} \times [0, 1]} \equiv z_0.$$


(Equivalently: $Top_{\infty}^+ \simeq (Top_{\infty})_{p/}$ where $p : \Delta^0 \rightarrow Top_{\infty}$ is defined by $[0] \mapsto pt.$)

The computation from last time changes only slightly. If we define in Top_{∞}^+

$$\Sigma X_{\bullet} := colim \left(\begin{array}{ccc} X_{\bullet} & \longrightarrow & pt \\ \downarrow & & \\ pt & & \end{array} \right),$$

then ΣX_{\bullet} is the reduced suspension

Example 22.2.



- If X is nice, the reduced and ordinary suspensions are homotopy equivalent (So, we will often leave the word "reduced" implicit.).
- Unlike Top_∞ , there is a unique morphism in Top_∞^+ from pt to any pointed space X .
- In other words, the object pt is both initial and final in Top_∞^+ .

Definition 22.3. A zero object in an ∞ -category is an object which is both initial and final.

Example 22.4. $\{0\}$ is a zero object in the category of abelian groups.

Proposition 22.5. *If K is an ∞ -category, then full subcategory of zero objects is contractible (i.e. is homotopy equivalent to Δ^0) if it is non-empty.*

- If we now also define

$$\Omega X_\bullet := \lim \begin{pmatrix} & pt \\ & \downarrow \\ pt \longrightarrow & X_\bullet \end{pmatrix},$$

then the result is the same, but the data of this diagram only depends on an object C (in Top_∞^+), as opposed to an object plus a morphism (in Top_∞).

- In particular, the definitions of ΣX_\bullet and ΩX_\bullet are now formally symmetric.
- In fact, they are adjoint (in an "enhanced" sense): ... next time.

23. 10/22/2021

Last time: Pointed spaces and zero objects

Today Digression: The ∞ -category of ∞ -categories (and of spaces)

Definition 23.1. A **simplicial category** is a category enriched in simplicial sets. I.e. it is the data of

- a collection $Ob(\mathcal{C})$ of objects
- for each $X, Y \in Ob(\mathcal{C})$ a simplicial set $Map_{\mathcal{C}}(X, Y) \in Set_{\Delta}$
- for each $X, Y \in Ob(\mathcal{C})$ a morphism

$$Map_{\mathcal{C}}(X, Y) \times Map_{\mathcal{C}}(Y, Z) \rightarrow Map_{\mathcal{C}}(X, Z)$$

in Set_{Δ} , which are collectively associative.

We write Cat_{Δ} for the category of simplicial categories and simplicial functors.

Example 23.2. Given any $J, K \in Set_{\Delta}$, we defined $Map_{Set_{\Delta}}(J, K) \in Set_{\Delta}$ by the formula

$$Hom_{Set_{\Delta}}(\Delta^n, Map_{Set_{\Delta}}(J, K)) \cong Map_{Set_{\Delta}}(J, K)_n := Hom_{Set_{\Delta}}(\Delta^n \times J, K).$$

Exercise This construction gives Set_{Δ} the structure of a simplicial category.

Example 23.3. Let \mathcal{C} be a topological category. Then we defined $\Pi(\mathcal{C}) \in Cat_{\Delta}$ by setting $Ob(\Pi(\mathcal{C})) := Ob(\mathcal{C})$ and $Map_{\Pi(\mathcal{C})}(X, Y) := \Pi(Map_{\mathcal{C}}(X, Y))$ for all $X, Y \in Ob(\mathcal{C})$. This preserves the composition law since $\Pi(-)$ preserves products.

- We can now relate topological and ∞ -categories by defining a simplicial nerve:

$$Cat_{top} \xrightarrow{\Pi(-)} Cat_{\Delta} \xrightarrow{N(-)} Set_{\Delta}$$

Definition 23.4. Define a functor $\Delta \xrightarrow{C[-]} Cat_{\Delta}$ as follows. Given $0 \leq i \leq j$ let P_{ij} denote the partially ordered set

$$P_{ij} = \{I \subset [i, j] \subset \mathbb{N} \mid i, j \in I\}$$

ordered by inclusions. Then $Ob(C[n]) = \{0, \dots, n\}$ and for $i, j \in \{0, \dots, n\}$

$$Map_{C[n]}(i, j) = \begin{cases} \emptyset & i > j \\ N(P_{ij}) & i \leq j \end{cases}$$

For $i \leq j \leq k$ the composition rule

$$Map_{C[n]}(i, j) \times Map_{C[n]}(j, k) \rightarrow Map_{C[n]}(i, k)$$

using the natural map of partially ordered sets

$$P_{ij} \times P_{jk} \rightarrow P_{ik}$$

given by taking unions.


The simplicial nerves $N(\mathcal{C})$ of a simplicial category is then defined as before:

$$\begin{array}{ccccc} \Delta^{op} & \xrightarrow{C[-]} & Cat_{\Delta}^{op} & \xrightarrow{Hom_{Cat_{\Delta}}(-, \mathcal{C})} & Set \\ & \searrow & & \nearrow & \\ & & N(-) & & \end{array}$$

Example 23.5. Given $0 \leq i \leq j \leq n$, write $p_{ij} \in \text{Map}_{C[n]}(i, j)_0$ for the element corresponding to $\{i, j\} \subset [i, j]$. Then $p_{jk} \circ p_{ij} \neq p_{ik}$ in general, but they are related by a "universal homotopy". For example when $n = 2$, we have

$$P_{02} = \{\{0, 2\}, \{0, 1, 2\}\} = \{p_{02}, p_{12} \circ p_{01}\} \cong ([1] \text{ as a poset})$$

In a picture:



24. 10/25/2021

Proposition 24.1. If $\mathcal{C} \in \text{Cat}_\Delta$ has the property that $\text{Map}_{\mathcal{C}}(X, Y)$ is a Kan complex for all $X, Y \in \text{Ob}(\mathcal{C})$, then $N(\mathcal{C})$ is an ∞ -category.

Example 24.2. Recall the construction $\text{Cat}_{\text{top}} \xrightarrow{\Pi(-)} \text{Cat}_\Delta$. If $\mathcal{C} \in \text{Cat}_{\text{top}}$, then $\Pi(\mathcal{C})$ satisfies hypothesis of Proposition 24.1. So ∞ -category behaves as well as topological categories.

Definition 24.3. Let Top_{top} be the category of spaces (CGHaus, or CW-complexes) and space of maps between them, i.e. $\text{Map}_{\text{top}}(X, Y)$ is characterized by $\text{Hom}_{\text{top}}(Z, \text{Map}_{\text{top}}(X, Y)) \simeq \text{Hom}_{\text{top}}(X \times Z, Y)$. Then the ∞ -category of spaces is $\text{Top}_\infty = N(\Pi(\text{Top}_{\text{top}}))$.


Definition 24.4. Write Set_Δ^Δ for the simplicial category of simplicial sets. Let $\text{Cat}_\infty^\Delta \subseteq \text{Set}_\Delta^\Delta$ be the simplicial subcategory whose objects are ∞ -categories. This is the **simplicial category of ∞ -categories**. Let $\text{Cat}_\infty^{\Delta, \text{Kan}} \subseteq \text{Cat}_\infty^\Delta$ be the simplicial subcategory where for each J, K , we let $\text{Map}_{\text{Cat}_\infty^{\Delta, \text{Kan}}}(J, K)$ be the largest Kan complex in $\text{Map}_{\text{Cat}_\infty^\Delta}(J, K)$.

Now we define **∞ -category of ∞ -categories** to be $N(\text{Cat}_\infty^{\Delta, \text{Kan}})$.

Exercise: Let K be a simplicial set and $\{K_i\}_{i \in \mathcal{I}}$ be a collection of simplicial subsets which are all Kan complexes, then $\coprod_{j \in \mathcal{I}} K_j$ is again a Kan complex, so there is a well-defined largest Kan complex inside K .

25. 10/27/2021

Example 25.1. A 2-simplex in Cat_∞ is pictured as



where J, K, L are ∞ -categories, F_1, F_2, F_3 are functors, and ϕ is a natural isomorphism $\phi: F_2 \circ F_1 \Rightarrow F_3$. (i.e. ϕ is an isomorphism in the ∞ -category $\text{Fun}(J, L)$, which means ϕ is sent to isomorphism in the category $H_0(\text{Fun}(J, L))$.)

Important Fact([Lur09]): \mathbf{Cat}_∞ admits arbitrary limits and colimits.

Definition 25.2. Let $\mathbf{Top}_\infty \subseteq \mathbf{Cat}_\infty$ be the full subcategory whose objects are Kan complexes.

Fact. By construction, $\Pi(\mathbf{Top}_{\text{top}})$ is a simplicial category of $\mathbf{Cat}_\infty^{\Delta, \text{Kan}}$. The induced functor $N(\Pi(\mathbf{Top}_{\text{top}})) \rightarrow N(\mathbf{Cat}_\infty^{\Delta, \text{Kan}}) = \mathbf{Cat}_\infty$ factors through an equivalence $N(\Pi(\mathbf{Top}_{\text{top}})) \rightarrow \mathbf{Top}_{\text{top}} \rightarrow \mathbf{Top}_\infty \subseteq \mathbf{Cat}_\infty$, so our two definitions of \mathbf{Top}_∞ agree up to equivalence.

Definition 25.3. The ∞ -category $\mathbf{Cat}_{\text{ord}} \subseteq \mathbf{Cat}_\infty$ of ordinary categories (or 1-categories) is the full subcategory whose objects are (nerves of) ordinary categories.

26. 10/29/2021

Last time: \mathbf{Cat}_∞ and Yoneda

Theorem 26.1. Let \mathcal{C} be an ∞ -category.

(1) There exists a mapping space functor

$$\text{Map}_{\mathcal{C}}(-, -) : \mathcal{C}^{\text{op}} \times \mathcal{C} \rightarrow \mathbf{Top}_\infty$$

compatible with our previous constructions (i.e. $\text{Map}_{\mathcal{C}}(X, Y)$ is homotopy equivalent to $\text{Hom}_{\mathcal{C}}^L(X, Y)$ for all $X, Y \in \text{Ob}(\mathcal{C})$).

(2) (Yoneda) The induced functor $\mathcal{C} \rightarrow \text{Fun}(\mathcal{C}^{\text{op}}, \mathbf{Top}_\infty)$ is fully faithful (i.e. it induces homotopy equivalences of mapping spaces.)

Adjoint Functors

Definition 26.2. Let $\mathcal{C} \xrightleftharpoons[G]{F} \mathcal{D}$ be functors between ∞ -categories \mathcal{C} and \mathcal{D} . Then we say (F, G) are an adjoint pair (or F is a left adjoint of G , or G is a right adjoint of F) if there exists a natural transformation

$$u : \text{Id}_{\mathcal{C}} \longrightarrow G \circ F$$

such that for all $X \in \text{Ob}(\mathcal{C})$ and $Y \in \text{Ob}(\mathcal{D})$, the composition

$$\text{Map}_{\mathcal{D}}(F(X), Y) \xrightarrow[1]{\quad} \text{Map}_{\mathcal{C}}(G(F(X)), G(Y)) \xrightarrow[2]{\quad} \text{Map}_{\mathcal{C}}(X, G(Y)) \quad (*)$$

1: because G is a functor

2: compose with $u_X : X \rightarrow G(F(X))$

is a homotopy equivalence.

If \mathcal{C} and \mathcal{D} are ordinary categories, the mapping spaces in $(*)$ are discrete, and we recover the classical definition of adjoint functors.

We call u the unit of the adjunction. One can show there is a counit natural transformation $V : F \circ G \longrightarrow \text{Id}_{\mathcal{D}}$ which satisfies a similar property to u .

Theorem 26.3. Any two right adjoints of F are isomorphic, and the space of isomorphisms compatible with their unit is contractible.

Theorem 26.4. Write $LFun(\mathcal{C}, \mathcal{D}) \subset Fun(\mathcal{C}, \mathcal{D})$ for the full subcategory of left adjoints, similarly for $RFun(\mathcal{D}, \mathcal{C}) \subset Fun(\mathcal{D}, \mathcal{C})$. Then there is a canonical equivalence $LFun(\mathcal{C}, \mathcal{D}) \xrightarrow{\sim} RFun(\mathcal{D}, \mathcal{C})$ such that the image of any left adjoint is a right adjoint.

Back to Loops and Suspensions

Recall that a zero object in an ∞ -category \mathcal{C} is an object which is both initial and final.

Definition 26.5. Let \mathcal{C} be a pointed ∞ -category with zero object $0 \in Ob(\mathcal{C})$. We write $\Omega : \mathcal{C} \rightarrow \mathcal{C}$ and $\Sigma : \mathcal{C} \rightarrow \mathcal{C}$ for the functors given by

$$\Omega(X) = \lim \left(\begin{array}{ccc} & & 0 \\ & & \downarrow \\ 0 & \longrightarrow & X \end{array} \right) \quad \Sigma(X) = \operatorname{colim} \left(\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \\ 0 & & \end{array} \right)$$

if the needed limits and colimits exist.

Note: We are using a fact that we have not explicitly proved, which is that limits and colimits are functorial under morphisms of diagrams.

Exercise: If \mathcal{C} is an ordinary category, then $\Omega(X) \equiv 0$ and $\Sigma(X) \equiv 0$ for all $X \in Ob(\mathcal{C})$.

Theorem 26.6. Let \mathcal{C} be an ∞ -category in which Ω and Σ are well-defined. Then (Σ, Ω) is an adjoint pair.

27. 11/1/2021

Last time: Adjoint functors, loops, and suspensions.

Recall the following two definitions:

Definition 27.1. Functors $\mathcal{C} \xrightleftharpoons[G]{F} \mathcal{D}$ between ∞ -categories are *adjoint* if there exists functorial isomorphisms

$$\operatorname{Map}_{\mathcal{D}}(F(X), Y) \simeq \operatorname{Map}_{\mathcal{C}}(X, G(Y))$$

in Top_{∞} for all $X \in Ob(\mathcal{C})$, $Y \in Ob(\mathcal{D})$ (i.e. isomorphisms induced from a unit or counit transformation).

Definition 27.2. Let \mathcal{C} be a pointed ∞ -category. Then, the loop and suspension functors

$$\Omega_{\mathcal{C}}, \Sigma_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$$

are defined by

$$\Omega_{\mathcal{C}}(X) = \lim \left(\begin{array}{ccc} & & 0 \\ & & \downarrow \\ 0 & \longrightarrow & X \end{array} \right) \quad \Sigma_{\mathcal{C}}(X) = \operatorname{colim} \left(\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & & \\ 0 & & \end{array} \right)$$

provided these limits and colimits exist.

Convenient assumption: call a simplicial set *finite* if it has finitely many nondegenerate simplices.

Examples:

$$\Lambda_0^2 = N \left(\begin{array}{ccc} & & \cdot \\ & \downarrow & \\ \cdot & \longrightarrow & \cdot \end{array} \right) \quad \Lambda_2^2 = N \left(\begin{array}{ccc} \cdot & \longrightarrow & \cdot \\ \downarrow & & \\ \cdot & & \end{array} \right)$$

Definition 27.3. We say that an ∞ -category \mathcal{C} *admits finite limits* (resp. *colimits*) if any diagram $p : K \rightarrow \mathcal{C}$ with K finite has a limit (resp. colimit).

Theorem 27.4. Let \mathcal{C} be a pointed ∞ -category which admits finite limits and colimits. Then,

$$\Sigma_{\mathcal{C}}, \Omega_{\mathcal{C}}$$

are an adjoint pair.

Proof. (Sketch)

We claim that any $X, Y \in \text{Ob}(\mathcal{C})$, we have isomorphisms:

$$\text{Map}_{\mathcal{C}}(\Sigma(X), Y) \simeq \left\{ \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \longrightarrow & Y \end{array} \right\} \simeq \text{Map}_{\mathcal{C}}(X, \Omega(Y))$$

where the middle diagram is in $\text{Fun}(\Lambda^1 \times \Lambda^1, \mathcal{C})$.

To see the RHS, consider the following projection maps:

$$\left\{ \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \longrightarrow & Y \end{array} \right\} \leftarrow \left\{ \begin{array}{ccc} X & \xrightarrow{\quad} & \Omega(Y) \longrightarrow 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \xrightarrow{\quad} & Y \end{array} \right\} \rightarrow \{\text{diagrams } X \rightarrow \Omega(Y)\}$$

Both projections are surjective, so (fact from homotopy theory) they are homotopy equivalences if and only if their fibers are contractible.

To see the left arrow, if p is the diagram:

$$\begin{array}{ccc} & & 0 \\ & & \downarrow \\ 0 & \longrightarrow & Y \end{array}$$

then $\Omega(Y)$ is, by definition, the final object of \mathcal{C}/p . Hence, a diagram $(*)$

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \longrightarrow & Y \end{array}$$

is an object of \mathcal{C}/p . A diagram as in the middle is a morphism from

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \longrightarrow & Y \end{array} \rightarrow \begin{array}{ccc} \Omega(Y) & \longrightarrow & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \longrightarrow & Y \end{array} \in \mathcal{C}/p$$

So the fiber of the left projection over the diagram $(*)$ is the space of maps as the diagram above. But these are contractible, because the latter is a final object in \mathcal{C}/p .

To see the right arrow:

$$\left\{ \begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & \searrow & \downarrow \\ 0 & \xrightarrow{\quad} & Y \end{array} \right\} \rightarrow \{\text{diagrams } X \rightarrow \Omega(Y)\}$$

Given $X \xrightarrow{f} \Omega(Y)$, the fiber of this over f is roughly the extra data needed to factor

$$X \xrightarrow{f} \Omega(Y) \rightarrow Y$$

through 0. Because 0 is a zero object, each of the other simplices that make up this data are unique up to contractible choices. \square

28. 11/03/2021

We proceed the discussion of Adjunction of Loop and Suspension functor.

Theorem 28.1. *Let \mathcal{C} be a pointed ∞ -category with finite limits and colimits, then $\Sigma_{\mathcal{C}}, \Omega_{\mathcal{C}}$ are adjoint pair of functors*

Remark 28.2. This immediately suggests a distinguish class of ∞ -categories

Definition 28.3. ∞ -categories is stable if it is pointed, has finite limits and colimits, and Σ_C, Ω_C are autoequivalence.

- General Principle: the more symmetries a mathematical object has, the easier it is to study

- So the notion of stable ∞ -category just axiomatizes the class of ∞ -category that have maximal intrinsic symmetry.

Exercise 28.4. If \mathcal{C} is an pointed ordinary, then $\Sigma_{\mathcal{C}}(X) \cong \Omega_{\mathcal{C}}(X) \cong 0$ for all $Ob(\mathcal{C})$

- In particular, an ordinary category can never be stable.

- This challenges the default intuition that ordinary categories are the simplest ∞ -category they involve less complicated data, but they have less potential for symmetry.

- we can give another characterization of stable ∞ -category using the following notion:

Definition 28.5. Let $f : X \rightarrow Y$ be a morphism in a pointed ∞ -category. Then the fiber and cofiber of f are defined as:

$$fib(f) := \lim \left\{ \begin{array}{ccc} & K & \\ & \downarrow f & \\ 0 & \longrightarrow & Y \end{array} \right\} \text{ and } cof(f) := \lim \left\{ \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow f & & \\ Y & & \end{array} \right\}, \text{ when they exist.}$$

Example 28.6. Suppose \mathcal{C} is an abelian category. Then the fiber and cofiber of a morphism $f : X \rightarrow Y$ in \mathcal{C} are just its kernel and in the usual sense.

Example 28.7. Let $f : X \rightarrow Y$ be a morphism in Top_{∞}^* . Let's compute $cof(f)$.

Let p be the diagram $\left\{ \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow f & & \\ Y & & \end{array} \right\}$, then by definition $cof(f)$ is the initial object of

$$(Top_{\infty}^*)_{p/}. \text{ Then } Ob((Top_{\infty}^*)_{p/}) = \left\{ \text{diagrams } \begin{array}{ccccc} X & \longrightarrow & 0 & & \\ & \searrow h_1 & \downarrow & & \\ \downarrow f & & & \searrow g & \\ Y & \longrightarrow & Z & & \end{array} \right\}, \text{ Combine } h_1 \text{ and } h_2,$$

$$= \left\{ (Z, g, h) \text{ where } g : Y \rightarrow Z \text{ is a pointed map and } h : X \times [0, 1] \rightarrow Z \text{ such that } h|_{X \times \{0\}} = g \cdot f \text{ and } h|_{X \times \{1\} \cup X \times [0, 1]} = z_0 \right\}$$

- As with the reduced suspension we can construct an initial packaging of such data by forming a quotient space:

$$cof(f) = (X \times [0, 1]) \sqcup Y / \sim$$

where \sim denotes $\left\{ \begin{array}{l} (x, 0) \sim f(x) \text{ for } \forall x \in X \\ (x, 1) \sim (x', 1) \text{ for } \forall x, x' \in X \\ (x_0, a) \sim (x_0, b) \text{ for } \forall a, b \in [0, 1] \end{array} \right.$, this is called reduced mapping cone of f

Example 28.8. Consider $f : S^1 \rightarrow \mathbb{R}^2$, the inclusion of circle, we can then take union of a cylinder $S^1 \times [0, 1]$ with point in $S^1 \times \{0\}$ (which spans to a line) and \mathbb{R}^2 with a point, and glue the points. With quotient relation we obtain a geometric cone, which is the cofiber of f .

29. 11/5/2021

Last Time: Stable ∞ -categories, fibers, and cofibers

Example 29.1. Let $F : X_\bullet \rightarrow Y_\bullet$ be a morphism in Top_∞^* , and let p be the diagram used to define $\text{fib}(f)$. That is, $p : \begin{array}{c} \cdot \\ \downarrow \\ \cdot \end{array} \rightarrow \text{Top}_\infty^*$. Then

$$\begin{aligned}
 \text{Ob}((\text{Top}_\infty^*)/p) &= \left\{ \text{diagrams } \begin{array}{ccc} Z & \xrightarrow{g} & X \\ \downarrow & \searrow h_1 & \downarrow f \\ 0 & \xrightarrow{y_0} & Y \end{array} \right\} \\
 (29.2) \quad &= \left\{ (Z, g, h) \text{ where } g : Z \rightarrow X \text{ is a pointed map and} \right. \\
 &\quad \left. h : Z \times [0, 1] \rightarrow Y \text{ is such that} \right. \\
 &\quad \left. h|_{Z \times \{0\}} \equiv y_0 \text{ and } h|_{Z \times \{1\}} = f \circ g \right\} \\
 &= \left\{ (Z, g, \bar{h}), \text{ where } \bar{h} : Z \rightarrow \text{Maps}([0, 1], Y) \right. \\
 &\quad \left. \text{takes } z \in Z \text{ to } h(Z, -) : [0, 1] \rightarrow Y \right\},
 \end{aligned}$$

where h is obtained from combining h_1 and h_2 . Thus, the universal data of this kind is the subspace of $X \times \text{Maps}([0, 1], Y)$ satisfying the needed conditions:

$$\text{fib}(f) = \left\{ (x, \gamma) \in X \times \text{Map}([0, 1], Y) \text{ s.t. } \gamma(0) = y_0 \text{ and } \gamma(1) = f(x) \right\}.$$

Note that this recovers $\Omega(Y_\bullet)$ when $X = pt$.

Aside: When there is potential for ambiguity we call $\text{fib}(f)$ the **homotopy** fiber of f . The **set-theoretic** fiber of f is just $f^{-1}(y_0) \subset X_\bullet$. When the set-theoretic fibers of f don't change too wildly (technically: when f is a Serre fibration) the set-theoretic fiber is weakly homotopy equivalent to the homotopy fiber.

We can also characterize stable ∞ -categories in terms of how fibers and cofibers interact.

Definition 29.3. Let \mathcal{C} be a pointed ∞ -category. A **triangle** in \mathcal{C} is a diagram $\Lambda^1 \times \Lambda^1 \rightarrow \mathcal{C}$ of the form

$$\begin{array}{ccc} Z & \xrightarrow{g} & X \\ \downarrow \phi & \searrow h_1 & \downarrow f \\ 0 & \xrightarrow{y_0} & Y \end{array}$$

h_2 (on the diagonal from Z to Y)

That is, it is the data of a pair of morphisms f and g , a 2-simplex h_1 , identifying ϕ as a composition of f and g , and a 2-simplex h_2 factoring ϕ through 0 (a **null homotopy** of ϕ). We'll often write this data as $X \xrightarrow{f} Y \xrightarrow{g} Z$ leaving (ϕ, h_1, h_2) implicit.

Example 29.4. If \mathcal{C} is a pointed ordinary category, then (ϕ, h_1, h_2) are determined by f and g , and f and g define a triangle iff $g \circ f = 0$.

Note: Consider the composition $\mathrm{Map}_{\mathcal{C}}(X, 0) \times \mathrm{Map}_{\mathcal{C}}(0, Z) \rightarrow \mathrm{Map}_{\mathcal{C}}(X, Z)$. The domain is contractible, so we can talk about “the zero morphism from X to Z ”, understanding that this is well-defined up to a contractible space of choices.

Definition 29.5. We say a triangle is a **fiber sequence** if it is a pullback diagram (so $X = \mathrm{fib}(g)$) and a **cofiber sequence** if it is a pushout diagram (so $Z = \mathrm{cof}(f)$).

Example 29.6. If \mathcal{C} is a pointed ordinary category then a triangle is a fiber sequence (resp. cofiber) iff $0 \rightarrow X \xrightarrow{f} Y \xrightarrow{g} Z \rightarrow 0$ is **left exact** (resp. right exact).

Theorem 29.7. An ∞ -category is stable iff

- i) it is pointed
- ii) every morphism admits a fiber and cofiber
- iii) every fiber sequence is a cofiber sequence, and every cofiber sequence is a fiber sequence.

30. 11/12/2021

Remark 30.1. If \mathcal{C} is pointed, the space $\mathrm{Map}_{\mathcal{C}}(Y, X)$ is naturally a pointed space, and the same is true for the corresponding functor $\mathcal{C} \rightarrow \mathrm{Top}_{\infty}^*$.

Theorem 30.2. If \mathcal{C} is a stable ∞ -category, then $\mathrm{Ho}(\mathcal{C})$ is additive.

Proof. First note that for any ∞ -category \mathcal{D} , if X is an initial object of \mathcal{D} , then its image under the natural functor $\mathcal{D} \rightarrow \mathrm{Ho}(\mathcal{D})$ is initial in $\mathrm{Ho}(\mathcal{D})$. Similarly for final objects.

(The key point is, X is initial in $\mathrm{Ho}(\mathcal{D}) \Leftrightarrow \mathrm{Hom}_{\mathrm{Ho}(\mathcal{C})}(X, Y) \simeq \pi_0 \mathrm{Map}_{\mathcal{D}}(X, Y)$ has a unique element $\forall Y$, but by definition, X is initial in $\mathcal{D} \Leftrightarrow \mathrm{Map}_{\mathcal{D}}(X, Y)$ is contractible for all Y .)

In particular, $\mathrm{Ho}(\mathcal{C})$ has a zero object since \mathcal{C} does. Similarly, $\mathcal{C} \rightarrow \mathrm{Ho}(\mathcal{C})$ preserves products and coproducts. The key point here is that if $p: \mathcal{S} \rightarrow \mathcal{C}$ is a diagram indexed by a set \mathcal{S} (more

formally, \mathcal{S} is a simplicial set $\prod_s \Delta^0 \rightarrow \mathcal{C}$, then $\text{Ob}(\mathcal{C}_{/p}) = \{(X, \{f_s: X \rightarrow p(s)\}_{s \in \mathcal{S}})\}$ only involves objects and morphisms (rather than any high-dimensional simplices). From this, one can show that $\text{Ho}(\mathcal{C}_{/p}) \simeq \text{Ho}(\mathcal{C})_{/p}$. In particular, $\mathcal{C}_{/p} \rightarrow \text{Ho}(\mathcal{C}_{/p}) \simeq \text{Ho}(\mathcal{C})_{/p}$ preserves final objects by the above argument, so $\mathcal{C} \rightarrow \text{Ho}(\mathcal{C})$ preserves products and similarly coproducts.

But then $\text{Ho}(\mathcal{C})$ has finite products and coproducts since \mathcal{C} does by definition.

Now we verify that the mapping space is an abelian group. For any $X, Y \in \text{Ob}(\mathcal{C})$, we have $\text{Map}_{\mathcal{C}}(X, Y) \simeq \text{Map}_{\mathcal{C}}(\Omega^2 X, \Omega^2 Y) \simeq \Omega \text{Map}_{\mathcal{C}}(\Omega^2 X, \Omega Y) \simeq \Omega^2 \text{Map}_{\mathcal{C}}(\Omega^2 X, Y)$, the last two identifications coming from the fact that the functor $\text{Map}_{\mathcal{C}}(\Omega X, -)$ preserves limits. Hence we also have $\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y) = \pi_0 \text{Map}_{\mathcal{C}}(X, Y) = \pi_0 \Omega^2 \text{Map}_{\mathcal{C}}(\Omega^2 X, Y) = \pi_2 \text{Map}_{\mathcal{C}}(\Omega^2 X, Y)$ = the set of homotopy classes of maps $[0, 1]^2 \rightarrow \text{Map}_{\mathcal{C}}(\Omega^2 X, Y)$ taking $\partial[0, 1]^2$ to the base point.

Recall that for any pointed space Z_{\bullet} , $\pi_2 Z_{\bullet}$ is the set of homotopy classes of maps $[0, 1]^2 \rightarrow Z$ that takes $\partial[0, 1]^2$ to the base-point z_0 . (A reference of this is chapter 4 of [Hat01]) But π_2 is automatically an abelian group. In particular, $\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y)$ is an abelian group. \square

30.1. The Second Homotopy Group. Let (X, x_0) be a pointed topological space, then the **second homotopy group** $\pi_2(X, x_0)$ is defined to be the set

$$\pi_2(X, x_0) = \{f: [0, 1]^2 \rightarrow X \mid f(\partial[0, 1]^2) = x_0\} / f \sim g \Leftrightarrow f, g \text{ are homotopic rel } \partial[0, 1]^2.$$

We define the multiplication of f and g in $\pi_2(X, x_0)$ to be the map

$$f \vee g(s, t) := \begin{cases} f(2s, t), & \text{when } 0 \leq s \leq \frac{1}{2}; \\ g(2s - 1, t), & \text{when } \frac{1}{2} \leq s \leq 1. \end{cases}$$

The following Proposition is a special case of the one in [Hat01]:

Proposition 30.3. $\pi_2(X, x_0)$ with the product structure given above is an abelian group.

Proof. We can construct a homotopy between $f \vee g$ and $g \vee f$ as follows: For $0 \leq \ell \leq \frac{1}{3}$, define $h: [0, 1]^2 \times [0, \frac{1}{3}] \rightarrow X$ to be the map

$$h(s, t, \ell) = \begin{cases} f(2s, (1 + 3\ell)t), & \text{if } 0 \leq s \leq \frac{1}{2} \text{ and } t \leq \frac{1}{1+3\ell}; \\ g(2s - 1, (1 + 3\ell)t - 3\ell), & \text{if } \frac{1}{2} \leq s \leq 1 \text{ and } \frac{3\ell}{1+3\ell} \leq t \leq 1; \\ x_0, & \text{otherwise.} \end{cases}$$

and when $\frac{1}{3} \leq \ell \leq \frac{2}{3}$, we let $h: [0, 1]^2 \times [\frac{1}{3}, \frac{2}{3}] \rightarrow X$ to be


$$h(s, t, \ell) = \begin{cases} f(2s - 3\ell + 1, 2t), & \text{when } \frac{3}{2}\ell - \frac{1}{2} \leq s \leq \frac{3}{2}\ell \text{ and } 0 \leq t \leq \frac{1}{2}; \\ g(2s + 3\ell - 2, 2t - 1), & \text{when } 1 - \frac{3}{2}\ell \leq s \leq \frac{3}{2} - \frac{3}{2}\ell \text{ and } \frac{1}{2} \leq t \leq 1; \\ x_0, & \text{otherwise.} \end{cases}$$

When $\frac{2}{3} \leq \ell \leq 1$, we define $h: [0, 1]^2 \times [\frac{2}{3}, 1] \rightarrow X$ to be

$$h(s, t, \ell) = \begin{cases} f(2s - 1, \frac{2t}{3\ell - 1}), & \text{if } \frac{1}{2} \leq s \leq 1 \text{ and } 0 \leq t \leq \frac{3}{2}\ell - 1; \\ g(2s, \frac{2t + 3\ell - 3}{3\ell - 1}), & \text{if } 0 \leq s \leq 1 \text{ and } \frac{3}{2} - \frac{3\ell}{2} \leq t \leq 1; \\ x_0, & \text{otherwise.} \end{cases}$$

These three h s are all continuous maps and agree along their intersections, hence they glue to a single $h: [0, 1]^3 \rightarrow X$ which gives a homotopy from $f \vee g$ to $g \vee f$. \square

The following picture gives an intuition of the rigorous construction in the proof:



31. 11/15/2021

Last time: \mathcal{C} stable $\Rightarrow Ho(\mathcal{C})$ additive

Given $X, Y \in Ob(\mathcal{C})$ we can see the group structure on $Hom_{Ho(\mathcal{C})}(X, Y)$ more explicitly by recalling that

$$Map_{\mathcal{C}}(X, Y) \simeq Map_{\mathcal{C}}(X, \Omega \Sigma Y) \simeq \left\{ \begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & \searrow h_2 & \downarrow \\ 0 & \xrightarrow{h_1} & \Sigma Y \end{array} \right\}$$

where \simeq represents homotopy equivalence.

We can paste two such diagrams together

$$\left(\begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & \searrow h_2 & \downarrow \\ 0 & \xrightarrow{h_1} & \Sigma Y \end{array} , \begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & \searrow h'_2 & \downarrow \\ 0 & \xrightarrow{h'_1} & \Sigma Y \end{array} \right) \mapsto \begin{array}{ccc} X & \xrightarrow{\quad} & 0 \\ \downarrow & \searrow h_2 & \downarrow \\ 0 & \xrightarrow{h_1} & \Sigma Y \end{array}$$

(The resulting diagram also includes curved arrows labeled h'_1 and h'_2 from X to ΣY and 0 respectively.)

(really we should be careful about the fact that the morphism $X \rightarrow 0$ is only unique in a homotopical sense).

One can "replace" the upper right part with a single 2-simplex h_2'' to get a diagram of the form

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow h_2'' & \downarrow \\ 0 & \xrightarrow{h_1} & \Sigma Y \end{array}$$

where h_2' and h_2'' have identical top and right edges (i.e. define h_2'' using horn-filling properties of ∞ -categories and properties of zero objects).

This replacement is not unique, but different replacements are connected by paths in $\text{Map}_{\mathcal{C}}(X, Y)$. Hence the resulting binary operation on $\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y)$ is well-defined.

Another important feature of stable ∞ -categories is that we can "rotate triangles".

Proposition 31.1. *Let \mathcal{C} be a stable ∞ -category and*

$$X \xrightarrow{f} Y \xrightarrow{g} Z$$

a fiber sequence in \mathcal{C} . Then $\text{cof}(g) \simeq \Sigma X \simeq \Sigma(\text{fib}(g))$. In other words, there exists a fiber sequence of the form

$$Y \xrightarrow{g} Z \xrightarrow{h} \Sigma X$$

(these two sequences differ by a "rotation"). Thus in a stable ∞ -category fibers and cofibers are closely related (contrast this with how things work in an abelian category).

Corollary 31.2. *There exists a fiber sequence of the form*

$$Z \xrightarrow{h} \Sigma X \xrightarrow{*} \Sigma Y$$

(in fact this morphism $()$ is $-\Sigma f$, where "minus" refers to the additive group structure).*

Remark 31.3. Our discussion of the group structure on $\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y)$ via "diagram pasting" can be extended to prove that reversed diagrams

$$\begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow h_2 & \downarrow \\ 0 & \xrightarrow{h_1} & \Sigma Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X & \longrightarrow & 0 \\ \downarrow & \searrow h_1 & \downarrow \\ 0 & \xrightarrow{h_2} & \Sigma Y \end{array}$$

correspond to inverse elements of $\text{Hom}_{\text{Ho}(\mathcal{C})}(X, Y)$ with respect to the abelian group structure.

32. 11/19/2021

Last time we talked about rotation of triangles and generalized the notion of **exact sequence** from **Abelian category** to fiber sequence in stable ∞ -category. From the rotation we have that cofiber and fiber are quite the same object, which is different from **Abelian category**.

Definition 32.1. Let \mathcal{C} and \mathcal{D} be ∞ -categories with finite limits and colimits. Then $F : \mathcal{C} \rightarrow \mathcal{D}$ is left (resp. right) exact if it preserves finite limits (resp. colimits).

Theorem 32.2. *If \mathcal{C} and \mathcal{D} are stable ∞ -categories, then $F : \mathcal{C} \rightarrow \mathcal{D}$ is left exact if and only if it is right exact.*

Proof. Recall from last time that F is left exact $\iff F$ preserves direct sums and fibers and that F is right exact $\iff F$ preserves direct sums and cofibers. Hence we just need to show that if F is left exact, then it preserves cofibers, and if F is right exact, then it preserves fibers.

Suppose that F is left exact, and let $g : X \rightarrow Y$ be a morphism in \mathcal{C} . Note that F preserves Ω since it is left exact, that is, $F(\Omega Z) = \Omega F(Z)$. Hence, it also preserves $[n]$ for any n . Then

$$\begin{aligned} F(\text{cof}(g)) &\simeq F(\Sigma \text{fib}(g)) \\ &\simeq \Sigma F(\text{fib}(g)) \\ &\simeq \Sigma \text{fib}(F(g)) \\ &\simeq \text{cof}(F(g)) \end{aligned}$$

By a dual argument, F preserves fibers if it is right exact. □

Summary of last few lectures:

(1) Stable ∞ -categories have a number of distinguished features:

- (a) they have an intrinsic \mathbb{Z} -symmetry $X \rightarrow X[n]$
- (b) they can be characterized in a similar (but simpler way) as Abelian categories
- (c) they have additive homotopy categories
- (d) right exact functors between them are left exact, and vice versa.

A natural question arises: are there actual examples? The answer is Yes. This is because there are universal constructions for producing stable ∞ -categories from non-stable ∞ -categories.

(2) We will study a few such constructions, starting with the **Spanier-Whitehead category** $SW(\mathcal{C})$ of a pointed ∞ -category \mathcal{C} with finite colimits (a.k.a, the category of Σ -spectrum objects of \mathcal{C}).

Analogy: Consider $\mathbb{Z}[x]$ together with the multiplication operator $x : \mathbb{Z}[x] \rightarrow \mathbb{Z}[x]$ that takes $f(x)$ to $xf(x)$, which is a homomorphism of $\mathbb{Z}[x]$ -modules as Abelian groups. This operator is not invertible, but there's a universal construction which makes it invertible: the

localization $\mathbb{Z}[x, x^{-1}]$. If we analogize $(\mathcal{C}, \Sigma_{\mathcal{C}})$ with $(\mathbb{Z}[x], x)$, is there a corresponding analogue of $\mathbb{Z}[x, x^{-1}]$?

Note that we have a filtration

$$\mathbb{Z}[x] \subset x^{-1}\mathbb{Z}[x] \subset x^{-2}\mathbb{Z}[x] \subset \cdots \subset \mathbb{Z}[x, x^{-1}].$$

In other words, $\mathbb{Z}[x, x^{-1}] \simeq \operatorname{colim}_{n \in \mathbb{N}} x^{-n}\mathbb{Z}[x]$ as $\mathbb{Z}[x]$ -modules as Abelian groups. Note that as $\mathbb{Z}[x]$ -modules, each $x^{-n}\mathbb{Z}[x]$ is isomorphic to $\mathbb{Z}[x]$ itself.

33. 11/22/2021

Last time: exactness

Given the definition of a stable ∞ -category, there exists a natural question that we will explore today. Given a non-stable ∞ -category \mathcal{C} , can we "approximate" it with a stable ∞ -category \mathcal{C}' ? The answer is yes, and one method of doing this is by inverting the suspension function Σ .

We begin by an analogy from basic algebra. Given a $\mathbb{Z}[x]$ -module M , the localization $x^{-1}M$ is a $\mathbb{Z}[x]$ -module in which x acts invertibly, and is universal with respect to homomorphisms from M to modules with this property. This means that if x acts invertibly on any $\mathbb{Z}[x]$ -module N , then any homomorphism $\varphi : M \rightarrow N$ factors uniquely through the localization $\varphi_M : M \rightarrow x^{-1}M$. Equivalently, composition with φ_M induces a bijection $\operatorname{Hom}_{\mathbb{Z}[x]\text{-mod}}(x^{-1}M, N) \cong \operatorname{Hom}_{\mathbb{Z}[x]\text{-mod}}(M, N)$.

For the case $M = \mathbb{Z}[x]$ we have that $x^{-1}\mathbb{Z}[x]$ is $\mathbb{Z}[x^{\pm 1}]$, the Laurent polynomials. In general, we also have

$$x^{-1}M \cong M \otimes_{\mathbb{Z}[x]} \mathbb{Z}[x^{\pm 1}].$$

Now note that the filtration by degree presents the Laurent polynomials as a colimit of free modules as

$$\mathbb{Z}[x] \cong \operatorname{colim}_{\mathbb{N}} (\mathbb{Z}[x] \subset x^{-1}\mathbb{Z}[x] \subset x^{-2}\mathbb{Z}[x] \subset \cdots).$$

We can use the fact that $x^{-n}\mathbb{Z}[x] \cong \mathbb{Z}[x]$ as a $\mathbb{Z}[x]$ module to rewrite the above colimit. Under this isomorphism, the morphisms in the filtered system above are just multiplication by x which gives us

$$\mathbb{Z}[x] \cong \operatorname{colim}_{\mathbb{N}} (\mathbb{Z}[x] \xrightarrow{x} \mathbb{Z}[x] \xrightarrow{x} \mathbb{Z}[x] \xrightarrow{x} \cdots).$$

We will make a similar construction for an ∞ -category \mathcal{C} for which $\Sigma_{\mathcal{C}}$ is well defined, replacing multiplication by x with the suspension $\Sigma_{\mathcal{C}}$.

Definition 33.1. Let \mathcal{C} be a pointed ∞ -category with finite colimits. Then the *Spanier-Whitehead category* $\operatorname{SW}(\mathcal{C})$ of \mathcal{C} is the colimit

$$\operatorname{SW}(\mathcal{C}) = \operatorname{colim}_{\mathbb{N}} (\mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \cdots)$$

in $\operatorname{Cat}_{\infty}$.

We can now compare $SW(\mathcal{C})$ to $\mathbb{Z}[x]$. $SW(\mathcal{C})$ is the target of a collection of functions

$$\{i_n : \mathcal{C} \rightarrow SW(\mathcal{C})\}$$

just like $\mathbb{Z}[x^{\pm 1}]$ is the target of a collection of $\mathbb{Z}[x]$ -module homomorphisms

$$\{\iota_n : \mathbb{Z}[x] \xrightarrow{x^{-n}} \mathbb{Z}[x^{\pm 1}]\}.$$

Every object of $SW(\mathcal{C})$ is of the form $i_n(x)$ for some $x \in Ob\mathcal{C}$ and $n \in \mathbb{N}$ just like any element of $\mathbb{Z}[x^{\pm 1}]$ is of the form $f(x) \in \mathbb{Z}[x]$ and $n \in \mathbb{N}$.

34. 11/29/2021

Last time: the Spanier-Whitehead category.

Definition 34.1. Let \mathcal{C} be a pointed ∞ -category with finite colimits. Then

$$SW(\mathcal{C}) := \operatorname{colim}_{\mathbb{N}} (\mathcal{C} \xrightarrow{\Sigma} \mathcal{C} \xrightarrow{\Sigma} \mathcal{C} \cdots)$$

where the colimit is taken in Cat_{∞} .

- Explicitly, we have a functor $i_n : \mathcal{C} \rightarrow SW(\mathcal{C})$ for every $n \in \mathbb{N}$, and every object of $SW(\mathcal{C})$ is in the essential image of i_n for all $n \gg 0$.

- This corresponds to the fact that every Laurent polynomial is in the image of

$$\mathbb{Z}[x] \rightarrow \mathbb{Z}[x, x^{-1}]$$

$$f(x) \mapsto x^{-n}f(x)$$

for $n \gg 0$.

- Given $m, n \in \mathbb{N}$ and $X, Y \in Ob(\mathcal{C})$, we moreover have

$$Map_{SW(\mathcal{C})}(i_n(X), i_n(Y)) \simeq \operatorname{colim}_{k \geq m, m} Map_{\mathcal{C}}(\Sigma^{k-m}X, \Sigma^{k-n}Y)$$

where the colimit is taken in Top_{∞}^* .

- We also have that

$$Hom_{Ho(SW(\mathcal{C}))}(i_n(X), i_n(Y)) \simeq \operatorname{colim}_{k \geq m, m} Hom_{Ho(\mathcal{C})}(\Sigma^{k-m}X, \Sigma^{k-n}Y)$$

where the colimit is taken in Set . (Rightarrow we can describe $Ho(SW(\mathcal{C}))$ purely in terms of $Ho(\mathcal{C})$)

Theorem 34.2. (1) $SW(\mathcal{C})$ is stable.

(2) Let \mathcal{D} be any stable ∞ -category. Then composition with $i_0 : \mathcal{C} \rightarrow SW(\mathcal{C})$ induces an equivalence

$$Fun^{rex}(SW(\mathcal{C}), \mathcal{D}) \simeq Fun^{rex}(\mathcal{C}, \mathcal{D})$$

between the categories of right exact functors from \mathcal{C} and $SW(\mathcal{C})$ to \mathcal{D} .

- Recall that the universal property of $x^{-1}M$ was characterized as follows: if N is any $\mathbb{Z}[x]$ -module on which x acts invertibly, composition with $M \xrightarrow{\varphi_x} x^{-1}M$ induces a bijection

$$\mathrm{Hom}_{\mathbb{Z}[x]\text{-mod}}(x^{-1}M, N) \simeq \mathrm{Hom}_{\mathbb{Z}[x]\text{-mod}}(M, N)$$

between the sets of $\mathbb{Z}[x]$ -module homomorphisms. So (2) says that swc is characterized by a universal property just like $x^{-1}M$ is.

Proof. (1) We can describe $\Sigma_{SW(\mathcal{C})}$ and its inverse via maps of directed systems: By construction the endomorphism of swc induced by the red arrow is inverse to the one induced by the blue arrow, which is just $\Sigma_{SW(\mathcal{C})}$.

(2) Note that $\mathcal{D} \simeq SW(\mathcal{D})$ since \mathcal{D} is stable. Then the inverse of composition with i_0 take a right-exact functor $F : \mathcal{C} \rightarrow \mathcal{D}$ to the functor $\tilde{F} : SW(\mathcal{C}) \rightarrow \mathcal{D}$ defined by (Note that since swc and sD are stable, the extension $\tilde{F} : SW(\mathcal{C}) \rightarrow \mathcal{D}$ is automatically left exact even if F is not.)

□

Bad example: if \mathcal{C} is an ordinary abelian category, then $\Sigma_{\mathcal{C}} : \mathcal{C} \rightarrow \mathcal{C}$ is the zero functor, i.e. $\Sigma_{\mathcal{C}}(X) \simeq 0$ for all $X \in \mathrm{Ob}(\mathcal{C})$, hence $SW(\mathcal{C})$ is just the zero category, i.e. it is contractible.

35. 12/1/2021

Last time : We defined the Spanier-Whitehead category $SW(\mathcal{C})$ of a pointed ∞ -category \mathcal{C} .

Definition 35.1. Given a pointed ∞ -category \mathcal{C} with finite colimits, we define

$$SW(\mathcal{C}) := \mathrm{colim}_{\mathbb{N}} (\mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Sigma_{\mathcal{C}}} \dots)$$

in \mathbf{Cat}_{∞} .

Theorem 35.2. (1) $SW(\mathcal{C})$ is stable.

(2) (Universal property of $SW(\mathcal{C})$) Let \mathcal{D} be any stable ∞ -category. Then composition with $i_0 : \mathcal{C} \rightarrow SW(\mathcal{C})$ induces an equivalence

$$\mathrm{Fun}^{rex}(SW(\mathcal{C}), \mathcal{D}) \xrightarrow{\sim} \mathrm{Fun}^{rex}(\mathcal{C}, \mathcal{D})$$

Example 35.3. $\mathbf{Sp}^{fin} := SW(\mathbf{Top}_{\infty}^{*,fin})$, the homotopy category of finite spectra or the finite stable homotopy category.

We say a CW complex is finite if it can be built out of finitely many cells (e.g. S^n for any $n \in \mathbb{N}$, any algebraic subset of \mathbb{R}^n etc.). Now let $\mathbf{Top}_{\infty}^{*,fin} \subset \mathbf{Top}_{\infty}^*$ denote the full sub ∞ -category of finite CW complexes. Note if X is a finite pointed CW complex then so is ΣX (In fact any finite colimit of finite CW complexes is again finite.).

Identifying $X, Y \in \mathbf{Top}_{\infty}^{*,fin}$ with their image under $i_0 : \mathbf{Top}_{\infty}^{*,fin} \rightarrow \mathbf{Sp}^{fin}$. Let's compare

$$\mathrm{Hom}_{\mathrm{Ho}(\mathbf{Top}_{\infty}^{*,fin})}(X, Y) \quad \text{and} \quad \mathrm{Hom}_{\mathbf{Sp}^{fin}}(X, Y)$$

the former is the set of homotopy classes of continuous maps from X to Y and is written as $[X, Y]$.

As we have seen before, the latter can be written as $\operatorname{colim}_{\mathbb{N}}[\Sigma^n X, \Sigma^n Y]$. This colimit is often written $[X, Y]_s$ and is called stable homotopy classes of maps $X \rightarrow Y$.

Theorem 35.4. (*Freudenthal Suspension Theorem*) *If X and Y are finite CW complexes then for sufficiently large n ,*

$$[\Sigma^n X, \Sigma^n Y] \rightarrow [\Sigma^{n+1} X, \Sigma^{n+1} Y]$$

is a bijection. (So homotopy classes of maps "stabilize" under suspension)

Classical Viewpoint : "Stable homotopy theory" (e.g. the study of $[X, Y]_s$) is a more accessible approximation of ordinary homotopy theory. For example, $[X, Y]_s$ has an abelian group structure.

Definition 35.5. Let \mathcal{C} be a pointed ∞ -category with finite limits. The category of spectrum objects (or Ω -spectrum) is the limit

$$\mathbf{Sp}(\mathcal{C}) := \lim_{\mathbb{N}} (\cdots \xrightarrow{\Omega_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Omega_{\mathcal{C}}} \mathcal{C} \xrightarrow{\Omega_{\mathcal{C}}} \mathcal{C})$$

in \mathbf{Cat}_{∞} .

We denote the functor which projects onto the n^{th} term of the diagram as

$$\Omega_{\mathcal{C}}^{\infty-n} : \mathbf{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$$

Explicitly, an object X of $\mathbf{Sp}(\mathcal{C})$ is determined by the objects $\{X_n := \Omega_{\mathcal{C}}^{\infty-n}(X)\}_{n \in \mathbb{N}}$ and the induced isomorphism $\{X_n \simeq \Omega_{\mathcal{C}}(X_{n+1})\}_{n \in \mathbb{N}}$.

Analogy : $\mathbb{Z}[[X]] \simeq \lim_{\mathbb{N}} \mathbb{Z}[X]/(X^n)$ and the data of a formal power series is the data of a collection $\{f_n(X) \in \mathbb{Z}[X]/(X^n)\}$ such that $f_n(X) \equiv f_{n+1}(X) \pmod{X^n}$.

36. 12/3/2021

Last time : We discussed stable homotopy theory, Ω -spectra.

Definition 36.1. Let \mathcal{C} be a pointed ∞ -category with finite limits. The category $\mathbf{Sp}(\mathcal{C})$ of Ω -spectrum objects of \mathcal{C} is the limit

$$\mathbf{Sp}(\mathcal{C}) = \lim_{n \in \mathbb{N}} (\cdots \xrightarrow{\Omega} \mathcal{C} \xrightarrow{\Omega} \mathcal{C} \xrightarrow{\Omega} \mathcal{C})$$

in \mathbf{Cat}_{∞} .

Writing $\{\Omega^{\infty-n} : \mathbf{Sp}(\mathcal{C}) \rightarrow \mathcal{C}\}_{n \in \mathbb{N}}$ for the tautological functors the data of an object $X \in \mathbf{Sp}(\mathcal{C})$ is equivalent to the data of $\{X_n = \Omega^{\infty-n}(X)\}_{n \in \mathbb{N}}$ together with isomorphisms $\{X_n \simeq \Omega(X_{n+1})\}$

Key Example : We call $\mathbf{Sp} := \mathbf{Sp}(\mathbf{Top}_\infty^*)$ the category of spectra or the stable homotopy category. Note that given an object X of \mathbf{Sp} we have $X_0 \simeq \Omega^n(X_n)$ for every n . We say X_0 is an infinite loop space, which is a special property.

$\mathbf{Sp}(\mathcal{C})$ has a dual universal property to $\mathcal{SW}(\mathcal{C})$.

Theorem 36.2. (1) $\mathbf{Sp}(\mathcal{C})$ is stable.

(2) Let \mathcal{D} be any stable ∞ -category, then composition with $\Omega^\infty : \mathbf{Sp}(\mathcal{C}) \rightarrow \mathcal{C}$ induces an equivalence

$$\mathrm{Fun}^{lex}(\mathcal{D}, \mathbf{Sp}(\mathcal{C})) \xrightarrow{\sim} \mathrm{Fun}^{lex}(\mathcal{D}, \mathcal{C})$$

of categories of left exact functors.

If \mathcal{C} has both finite limits and colimits then we get a tautological functor

$$\mathcal{SW}(\mathcal{C}) \rightarrow \mathbf{Sp}(\mathcal{C})$$

which is fully faithful but not an equivalence.

Note that $\mathcal{SW}(\mathcal{C})$ or $\mathbf{Sp}(\mathcal{C})$ are trivial/degenerate when \mathcal{C} is an ordinary category.

Question : Is there a natural way of "approximating" an abelian category \mathcal{C} with a stable ∞ -category?

Answer : Yes, via parsing to the simplicial objects of \mathcal{C}

Recall that we described \mathbf{Top}_∞ via Kan complexes and simplicial sets -

$$\mathbf{Top}_\infty \simeq \mathcal{N}_\Delta(\mathrm{Kan}) \subset \mathcal{N}_\Delta(\mathrm{Set}_\Delta)$$

where \mathcal{N}_Δ denotes the simplicial nerve and the categories inside are thought of as simplicial categories in a natural way. Also recall that given any category \mathcal{C} we have the category $\mathcal{C}_\Delta := \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathcal{C})$ of simplicial objects of \mathcal{C} . Our idea would be to repeat the process

$$\mathrm{Set} \rightsquigarrow \mathrm{Kan} \subset \mathrm{Set}_\Delta \rightsquigarrow \mathbf{Sp}^{fin} \subset \mathbf{Sp}$$

with a different ordinary category \mathcal{C} in place of Set . This construction turns out to be particularly nice when \mathcal{C} is additive.

Proposition 36.3. Let \mathcal{C} be an additive category. Then \mathcal{C}_Δ can be canonically enhanced to a simplicial category whose mapping spaces are Kan complexes.

REFERENCES

- [Hat01] Allen Hatcher. *Algebraic Topology*. Cambridge University Press, Cambridge, England, UK, Dec 2001. [47](#)
- [Lur09] Jacob Lurie. *Higher Topos Theory (AM-170)*. Princeton University Press, Princeton, NJ, USA, Jul 2009. [40](#)