

STA 1013 : Statistics through Examples

Lecture 7: Measure of Center

Hwiyoung Lee

September 16, 2019

Department of Statistics, Florida State University

1. Measure of Center

1.1 Mean

1.2 Median

1.3 Mode

2. Weighted Mean

Measure of Center

Measure of Center

Measure of Center

- Mean
- Median
- Mode

Mean

The **mean** is what we most commonly call the average value

$$\text{Mean} = \frac{\text{sum of all values}}{\text{total number of values}} = \frac{\sum_{i=1}^n x_i}{n} = \bar{x}$$

- x_i : i th observation
- Example : 1, 2, 2, 4, 5, 10

$$\frac{1 + 2 + 2 + 4 + 5 + 10}{6} = 4$$

Mean

- Most familiar measure of center
- Takes every value into account
- Affected by outliers

Example : Annual Income

Name	Annual Income
Tom	\$ 32,000
Larry	\$ 36,000
Susan	\$ 39,000
Paul	\$ 41,000
Marcus	\$ 50,000
Randy	\$ 57,000
Sandy	\$ 60,000
Tim	\$ 75,000
Pam	\$ 80,000
Kim	\$ 95,000

Mean ?

Example : Annual Income

Bill Gates moves to town

Name	Annual Income
Tom	\$ 32,000
Larry	\$ 36,000
Susan	\$ 39,000
Paul	\$ 41,000
Marcus	\$ 50,000
Randy	\$ 57,000
Sandy	\$ 60,000
Tim	\$ 75,000
Pam	\$ 80,000
Kim	\$ 95,000
Bill Gates	\$ 5,000,000,000

Mean ?

Outlier

a value that is much higher or much lower than almost all other values.

- Bill Gates's annual income \$ 5,000,000,000 would be an outlier
- An outlier can pull the mean significantly upward (or downward)
- Making the mean unrepresentative of the data set as a whole

Example : Outlier

Example

Imagine that the 5 graduating seniors on a college basketball team receive the following first-year contract offers to play in the NBA (zero indicates that the player did not receive a contract offer):

0	0	0	0	\$ 10,000,000
---	---	---	---	---------------

Table 1: Graduating seniors NBA contract offer

- Is it therefore fair to say that the average senior on this basketball team received a \$2 million contract offer?
- Not really. The single player receiving the large offer makes the mean much larger than it would be otherwise.

Median : the middle value of the ordered data

- Half of the observations are larger and half are smaller than the Median.
- How to get ?
 1. Sort the data ascending order : the lowest value to the highest value.
 2. The Median is the $(n + 1)/2$ th observation
 3. If n is odd, the median is the middle observation of the ordered array.
 4. If n is even, it is midway between the two central observations.
- Median is robust to outliers

Example ($n = \text{odd}$)

Example : 2, 4, 1, 10, 8, 9, 5

- Sort : 1, 2, 4, 5, 8, 9, 10
- Median : $(7 + 1)/2$ th observation
 - 4 th observation : 5
- Q: What happens to the median if we change the 10 to 1,000?

- Mean :
- Q: What happens to the mean if we change the 10 to 1,000?

Example ($n = \text{even}$)

Example : 5, 11, 1, 13, 6, 9, 8, 3

- Sort : 1, 3, 5, 6, 8, 9, 11, 13
- Median : $(8 + 1)/2$ th observation
 - 4.5 th observation : average of 4 th and 5 th observation
- Q: What happens to the median if we change the 1 to -1,000?
- Mean :
- Q: What happens to the mean if we change the 1 to -1,000?

Example : Annual Income

Bill Gates moves to town

Name	Annual Income
Tom	\$ 32,000
Larry	\$ 36,000
Susan	\$ 39,000
Paul	\$ 41,000
Marcus	\$ 50,000
Randy	\$ 57,000
Sandy	\$ 60,000
Tim	\$ 75,000
Pam	\$ 80,000
Kim	\$ 95,000
Bill Gates	\$ 5,000,000,000

Median ?

Midrange

- The value that midway between the maximum and minimum in the data set

$$\text{Midrange} = \frac{\text{Max} + \text{Min}}{2}$$

- Easy to calculate
- Example : 1, 2, 4, 5, 7, 9, 10, 11
 - Midrange :
 - Median :

Midrange

- Because the midrange uses only the maximum and minimum values (ignore all other values), it is very **sensitive to those extremes**
- Example : 0, 5, 5, 5, 6, 7, 8, 9, 100
 - Midrange :
 - Median :
- Not popular (Rarely used)

Mode

Mode : the value of the data that occurs with the greatest frequency.

Example : 1, 2, 3, 3, 3, 4, 5

- Mean :
- Median :
- Mode :

Mode for skewed data

Example : 5, 5, 5, 6, 6, 7, 8, 9, 10, 11, 100 , 500, 1,000,000

- Mean :
- Median :
- Mode :

Mode for Right skewed data

Example : 5, 5, 5, 6, 6, 7, 8, 9, 10, 11, 100 , 500, 1,000,000

- Mean : 76,974.77
- Median : 8
- Mode : 5

Mode < Median < Mean

Mode for skewed data

Example : -1,000,000, -500, -100, 6, 6, 7, 7, 7, 8, 8, 8, 8, 10

- Mean :
- Median :
- Mode :

Mode for Left skewed data

Example : -1,000,000, -500, -100, 6, 6, 7, 7, 7, 8, 8, 8, 8, 10

- Mean :-7,732.692
- Median : 7
- Mode : 8

Mode > Median > Mean

Mean, Median always **exist** and are always **unique**

1. The mode **may not exist**

- Example : 1, 2, 3, 4, 5, 6, 7, 8, 9

2. The mode **may not be unique**

- Example : 1, 2, 2, 2, 5, 6, 7, 7, 7, 8, 8, 9, 10, 10, 10

Mode for continuous data

Example : Weight data

195.6	200.4	165.6	165.3	191.7	169.3	153.2
189.5	170.4	149.3	185.3	150.3	179.6	160.3
198.5	163.2	166.3	197.3	201.3	168.2	198.4

- Mean : 177.0952

149.3	150.3	153.2	160.3	163.2	165.3	165.6
166.3	168.2	169.3	170.4	179.6	185.3	189.5
191.7	195.6	197.3	198.4	198.5	200.4	201.3

Table 2: Sorted weight data

- Median : 170.4

For continuous data, Mode usually does not exist !!

Mode for binned data

Binned weight data

Weight	Count
140 ~ 149.9	1
150 ~ 159.9	2
160 ~ 169.9	7
170 ~ 179.9	2
180 ~ 189.9	2
190 ~ 199.9	5
200 ~ 209.9	2

- **Mode** : the bin with the highest frequency (160 ~ 169.9)
- Binned data will be used for the Histogram

Weighted Mean

Motivation

Suppose two different groups : A, B

Group	n	Mean
A	10	20
B	30	40

- The overall mean of 40 people

$$\frac{20 + 40}{2} = 30 ?$$

Totally wrong !!

The sizes of the two different groups must be taken into account

Motivation

Recall the mean formula :

$$\text{Mean} = \frac{\sum_{i=1}^n x_i}{n} = \frac{\text{Total sum of all data values}}{\text{The number of all data values}}$$

Since,

- Group A

$$\text{Total sum} : 10 \times 20 = 200 \left(\because \frac{\text{Total sum of A}}{10} = 20 \right)$$

- Group B

$$\text{Total sum} : 30 \times 40 = 1,200 \left(\because \frac{\text{Total sum of B}}{30} = 40 \right)$$

Thus,

$$\text{Mean} = \frac{10 \times 20 + 30 \times 40}{40} = \frac{200 + 1,200}{40} = 35$$

Motivation

State	n	Mean
Alabama	n_1	\bar{x}_1
Alaska	n_2	\bar{x}_2
\vdots	\vdots	\vdots
Wisconsin	n_{49}	\bar{x}_{49}
Wyoming	n_{50}	\bar{x}_{50}

Don't do

$$\bar{x}_{usa} = \frac{\bar{x}_1 + \bar{x}_2 + \cdots + \bar{x}_{49} + \bar{x}_{50}}{50} \quad \text{Wrong !!}$$

The sizes of the 50 different states must be taken into account

$$\bar{x}_{usa} = \frac{n_1\bar{x}_1 + n_2\bar{x}_2 + \cdots + n_{49}\bar{x}_{49} + n_{50}\bar{x}_{50}}{n_1 + n_2 + \cdots + n_{49} + n_{50}}$$

Weighted Mean

A weighted mean accounts for variations in the relative importance of data values.

$$\begin{aligned}\text{Weighted mean} &= \frac{\text{Sum of (each data value} \times \text{its weight)}}{\text{sum of all weights}} \\ &= \frac{\sum_{i=1}^n w_i x_i}{\sum_{i=1}^n w_i}\end{aligned}$$

- Each data value is assigned a weight (w_i)
- Weighted means are appropriate whenever the data values vary in their degree of importance

Example : Grade

Grade

Suppose your course grade is based on four quizzes and one final exam. Each quiz counts as 15% of your final grade, and the final counts as 40%. Your quiz scores are 75, 80, 84, and 88, and your final exam score is 96. What is your overall score?

solution :

$$\frac{15 \times 75 + 15 \times 80 + 15 \times 84 + 15 \times 88 + 40 \times 96}{15 + 15 + 15 + 15 + 40} = \frac{8745}{100} = 87.45$$

Exercise : Grade

Grade

Suppose your course grade is based on four quizzes and one final exam. Your lowest quiz grade will be dropped, and the other three will equally worth 20% of your grade, and the final counts as 40%. Your quiz scores are 100, 70, 64, and 88, and your final exam score is 96. What is your overall score?

solution :

Example : GPA

GPA

Randall has 38 credits with a grade of A, 22 credits with a grade of B, and 7 credits with a grade of C. What is his grade point average (GPA)? Base the GPA on values of 4.0 points for an A, 3.0 points for a B, and 2.0 points for a C.

solution :

$$\frac{38 \times 4 + 22 \times 3 + 7 \times 2}{38 + 22 + 7} = \frac{232}{67} = 3.46$$

Exercise : GPA

GPA

Lee has 30 credits with a grade of A, 15 credits with a grade of B, 5 credits with a grade of C, and 3 credits with F. What is his grade point average (GPA)? Base the GPA on values of 4.0 points for an A, 3.0 points for a B, 2.0 points for a C, and 0 points for a F.

solution :