
1 NMR Shielding

Shielding Tensor:

$$\sigma_{\beta\alpha}^K = \left. \frac{d^2 E}{dB_\alpha dm_{K\beta}} \right|_{\mathbf{B}, \mathbf{m}_K=0} \quad (1)$$

How do I parameterize energy E with \mathbf{B} and \mathbf{m}_K ?

The one-electronic Hamiltonian in magnetic field:

$$h(\mathbf{r}, \mathbf{B}, \mathbf{m}) = \frac{1}{2} \boldsymbol{\pi}^2 - \phi(\mathbf{r}) \quad (2)$$

in which:

$$\boldsymbol{\pi} = -i\nabla + \mathbf{A} \quad (3)$$

is the kinetic momentum operator.

Vector potential:

$$\mathbf{A}_i = \mathbf{A}_0(\mathbf{r}_i) + \sum_K \mathbf{A}_K(\mathbf{r}_i) \quad (4)$$

with:

$$\mathbf{A}_0(\mathbf{r}_i) = \frac{1}{2} \mathbf{B} \times \mathbf{r}_0 \quad \mathbf{B} = \nabla \times \mathbf{A}(\mathbf{r}) \quad (5)$$

$$\mathbf{A}_K(\mathbf{r}_i) = \alpha^2 \frac{\mathbf{M}_K \times \mathbf{r}_K}{r_K^3} \quad \mathbf{B}_K(\mathbf{r}) = \nabla \times \mathbf{A}_K(\mathbf{r}) \quad (6)$$

The first part is contribution from the external magnetic field, the second part from the nuclear magnetic moments.

Now parameterize with MO coefficients / densities?

2 SCF Level

$$\begin{aligned}
 E^{\text{SCF}} &= \sum_i^N h_{ii} + \frac{1}{2} \sum_{ij}^N \langle ij || ij \rangle \\
 &= \sum_i \sum_{\mu\nu} C_{\mu i}^* h_{\mu\nu} C_{\nu i} + \frac{1}{2} \sum_{ij} \langle ij || ij \rangle
 \end{aligned} \tag{7}$$

$$D_{\mu\nu}^{\text{SCF}} = \sum_i C_{\mu i}^* C_{\nu i} \tag{8}$$

$$C_{\mu p}(\lambda) = \sum_q C_{\mu q}(0) U_{qp}(\lambda) \tag{9}$$

At SCF level, the NMR shielding tensor is given as:

$$\sigma_{\beta\alpha}^{\text{SCF},K} = \left. \frac{d^2 E^{\text{SCF}}}{dB_\alpha dm_{K_\beta}} \right|_{\mathbf{B}, \mathbf{m}_K=0} \tag{10}$$

Taking the first derivative against the nuclear magnetic moment gives:

$$\begin{aligned}
 \frac{dE^{\text{SCF}}}{dm_{K_\beta}} &= \frac{d}{dm_{K_\beta}} \left(\sum_{i\mu\nu} C_{\mu i}^* h_{\mu\nu} C_{\nu i} \right) \\
 &= \sum_{i\mu\nu} C_{\mu i}^* C_{\nu i} \frac{dh_{\mu\nu}}{dm_{K_\beta}} \\
 &= \sum_{\mu\nu} D_{\mu\nu}^{\text{SCF}} \frac{dh_{\mu\nu}}{dm_{K_\beta}}
 \end{aligned} \tag{11}$$

Note that the MO coefficients are variationally determined so $\frac{d\mathbf{C}}{dm_{K_\beta}} = \mathbf{0}$, and the basis function does not depend on the nuclear magnetic moment, i.e. $\frac{d\phi_\mu}{dm_{K_\beta}} = 0$.

Now taking the second derivative w.r.t. the external magnetic field:

$$\begin{aligned}
 \sigma_{\beta\alpha}^{\text{SCF},K} &= \frac{d^2 E^{\text{SCF}}}{dB_\alpha dm_{K_\beta}} \\
 &= \frac{d}{dB_\alpha} \left(\sum_{\mu\nu} D_{\mu\nu}^{\text{SCF}} \frac{dh_{\mu\nu}}{dm_{K_\beta}} \right) \\
 &= \sum_{\mu\nu} D_{\mu\nu}^{\text{SCF}} \frac{d^2 h_{\mu\nu}}{dB_\alpha dm_{K_\beta}} + \sum_{\mu\nu} \frac{dD_{\mu\nu}^{\text{SCF}}}{dB_\alpha} \frac{dh_{\mu\nu}}{dm_{K_\beta}}
 \end{aligned} \tag{12}$$

The response of SCF density to the magnetic field perturbation is:

$$\begin{aligned}
 \frac{dD_{\mu\nu}^{\text{SCF}}}{dB_\alpha} &= \frac{d}{dB_\alpha} \left(\sum_i C_{\mu i}^*(\mathbf{B}) C_{\nu i}(\mathbf{B}) \right) \\
 &= \frac{d}{dB_\alpha} \left(\sum_{ipq} C_{\mu p}^*(0) U_{pi}^*(\mathbf{B}) C_{\nu q}(0) U_{qi}(\mathbf{B}) \right) \\
 &= \sum_{ip} C_{\mu p}^*(0) \frac{dU_{pi}^*(\mathbf{B})}{dB_\alpha} C_{\nu i}(\mathbf{B}) + \sum_{iq} C_{\mu i}^*(\mathbf{B}) \frac{dU_{qi}(\mathbf{B})}{dB_\alpha} C_{\nu q}(0)
 \end{aligned} \tag{13}$$

$$= \sum_{ip} C_{\mu p}^* (U_{pi}^{B_\alpha})^* C_{\nu i} + \sum_{ip} C_{\mu i}^* U_{pi}^{B_\alpha} C_{\nu p} \quad (14)$$

The virtual-occupied block of \mathbf{U}^B is obtained from the CPSCF equations, and the occupied-occupied block is chosen according to the orthonormality condition.