

Chap 2 Note: Random Variables

1 Random Variables and CDF

Example Suppose that independent trials, each of which results in any of m possible outcomes with respective probabilities p_1, p_2, \dots, p_m , $\sum_{i=1}^m p_i = 1$, are continually performed. Let X denote the number of trials needed until each outcome has occurred at least once.

Explain Rather than directly considering $P\{X = n\}$ we will first determine $P\{X > n\}$, the probability that at least one of the outcomes has not yet occurred after n trials. Letting A_i denote the event that outcome i has not yet occurred after the first n trials, $i = 1, \dots, m$, then

$$\begin{aligned} P\{X > n\} &= P(\cup_{i=1}^m A_i) \\ &= \sum_{i=1}^m P(A_i) - \sum_{i < j} P(A_i A_j) + \dots \\ &\quad + \sum_{i < j < k} P(A_i A_j A_k) - \dots + (-1)^{m+1} P(A_1 \dots A_m) \end{aligned}$$

$P(A_i)$ is the probability that each of the first n trials results in a non- i outcome, and so by independence $P(A_i) = (1 - p_i)^n$

Similarly, $P(A_i A_j)$ is the probability that the first n trials all result in a non- i and non- j outcome, and so

$$P(A_i A_j) = (1 - p_i - p_j)^n$$

As all of the other probabilities are similar, we see that

$$\begin{aligned} P\{X > n\} &= \sum_{i=1}^m (1 - p_i)^n - \sum_{i < j} (1 - p_i - p_j)^n \\ &\quad + \sum_{i < j < k} (1 - p_i - p_j - p_k)^n - \dots \end{aligned}$$

Since $P\{X = n\} = P\{X > n - 1\} - P\{X > n\}$, we see, upon using the algebraic identity $(1 - a)^{n-1} - (1 - a)^n = a(1 - a)^{n-1}$, that

$$\begin{aligned} P\{X = n\} &= \sum_{i=1}^m p_i * (1 - p_i)^{n-1} - \sum_{i < j} (p_i + p_j) * (1 - p_i - p_j)^{n-1} \\ &\quad + \sum_{i < j < k} (p_i + p_j + p_k) (1 - p_i - p_j - p_k)^{n-1} - \dots \end{aligned}$$

CDF The cumulative distribution function(cdf) $F(\cdot)$ of the random variable X is defined for any real number b , $-\infty < b < \infty$, by

$$F(b) = P\{X \leq b\}$$

We have

$$P\{a < X \leq b\} = F(b) - F(a), \text{ for all } a < b$$

$$P\{X < b\} = \lim_{h \rightarrow 0^+} P\{X \leq b - h\} = \lim_{h \rightarrow 0^+} F(b - h)$$

2 Discrete Random Variables

2.1 The Bernoulli Random Variable

Suppose that a trail, or an experiment, whose outcome can be classified as either a "success" or as a "failure" is performed. If we let X equal 1 if the outcome is a success and 0 if it is a failure, then the probability mass function of X is given by

$$p(0) = P\{X = 0\} = 1 - p$$

$$p(1) = P\{X = 1\} = p$$

where $p, 0 \leq p \leq 1$, is the probability that the trail is a "success".

A random variable X is said to be a *Bernoulli random variable* if its probability mass function is given by the above equation for $p \in (0, 1)$.

2.2 The Binomial Random Variable

Suppose that n independent trails, each of which results in a "success" with probability p and in a "failure" with probability $1 - p$, are to be performed. If X represents the number of successes that occur in the n trails, then X is said to be a *binomial random variable* with parameters (n, p) .

The probability mass function of a binomial random variable having parameters (n, p) is given by

$$p(i) = \binom{n}{i} p^i * (1 - p)^{n-i}, i = 0, 1, \dots, n$$

where

$$\binom{n}{i} = \frac{n!}{(n-i)! * i!}$$

Note that, by the binomial theorem, the probabilities sum to one, that is,

$$\sum_{i=0}^{\infty} p(i) = \sum_{i=0}^n p^i * (1 - p)^{n-i} = (1 + (1 - p))^n = 1$$

2.3 The Geometric Random Variable

Suppose that independent trails, each having probability p of being a success, are performed until a success occurs. If we let X be the number of trails required until the first success, then X is said to be a *geometric random variable* with parameter p . Its probability mass function is given by

$$p(n) = P\{X = n\} = (1 - p)^{n-1} * p, n = 1, 2, \dots$$

To check that $p(n)$ is a probability mass function, we note that

$$\sum_{n=1}^{\infty} p(n) = \sum_{n=1}^{\infty} (1 - p)^{n-1} = 1$$

2.4 The Poisson Random Variable