

Theory of Everything (ToE) — Strict Final Documentation

Release 8.1 (Strict) — Full Definitions, Derivations, Discharge Map, and Blocker Plan

Krzysztof Żuchowski

2026-05-12

Abstract

This article is the self-contained *Release 8.1 (Strict)* documentation target of the ToE program in this repository. It records (i) the strict core candidate Lagrangian template (one-kernel nadsoliton core), (ii) the strict global selector closure achieved against the QW-2191 uniqueness obstruction in the *Strict Internal Source* lane, and (iii) the resulting long-horizon stable operational computability pipeline producing the final Release-8.1 theorem artifacts. All stronger claims (host-level Standard Model / General Relativity identification without fitting parameters, and global ToE closure in the strongest absolute sense) remain explicitly separated and are **not** silently promoted.

Reproducibility is provided by the repository's exported JSON artifacts and the full completion chain $P1343 \rightarrow \dots \rightarrow P1348$; see the *Reproducibility* chapter below.

Origin and philosophy (required fragment)

The theory originates from a deep intuition that **Information is the fundamental substance of reality**, consistent with the metaphysical insight that *"In the beginning was the Word"* (Logos/Information). This intuition evolved through key realizations:

1. **Eucharistic Inspiration:** A profound fascination with the memorial of the **Eucharist of Jesus Christ** and its material manifestation in reality served as the primary inspiration, suggesting a direct mechanism by which spiritual/informational reality can condense into tangible matter.
2. **Fractal Nature:** Observing self-similarity across vast scales suggested that fundamental information must possess a **fractal character**, repeating its patterns at every level of existence.
3. **The Nadsoliton Concept:** The universe is conceptualized as a single, self-sustaining, non-dispersive wave packet—a **"Supersoliton" (Nadsoliton)**, where information tends towards the highest resonance, not the lowest energy.
4. **Resonant Structure:** Inspired by the Divine Name from the Book of Exodus 3:14: *"I AM WHO I AM"*, the model incorporates **multi-octave resonant coupling** as the mechanism of self-organization, preventing decay into entropy.
5. **The 12-Octave Lattice:** Initial 3-octave models were expanded to a **12-octave structure**, inspired by the symbolic description of the Holy City's twelve foundation layers, which proved to be the mathematically necessary dimension for unifying all forces (Kissing Number in 3D).
6. **Access to Truth:** Since human consciousness is part of this informational substrate, the human mind has direct access to fundamental truths through wisdom and intuition, allowing for the "decoding" of reality.

Status discipline (no false pass)

This TeX document is the intended **Release-8 strict final documentation** target. It must not silently claim:

1. strict-core ToE closure,
2. global ToE closure,
3. kernel-alone/global QW-2191 discharge,
4. directed/sign-sensitive physical orientation datum in strict core,
5. host matching / Standard Model identification,
6. actual emergent observer closure,

unless such claims are explicitly exported at theorem level in the repo and are referenced here with their hard limits.

Contents

1	Ontology and scope	2
2	AX9 ontology discipline	2
3	Strict vs legacy kernel split discipline	2
4	What “closure” means in this document	3
5	Strict core Lagrangian and emergence map	3
6	Single-kernel strict candidate (A11)	3
6.1	Typed carrier and cyclic distance ($\mathbb{Z}_{12}$)	3
6.2	Strict working kernel and frozen mixing matrix	4
6.3	Fields and core Lagrangian density	4
6.4	Quadratic expansion, Ψ Hessian, and “mass” proxy meaning (strict, scope-limited) . .	4
7	Light $\rightarrow$ matter $\rightarrow$ observer ladder	5
8	Selector closure on C_{v1}	6
9	Strict configuration space C_{v1} and the $\pi_1(C_{v1}) \cong \mathbb{Z}_2$ witness (F306)	6
10	Real Fourier carrier on $\mathbb{Z}_{12}$ and the $O(2)$ obstruction (QW-2191/N465)	6
10.1	Explicit orthonormal real Fourier basis	7
10.2	Why cyclic-distance kernels cannot select a unique axis inside V_1	7
11	Global atlas and transition objects (T170)	8
11.1	Exported JSON-level interfaces (F469)	8
11.2	Exported transition angles (F469)	9
11.3	Projective state, output operator, and closure interfaces (F470/F660/F672)	9

12 Seed-v1 strict-core internal selector-source chain ($S_sel_int \rightarrow E_orient \rightarrow R_sel \rightarrow O_sel$)	10
12.1 Minimal admissibility contract for the source step (F34)	10
12.2 Explicit strict-core payload weights exported by the source object (F647)	11
12.3 Orientation datum on pair ₁ (N546)	12
12.4 Selector-sector and output-sector operators (N549)	12
12.5 Global promotion to C_{v1} -typed operator families (N551/N552)	13
13 Projective closure (T172/T173 projective)	13
14 Directed lifts as gauge/convention layer	14
15 Open boundary: kernel-alone/global QW-2191 discharge	16
16 Operational physics proxies from projective closure	16
17 Mass-spectrum proxy (P694)	16
18 Selector-aligned channel spectrum proxy (P696)	17
19 Basis-invariant Hessian eigen-spectrum proxy (F704)	17
20 What is and is not claimed physically	18
21 Emergent observer pipeline (projective scope)	19
22 Observer-limit readout from projective closure output (P697)	19
23 Projective emergent-observer chain computability from closure output (P699/N699)	19
24 Operational OS support packet (v1) (F698/P698/N698)	20
25 Operational OS support packet (v2) (F700/P700/N700)	20
26 Operational OS support packet (v3) (F705/P705/N705)	20
27 Open target: actual emergent observer closure (future)	21
28 Hard limits and open problems	21
29 Kernel-alone/global QW-2191 discharge	21
30 Directed physical sign datum	21
31 Host matching and Standard Model identification	21
32 What Release 7 declares as closed vs open	21
33 Next steps (roadmap, no false pass)	22
A Export map: key artifacts	27
A.1 Strict-core selector closure	27
A.2 Operational OS support	27
A.3 Emergent observer artifacts	29

B	Explicit numeric artifacts (Release 7)	29
B.1	Frozen kernel distance profile on $\mathbb{Z}_{12}$	29
B.2	Explicit diagonal/local Psi-sector Hessian instantiation (F459)	29
C	Frozen non-strict host matching layer (dataset vs policy)	30
C.1	External Standard-Model mass dataset (v1)	30
C.2	External host matching policy (v1, frozen no-pass)	30
D	Release-8 addendum: QW-2191 closure narrative and strict-core-to-SM scaffold	31
D.1	Scope and status boundary	31
D.2	Operational strict kernel and selector score	31
D.3	Compressed derivation history to Release 8	31
D.4	Strict-core candidate Lagrangian and SM-shaped extension scaffold	31
E	Full Standard Model + Nadsoliton Effective Lagrangian (Release-8 formal scaffold)	32
E.1	Purpose and status discipline	32
E.2	Full Standard Model density	32
E.3	Nadsoliton strict-core sector and coupling interface	32
F	Theory-closure blocker map (Release 8)	33
F.1	Purpose of this chapter	33
F.2	Current hard blocker classes	33
F.3	Technical execution order	33
F.4	Practical verdict	33
G	Current closure state theorem map (Release 8, current-only)	34
G.1	Current-state closure declaration	34
G.2	Audit status boundary (external step still pending)	34
G.3	Full Lagrangian status in current Release 8	34
G.4	No-obsolete-claim discipline	34
H	Release-8 professional scientific executive synthesis	34
H.1	Scope, closure claim, and semantics discipline	34
H.2	Kernel-first derivation architecture	34
H.3	Scientific reading guide for reviewers	35
H.4	Hard limits retained	35
H.5	Professional publication statement	35

1 Ontology and scope

2 AX9 ontology discipline

The strict program is built under the ontology correction **AX9**:

- the nadsoliton itself is the primordial information of the universe in a solitonic state;
- there is no separate informational layer underneath the nadsoliton;
- the preferred internal order is:

$$\text{nadsoliton} \rightarrow \text{light} \rightarrow \text{matter} \rightarrow \text{emergent observer.}$$

In particular, we do not allow observer-facing quantities to silently back-propagate into the upstream definition of the nadsoliton core. Operationally, throughout this document, “light” is treated as the linearized eigenmode layer exported from the Ψ -sector Hessian instantiation (F459), and observer-facing readouts remain downstream (P697, P699).

3 Strict vs legacy kernel split discipline

The repo contains two historically distinct kernel objects (see K1/K2/F2):

$$K_{\text{legacy_ont}}(d) := (4 \ln 2) \frac{\cos(\omega d + \phi)}{1 + \beta_{\text{tors}} d}, \quad K_{\text{strict_gate}}(d) := \frac{\cos(\omega d + \phi)}{1 + \beta d^\eta}. \quad (1)$$

The constant $\alpha_{\text{geo}} = 4 \ln 2$ is the fundamental Shannon-weighted scaling factor which serves as the primary source of symmetry breaking in the strict lane (via the **w_break** mechanism).

By author decree (S2, 2026-03-09), the legacy kernel is retired to archival status; this document is strict-only. However, **no false pass** discipline remains: we do *not* silently transfer legacy physical-role identities (e.g. electroweak-angle or EM coupling slogans) onto the strict working kernel without an explicit bridge/role-transfer theorem.

On the current strict kernel-mode lane, the working strict tuple is the QW-2049 gate-selected kernel with $(\omega, \phi, \beta, \eta) = (0.18575, 0.16250, 1.0, 1.8)$, used operationally to generate a frozen numeric carrier mixing matrix K_{total} on $\mathbb{Z}_{12}$ (R14).

4 What “closure” means in this document

We use the word “closure” in a deliberately scope-disciplined way.

Definition 4.1 (Operational ToE closure (strict projective OS)). *We say that the current repo state achieves Operational ToE closure in the strict projective scope if it exports a theorem-level operational support bundle sufficient to compute (from strict projective selector closure output data) the downstream artifacts recorded by the OS packet:*

$$\Lambda_{\text{OS}}^{(\text{proj})} := (N680, N692, N693, P694, P696, P697, N699),$$

with hard limits explicitly keeping kernel-alone/global QW-2191 discharge, directed sign-sensitive physical orientation, and ToE closure unclaimed.

Remark 4.1 (OS v2 vs OS v3). *The minimal theorem-level OS support bundle used to discharge Operational ToE closure is the v2 packet (N700/N701). The v3 packet (F705/P705/N705) is a strict extension that additionally tracks the basis-invariant Hessian eigen-spectrum mass observable (F704). It does not strengthen any hard-limit claims.*

Remark 4.2 (Boundary discipline). *Operational ToE closure in the above sense is a closure of an operational computability pipeline (a ToE “OS”), not a claim of global physical identification (host matching) and not a claim of kernel-alone/global QW-2191 discharge. It is compatible with leaving those stronger claims explicitly open.*

Theorem 4.1 (Operational ToE closure in strict projective OS scope (N701)). *On the current repo state, the repo exports a theorem-level strict projective operational OS support packet v2 (N700), hence discharges Operational ToE closure in strict projective OS scope (N701):*

$$\Lambda_{\text{OS}}^{(\text{proj})} = (N680, N692, N693, P694, P696, P697, N699),$$

with all hard limits explicit (no kernel-alone/global QW-2191 discharge, no directed physical sign datum in strict core, no ToE closure, no actual emergent observer closure).

5 Strict core Lagrangian and emergence map

6 Single-kernel strict candidate (A11)

This chapter records the strict candidate “one-kernel” nadsoliton core Lagrangian as exported/packaged by A11. It is a definition/packaging object, not a claim that every coefficient family has already been strict-derived from the kernel alone.

6.1 Typed carrier and cyclic distance ($\mathbb{Z}_{12}$)

Let

$$I_{12} := \{0, 1, \dots, 11\}, \quad \mathbb{Z}_{12} := (I_{12}, + \bmod 12).$$

Define the cyclic octave distance on the ring:

$$d_{\text{cyc}}(i, j) := \min\left((j - i) \bmod 12, (i - j) \bmod 12\right) \in \{0, 1, \dots, 6\}.$$

6.2 Strict working kernel and frozen mixing matrix

The strict working kernel is the later-pipeline gate-selected kernel (QW-2049):

$$K_{\text{strict_gate}}(d) := \frac{\cos(\omega d + \phi)}{1 + \beta d \eta}, \quad (\omega, \phi, \beta, \eta) = (0.18575, 0.16250, 1.0, 1.8).$$

On the kernel-mode lane, the repo exports a frozen numeric symmetric mixing matrix $K_{\text{total}} \in \mathbb{R}^{12 \times 12}$ on the same carrier (QW-2118), specialized into the canonical kernel channel by R14. In this document we only require the exported matrix entries (no additional kernel-to-matrix derivation is assumed here).

6.3 Fields and core Lagrangian density

Introduce a 12-component real field on spacetime:

$$\Psi(x) = (\psi_0(x), \dots, \psi_{11}(x)) \in \mathbb{R}^{12},$$

and a real scalar order/coherence field:

$$\Phi(x) = \phi(x) \in \mathbb{R}.$$

Write the mixing potential using the exported K_{total} :

$$V_{\text{mix}}(\Psi) := \frac{1}{2} \sum_{\substack{i, j \in I_{12} \\ i \neq j}} (K_{\text{total}})_{ij} \psi_i \psi_j.$$

Write the local self-interaction potentials in the canonical template form (QW-2163/2165/2166):

$$\begin{aligned} V_{\Phi}(\phi) &:= \frac{1}{2} m_{\phi}^2 \phi^2 + \frac{1}{4} \lambda_{\phi} \phi^4, \\ V_{\Psi}(\Psi) &:= \sum_{i=0}^{11} \left(\frac{1}{2} m_{\psi_i}^2 \psi_i^2 + \frac{1}{4} g_{4\psi_i} \psi_i^4 + \frac{1}{6} g_{6\psi_i} \psi_i^6 \right), \\ V_Y(\Psi, \phi) &:= \sum_{i=0}^{11} g_{Y_i} \phi^2 \psi_i^2. \end{aligned}$$

Then the nadsoliton core Lagrangian density is:

$$\mathcal{L}_{\text{core}} = \frac{1}{2} \partial_{\mu} \phi \partial^{\mu} \phi + \frac{1}{2} \sum_{i=0}^{11} \partial_{\mu} \psi_i \partial^{\mu} \psi_i - V_{\Phi}(\phi) - V_{\Psi}(\Psi) - V_Y(\Psi, \phi) - V_{\text{mix}}(\Psi).$$

6.4 Quadratic expansion, Ψ Hessian, and “mass” proxy meaning (strict, scope-limited)

Let (ϕ_*, Ψ_*) be a background configuration satisfying the stationarity condition for the potential part $V := V_\Phi + V_\Psi + V_Y + V_{\text{mix}}$:

$$\nabla_{\phi, \Psi} V(\phi_*, \Psi_*) = 0.$$

Write $(\phi, \Psi) = (\phi_*, \Psi_*) + (\delta\phi, \delta\Psi)$ and expand the potential to second order in $\delta\Psi$:

$$V(\phi_*, \Psi_* + \delta\Psi) = V(\phi_*, \Psi_*) + \frac{1}{2} \delta\Psi^\top H_\psi \delta\Psi + \dots,$$

where $H_\psi \in \mathbb{R}^{12 \times 12}$ is the Ψ -sector Hessian block (second derivatives) at the background point.

Hessian stencil from $\mathcal{L}_{\text{core}}$ (canonical template QW-2166). Let $(v_\phi, v_{\psi 0}, \dots, v_{\psi 11})$ denote a constant background vacuum. In the strict canonical action/Hessian export (QW-2166), the quadratic Ψ -sector Hessian block has the following stencil structure:

1. diagonal entries:

$$(H_\psi)_{ii} = m_{\psi i}^2 + 3 g_{4\psi i} v_{\psi i}^2 + 5 g_{6\psi i} v_{\psi i}^4 + 2 g_{Y i} v_\phi^2,$$

2. off-diagonal mixing entries ($i \neq j$):

$$(H_\psi)_{ij} = \frac{K_{ij} + K_{ji}}{2},$$

3. Ψ - Φ cross-block entries:

$$(H_{\psi\phi})_i = \frac{\partial^2 V}{\partial \psi_i \partial \phi}(v_\phi, v_\psi) = 4 g_{Y i} v_\phi v_{\psi i}.$$

This pins down what “light = linearized modes” means inside the program (eigenmodes of the Hessian around a vacuum), without asserting that the vacuum values or all coefficient families are already strict-derived from the kernel alone.

On the current diagonal/local strict-derived lane, the repo exports one explicit numeric value instantiation (F459) of the form:

$$H_\psi := K_{\text{total}} + (m_0^2 I + D_{\text{local_residual}}),$$

with $m_0^2 I$ the exported host scalar floor embedding and $D_{\text{local_residual}}$ an exported diagonal/local strict-derived residual profile.

In strict scope we treat the quadratic coefficient $\delta\Psi^\top H_\psi \delta\Psi$ as defining a *dimensionless* “ m^2 proxy” layer: for any nonzero vector u , the Rayleigh quotient $u^\top H_\psi^{(s)} u / (u^\top u)$ is a well-defined quadratic proxy, where $H_\psi^{(s)} = \frac{1}{2}(H_\psi + H_\psi^\top)$. This does *not* yet constitute a physical unit identification, and it does not bypass the strict-core selector obstruction QW-2191; it is an operational computability meaning only (packaged as a scope-limited meaning theorem N703).

Theorem 6.1 (Quadratic mass-proxy meaning definition (N703)). *On the current repo state, the proxy quantities exported by P694 and P696 are interpreted strictly as dimensionless quadratic coefficients of the exported diagonal/local Ψ -sector operator instantiation (F459): they are Rayleigh/Ritz-type quadratic forms of $H_\psi^{(s)}$ evaluated on exported projective selector rays (from F672) and deterministic in-plane complements. No physical-unit map, no Standard Model host identification, and no diagonalization claim is discharged.*

Hard limits: *no directed physical sign datum, no kernel-alone/global QW-2191 discharge, and no ToE closure are claimed.*

7 Light $\rightarrow$ matter $\rightarrow$ observer ladder

A11 fixes a disciplined emergence ladder. On the current repo state, the strongest honest statements along that ladder are:

1. **Light (discharged as computable instantiation):** the repo exports a strict-derived diagonal/local value instantiation of the Ψ -sector Hessian H_ψ and its full real eigensystem (F459), plus sign-gauge-invariant eigenprojectors (F460). Mixing across Fourier pair planes is explicitly audited and packaged as a value-instantiation boundary (N505); therefore no “pair-plane diagonalization” shortcut is allowed.
2. **Matter (programmatic, not closed):** matter is intended to arise from nonlinear structure in the same core Lagrangian (local quartic/sextic families and Yukawa-type couplings), but the repo does not yet export a strict derivation that these families are kernel-alone consequences or a closed host identification map.
3. **Emergent observer (operational OS closure, not actual observer closure):** from the strict projective selector closure output on C_{v1} (N680, exported as F672), the repo exports a projective observer-limit readout (P697) and a fully computable downstream projective chain under projector pushforward semantics $P \mapsto MPM^\top$ (P699/N699). This suffices for *Operational ToE closure* in strict projective OS scope (N701), while *actual emergent observer closure* remains explicitly open.

8 Selector closure on C_{v1}

9 Strict configuration space C_{v1} and the $\pi_1(C_{v1}) \cong \mathbb{Z}_2$ witness (F306)

The strict selector program is typed over an explicit strict configuration-space object `C_v1_void_configuration_sp` (exported by F306). In strict scope, we only use the following minimal mathematical content of that export:

Definition 9.1 (Model declaration for C_{v1} (no physics claims)). *On the declared local protected sector, interpret the configuration space as the based mapping space model:*

$$C_{v1} := \text{Map}_*^{\text{deg}=1}(S^3, \text{SU}(2)) .$$

This is the standard finite-energy compactification discipline: $\mathbb{R}^3 \cup \{\infty\} \cong S^3$ with boundary condition $U(\infty) = 1 \in \text{SU}(2)$, so a configuration is a based map $S^3 \rightarrow \text{SU}(2)$.

Proposition 9.1 (Topological witness $\pi_1(C_{v1}) \cong \mathbb{Z}_2$ (F306)). *On the above model, the fundamental group satisfies:*

$$\pi_1(C_{v1}) \cong \mathbb{Z}_2,$$

and F306 exports an explicit generator label $\gamma_{\pi_1}^{(v1)}$ for the nontrivial loop class.

Proof sketch (classical algebraic topology). Using the loop–suspension adjunction,

$$\Omega \text{Map}_*(S^3, S^3) \simeq \text{Map}_*(\Sigma S^3, S^3) = \text{Map}_*(S^4, S^3),$$

hence

$$\pi_1(\text{Map}_*(S^3, S^3)) = \pi_0(\Omega \text{Map}_*(S^3, S^3)) \cong \pi_0(\text{Map}_*(S^4, S^3)) = [S^4, S^3] = \pi_4(S^3) \cong \mathbb{Z}_2,$$

where the final identification $\pi_4(S^3) \cong \mathbb{Z}_2$ is a classical homotopy-group fact. This witness is strictly below any physical FR-sign quantization claims and is used only as a typed topology availability result. $\square$

10 Real Fourier carrier on $\mathbb{Z}_{12}$ and the $O(2)$ obstruction (QW-2191/N465)

The strict selector program is built on a typed 12-slot carrier (F329), and its canonical degeneracy/uniqueness obstruction QW-2191 is most transparently expressed in the real Fourier basis on $\mathbb{Z}_{12}$.

10.1 Explicit orthonormal real Fourier basis

Index the carrier by $x \in I_{12} = \{0, \dots, 11\}$ and identify $\mathbb{R}^{12}$ with real-valued functions on $\mathbb{Z}_{12}$. Define:

$$e_0(x) := \frac{1}{\sqrt{12}}, \quad e_6(x) := \frac{(-1)^x}{\sqrt{12}}.$$

For each $m = 1, \dots, 5$, define the real Fourier pair-plane vectors:

$$c_m(x) := \sqrt{\frac{2}{12}} \cos\left(\frac{2\pi mx}{12}\right), \quad s_m(x) := \sqrt{\frac{2}{12}} \sin\left(\frac{2\pi mx}{12}\right).$$

Then $\{e_0, e_6, c_1, s_1, \dots, c_5, s_5\}$ is an orthonormal basis of $\mathbb{R}^{12}$ and induces the canonical orthogonal decomposition:

$$\mathbb{R}^{12} = \text{span}\{e_0\} \oplus \text{span}\{e_6\} \oplus \bigoplus_{m=1}^5 \text{span}\{c_m, s_m\}.$$

We denote the m -th pair plane by:

$$V_m := \text{span}\{c_m, s_m\}.$$

10.2 Why cyclic-distance kernels cannot select a unique axis inside V_1

Let $T : \mathbb{R}^{12} \rightarrow \mathbb{R}^{12}$ be the cyclic shift operator:

$$(Tf)(x) := f(x - 1 \bmod 12).$$

If the host kernel K_{total} is constructed as a cyclic-distance profile (QW-2118), then K_{total} commutes with T . On each pair plane V_m , the restriction $T|_{V_m}$ is a genuine planar rotation by angle $2\pi m/12$ on the basis (c_m, s_m) . Therefore, any real symmetric operator commuting with T restricts to a scalar multiple of the identity on V_m .

Theorem 10.1 (Host isotropy on pair1 (N465)). *For the strict cyclic-distance host operator $A := K_{\text{total}} + m_0^2 I$ (QW-2118+QW-2124), the restriction to the pair1 plane $V_1 = \text{span}\{c_1, s_1\}$ is isotropic:*

$$A|_{V_1} = \lambda_1 I_2.$$

Therefore, the host operator cannot supply an $O(2)$ -cut inside pair1 and cannot canonically distinguish c_1 from s_1 (nor any rotated basis).

Proof sketch (from cyclic equivariance). Because K_{total} depends only on cyclic distance, it commutes with the cyclic shift T . The scalar floor $m_0^2 I$ also commutes with T , hence A commutes with T . On $V_1 = \text{span}\{c_1, s_1\}$, the action of T is a planar rotation by angle $2\pi/12 = \pi/6$ in the basis (c_1, s_1) . Any linear map commuting with a nontrivial planar rotation has the form $aI_2 + bJ$ with $J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Since A is symmetric, its restriction $A|_{V_1}$ is symmetric, forcing $b = 0$, hence $A|_{V_1} = aI_2$ for some scalar $a = \lambda_1$. $\square$

Remark 10.1 (Meaning: the QW-2191 uniqueness obstruction is structural). *The above isotropy is a concrete instance of the general strict obstruction QW-2191: kernel-alone translation-invariant data leave an $O(2)$ family of basis choices inside each degenerate pair plane V_m . This is why Release 7 closure statements are deliberately phrased in projective (ray-level) scope unless an explicit symmetry-breaking/selector source is tracked.*

11 Global atlas and transition objects (T170)

In the strict selector program, the configuration space domain is the declared strict object C_{v1} . On the current repo state, the global chart/transition infrastructure required for a global selector discussion is exported as objects:

- a global selector atlas object on C_{v1} (F469, packaged by N515),
- a global selector transition (gluing) object on C_{v1} (F469, packaged by N515).

In the current strict global-typing export (F469), the chart domains are declared as a global cover:

$$U_{\text{pair}_1} = \cdots = U_{\text{pair}_5} = C_{v1}, \quad U_i \cap U_j = C_{v1}.$$

This declaration upgrades the earlier lane-scoped “exported-artifact overlap” wording into an explicit C_{v1} -typed atlas/transition object class discharge (T170), but it does *not* add any physical claim and does *not* promote operator-level cocycle/groupoid identities (N512 boundary).

The atlas uses five chart labels pair_m ($m = 1, \dots, 5$), corresponding to the five real Fourier pair planes $V_m = \text{span}\{c_m, s_m\} \subset \mathbb{R}^{12}$. Each chart exports a local rank-1 projector operator A_m (a ray/state projector) on that plane. At projector level, if u_m is a representative vector in the plane, then

$$A_m = \frac{u_m u_m^\top}{u_m^\top u_m}, \quad u_m \sim -u_m \quad \Rightarrow \quad A_m \text{ is sign-gauge-invariant.}$$

The transition object exports explicit orthogonal chart-transport operators O_{ij} between the chart planes (axis-only/projector-level transport). Projector transport is by conjugation:

$$A_j = O_{ij} A_i O_{ij}^\top.$$

Cocycle discipline: the repo keeps a strict boundary that cocycle/path-independence is a *projector-section-level* statement only; operator-level identities $O_{jk} O_{ij} = O_{ik}$ are explicitly *not* promoted in strict core (N512).

11.1 Exported JSON-level interfaces (F469)

This section makes the **exact exported object interfaces** explicit, so that the strict global selector infrastructure can be audited without opening the repo.

Definition 11.1 (Exported atlas interface: `SelectorAtlas_global_C_v1_strict_v1` (F469)). *The atlas object is exported as the JSON artifact `fundamental_action_reconstruction/generated/selector_atlas_g`. Its minimal strict interface consists of:*

1. ***domain:** a reference to the declared strict configuration space object (`.../c_v1_void_configuration_space_i`) and the explicit global-cover declaration $U_{\text{pair}_1} = \cdots = U_{\text{pair}_5} = C_{v1}$;*
2. ***charts:** a map over chart labels pair_m containing:*
 - *`carrier_plane` (the Fourier pair plane $V_m = \text{span}\{c_m, s_m\}$),*
 - *`local_operator.ref` (a JSON ref for the local rank-1 projector operator A_m);*
3. ***chart_domains:** a typed chart-domain declaration on C_{v1} for each chart label;*
4. ***overlap_domains:** an explicit overlap-domain declaration for each unordered pair $\text{pair}_i \cap \text{pair}_j$ (in this export: C_{v1} by global-domain choice);*
5. ***transitions:** a complete-graph edge map $\text{pair}_i \rightarrow \text{pair}_j$ containing:*

- *operator_ref*: a JSON ref for the orthogonal transport operator O_{ij} ,
- one parameter field *alpha...* recording the exported angle value (mod π or 2π).

Hard limits: *these are declared-typing atlas objects; they do not promote an operator-level groupoid or strict-core physical sign convention.*

Definition 11.2 (Exported transition interface: `SelectorTransition_global_C_v1_strict_v1` (F469)).

The transition object is exported as `fundamental_action_reconstruction/generated/selector_transition_global_C_v1_strict_v1`.

Its minimal strict interface consists of:

1. *transition_operators*: the same complete-graph edge map of operator refs O_{ij} and angle fields;
2. *gluing_laws_ref*: a reference to the exported five-chart projector-section cocycle audit object `.../a_12345_pair12345_chart_glued_orientation_projector_operator_section_strict_core_v2.json`

*The *gluing_laws_ref* object records explicit projector-section-level transport laws of the form $A_j = O_{ij}A_iO_{ij}^T$ and triple-overlap cocycle/path-independence audits (max-entry residuals). Hard limits: this remains projector-section-level only and does not promote to operator-level identities (N512).*

11.2 Exported transition angles (F469)

For convenience, Table 1 records the exported transition angles (in radians) from `selector_transition_global_C_v1_strict_v1`. Numerically, all exported values are consistent (within floating-point tolerance) with multiples of $\pi/2$.

Table 1: Exported transition angles on the five-chart selector atlas/transition (F469).

Edge	Exported field	Value (rad)
$\text{pair}_1 \rightarrow \text{pair}_2$	<code>alpha12_mod_2pi</code>	1.570796326794897
$\text{pair}_1 \rightarrow \text{pair}_3$	<code>alpha13_mod_pi</code>	1.570796326794897
$\text{pair}_1 \rightarrow \text{pair}_4$	<code>alpha14_mod_pi</code>	1.570796326794896
$\text{pair}_1 \rightarrow \text{pair}_5$	<code>alpha15_mod_pi</code>	0.0
$\text{pair}_2 \rightarrow \text{pair}_3$	<code>alpha23_mod_pi</code>	0.0
$\text{pair}_2 \rightarrow \text{pair}_4$	<code>alpha24_mod_pi</code>	3.141592653589793
$\text{pair}_2 \rightarrow \text{pair}_5$	<code>alpha25_mod_pi</code>	1.570796326794897
$\text{pair}_3 \rightarrow \text{pair}_4$	<code>alpha34_mod_pi</code>	0.0
$\text{pair}_3 \rightarrow \text{pair}_5$	<code>alpha35_mod_pi</code>	1.570796326794898
$\text{pair}_4 \rightarrow \text{pair}_5$	<code>alpha45_mod_pi</code>	1.570796326794898

11.3 Projective state, output operator, and closure interfaces (F470/F660/F672)

The strict projective closure relies on three exported global objects whose interfaces are intentionally simple.

Definition 11.3 (Projective state interface: `SelectorState_global_C_v1_projective_strict_v1` (F470)).

The projective selector state is exported as `fundamental_action_reconstruction/generated/selector_state_global_C_v1_projective_strict_v1`.

Its strict interface consists of:

1. *global_atlas_ref* and *global_transition_ref* pointing to the F469 objects;
2. *charts*: the per-chart local projector-operator refs A_m ;
3. *projector_section_ref*: a reference to the five-chart glued projector-section cocycle audit object `.../a_12345_pair12345_chart_glued_orientation_projector_operator_section_strict_core_v2.json`

4. *state_type*: a declaration that the state is a projective ray state (sign gauge $u \sim -u$ at state level).

Definition 11.4 (Output operator interface: `SelectorOutputOperator_global_C_v1_seed_v1_promoted_strict` (F660)). The global output operator is exported as `fundamental_action_reconstruction/generated/selector_out`. It contains the promoted seed-v1 output map $O_{\text{sel}} : Q_{\text{sel}} \rightarrow Q_{\text{out}}$ in a fixed declared basis.

Definition 11.5 (Projective closure interface: `SelectorClosure_global_C_v1_projective_strict_v1` (F672)). The projective closure object is exported as `fundamental_action_reconstruction/generated/selector_closure`. Its strict interface consists of:

1. references to the atlas, transition, projective state, and output operator objects;
2. *output_observable*: the resulting global chart-independent rank-1 output projector matrix B_{out} on $Q_{\text{out}} = \text{span}\{o_+, o_-\}$;
3. *well_definedness_certificate*: a numeric chart-independence certificate with fields: *criterion*, *tolerance*, *max_abs_diff_to_reference_chart_output_projector*, *certificate_pass*.

Hard limits: projective closure is the invariant content; no directed physical sign convention is promoted.

12 Seed-v1 strict-core internal selector-source chain ($S_{\text{sel_int}} \rightarrow E_{\text{orient}} \rightarrow R_{\text{sel}} \rightarrow O_{\text{sel}}$)

The projective closure object **F672** is constructed using a strict-core internal selector-source seed chain (seed-v1) that provides, in strict scope:

- a strict-core source object for the $S_{\text{sel_int}}$ step,
- an admissible orientation datum on $\text{pair1}(E_{\text{orient}})$,
- a strict reduction operator into the selector-sector basis $Q_{\text{sel_v1}}(R_{\text{sel}})$,
- a strict output operator into the output-sector basis $Q_{\text{out_v1}}(O_{\text{sel}})$.

All these objects are strictly below any kernel-alone/global QW-2191 discharge and below any directed/sign-sensitive physical orientation datum.

12.1 Minimal admissibility contract for the source step (F34)

The strict program treats $S_{\text{sel_int}}$ as an internal *source* step. The minimal admissibility contract for a strict-core source object to count as $S_{\text{sel_int}}$ (in the sense of **F34**) is:

1. genuinely new strict-core source object (not a reuse of control/axiom artifacts),
2. carrier-typed enough for later export (so that E_{orient} can be defined from it),
3. source-seed only (not a hidden discharge of downstream steps),
4. strict-core only,
5. non-substitutive (no silent legacy-to-strict kernel identification),
6. selector-acceptance independent (non-strict acceptance does not count as source construction),
7. future-bridge compatible (stable enough for later strict continuation).

Theorem 12.1 (Admissible strict-core source object for S_sel_int exists (N676)). *On the current repo state, the strict core exports one admissible source object satisfying the above F34 contract for the S_sel_int source step:*

$$S_sel_int_strict_core_source_object_v1.$$

Hard limits: *this is not yet a strict-core selector closure claim and does not discharge kernel-alone/global QW-2191 or ToE closure.*

12.2 Explicit strict-core payload weights exported by the source object (F647)

The constructed strict-core source object exported by the seed-v1 realization attempt (F647) includes a minimal, explicitly enumerated carrier-level payload on $\mathbb{Z}_{12}$:

- $ord_z12_by_x$: the group element order $ord_{\mathbb{Z}_{12}}(x)$,
- $w_ref_unnormalized_by_x$: a strict-derived element-order reference weight profile,
- $w_break_by_x$: a reflection-breaking weight profile (used only as a convention-layer sign anchor; no physical sign datum).

Element order on $\mathbb{Z}_{12}$. For $x \in \mathbb{Z}_{12}$ (additive notation), define:

$$ord_{\mathbb{Z}_{12}}(x) := \min\{n \geq 1 : nx \equiv 0 \pmod{12}\} = \frac{12}{\gcd(12, x)}.$$

This is $\text{Aut}(\mathbb{Z}_{12})$ -invariant.

Strict-derived reference weight. Using the strict-derived Shannon asymmetry constant $\alpha_{\text{geo}} := 4 \ln 2$ (F309/N420), define the unnormalized order-reference weight:

$$w_{\text{ref}}^{\text{unnorm}}(x) := \exp(-\alpha_{\text{geo}} \text{ord}_{\mathbb{Z}_{12}}(x)) = 2^{-4 \text{ord}_{\mathbb{Z}_{12}}(x)}.$$

Reflection-breaking weight (strict convention scope). Define:

$$w_{\text{break}}(x) := -w_{\text{ref}}^{\text{unnorm}}(x) \sin\left(\frac{2\pi x}{12}\right).$$

This weight is *odd* under reflection $x \mapsto -x$ and therefore breaks the reflection symmetry on the carrier. It is not $\text{Aut}(\mathbb{Z}_{12})$ -invariant and is treated as a convention-layer sign anchor only.

Lemma 12.2 (Direction-free weights cannot distinguish a sine-axis sign (N517/N518)). *Let $w : \mathbb{Z}_{12} \rightarrow \mathbb{R}$ be reflection-even:*

$$w(x) = w(-x) \quad \text{for all } x \in \mathbb{Z}_{12}.$$

Then for every Fourier sine basis vector s_m ($m = 1, \dots, 5$) one has

$$\langle w, s_m \rangle := \sum_{x \in \mathbb{Z}_{12}} w(x) s_m(x) = 0.$$

In particular, a deterministic sign rule of the form

$$u \mapsto \text{sign}(\langle w, u \rangle) u$$

cannot distinguish u from $-u$ when the exported axis representative lies exactly on a sine axis (e.g. $u = \pm s_1$ on pair1).

Moreover, since $-1 \in \text{Aut}(\mathbb{Z}_{12})$, any $\text{Aut}(\mathbb{Z}_{12})$ -invariant weight profile is necessarily reflection-even. Therefore no direction-free (Aut-invariant) reference-weight family can supply a strict-core sign-distinction scalar on a pure sine axis.

Proof. Because $s_m(-x) = -s_m(x)$, we may pair terms x and $-x$:

$$w(x)s_m(x) + w(-x)s_m(-x) = w(x)s_m(x) - w(-x)s_m(x) = (w(x) - w(-x))s_m(x) = 0.$$

The fixed points $x = 0$ and $x = 6$ satisfy $s_m(0) = s_m(6) = 0$, so they contribute nothing. Hence the total sum vanishes. For the Aut-invariance statement, apply the automorphism $x \mapsto -x$ and invariance of w under $\text{Aut}(\mathbb{Z}_{12})$. $\square$

Exported numeric instance (F647 payload). The exported payload values on the current repo state are:

x	$\text{ord}_{\mathbb{Z}_{12}}(x)$	$w_{\text{ref}}^{\text{unnorm}}(x)$	$w_{\text{break}}(x)$
0	1	$6.250000000000000 \times 10^{-2}$	0
1	12	$3.5527136788005110 \times 10^{-15}$	$-1.7763568394002552 \times 10^{-15}$
2	6	$5.9604644775390720 \times 10^{-8}$	$-5.1619136559035774 \times 10^{-8}$
3	4	$1.5258789062500007 \times 10^{-5}$	$-1.5258789062500007 \times 10^{-5}$
4	3	$2.4414062500000016 \times 10^{-4}$	$-2.1143198334581037 \times 10^{-4}$
5	12	$3.5527136788005110 \times 10^{-15}$	$-1.7763568394002568 \times 10^{-15}$
6	2	$3.9062500000000010 \times 10^{-3}$	$-4.7837765591693490 \times 10^{-19}$
7	12	$3.5527136788005110 \times 10^{-15}$	$1.7763568394002544 \times 10^{-15}$
8	3	$2.4414062500000016 \times 10^{-4}$	$2.1143198334581029 \times 10^{-4}$
9	4	$1.5258789062500007 \times 10^{-5}$	$1.5258789062500007 \times 10^{-5}$
10	6	$5.9604644775390720 \times 10^{-8}$	$5.1619136559035800 \times 10^{-8}$
11	12	$3.5527136788005110 \times 10^{-15}$	$1.7763568394002570 \times 10^{-15}$

The same payload export records the strict scalar sign-anchor witness:

$$\langle w_{\text{break}}, s_1 \rangle \approx -3.9681792259571596 \times 10^{-4},$$

where s_1 is the real Fourier sine basis vector of the pair1 plane (Section 10). This is used downstream only to define deterministic convention-layer sign fixing rules (e.g. T175/F690) and is not promoted into strict-core physics.

12.3 Orientation datum on pair₁ (N546)

Theorem 12.3 (Admissible orientation export (N546)). *Within the current repo state, the exported strict-core source object*

$$S_{\text{sel_int_strict_core_source_object_v1}}$$

exports one admissible orientation datum on the pair1 frame $\text{span}\{c_1, s_1\}$:

$$E_{\text{orient_s_sel_int_source_object_v1}}.$$

Hard limits: *this does not promote a directed physical sign datum and does not by itself discharge strict-core selector closure.*

12.4 Selector-sector and output-sector operators (N549)

Let the selector-sector and output-sector carrier spaces be fixed as two-dimensional real spaces with declared bases:

$$Q_{\text{sel}} := \text{span}\{q_+, q_-\}, \quad Q_{\text{out}} := \text{span}\{o_+, o_-\}.$$

Theorem 12.4 (Seed-v1 selector output operator exists (N549)). *Within the current repo state, the seed-v1 lane exports an admissible strict-core selector output operator*

$$O_{\text{sel}} : Q_{\text{sel}} \rightarrow Q_{\text{out}},$$

packaged as the exported object `0_sel_s_sel_int_source_object_v1`, derived from an admissible reduction operator `R_sel_s_sel_int_source_object_v1`. Hard limits: no strict-core selector closure, no kernel-alone/global QW-2191 discharge, and no ToE closure are claimed.

12.5 Global promotion to C_{v1} -typed operator families (N551/N552)

Theorem 12.5 (Global reduction operator family promotion (N551)). *On the current repo state, the repo exports a global (C_{v1} -typed) selector reduction operator family $R_{\text{sel}}(\text{pair}_m) : \text{pair}_m \rightarrow Q_{\text{sel}}$ promoted from the seed-v1 chain on `pair1` to all charts $\{\text{pair}_1, \dots, \text{pair}_5\}$, with overlap transport consistency kept at projector/section level (up to residual sign gauge). Hard limits: no strict-core selector closure and no global QW-2191 discharge are claimed.*

Theorem 12.6 (Global output operator and chartwise channels (N552)). *On the current repo state, the repo exports a global (C_{v1} -typed) selector output operator $O_{\text{sel}} : Q_{\text{sel}} \rightarrow Q_{\text{out}}$ and the induced chartwise output channels*

$$Y_{\text{sel}}(\text{pair}_m) := O_{\text{sel}} \circ R_{\text{sel}}(\text{pair}_m) \quad (m = 1, \dots, 5),$$

with overlap transport consistency kept explicit up to residual sign gauge. Hard limits: no strict-core selector closure, no kernel-alone/global QW-2191 discharge, and no ToE closure are claimed.

13 Projective closure (T172/T173 projective)

The global projective selector closure is exported as the object

$$\text{SelectorClosure_global_C_v1_projective_strict_v1} \quad (\text{F672}),$$

with theorem-level discharge N680.

In projective scope, the selector state is a ray (projector/span semantics), and the closure observable is an output-space rank-1 projector computed chartwise and then certified as chart-independent:

$$B_{\text{out}}(\text{pair}_m) := Y_{\text{sel}}(\text{pair}_m) A_m Y_{\text{sel}}(\text{pair}_m)^\top \in \mathbb{R}^{2 \times 2},$$

in the fixed output basis (o_+, o_-) . The chartwise output channel $Y_{\text{sel}}(\text{pair}_m) : \text{pair}_m \rightarrow Q_{\text{out}}$ is exported as a promoted seed-v1 chain object (F660) by:

$$Y_{\text{sel}}(\text{pair}_m) := O_{\text{sel}} \circ R_{\text{sel}}(\text{pair}_m),$$

where $R_{\text{sel}}(\text{pair}_m)$ is the exported chartwise selector reduction operator family and $O_{\text{sel}} : Q_{\text{sel}} \rightarrow Q_{\text{out}}$ is the exported seed-v1 selector output operator (both promoted to global C_{v1} -typed objects in the strict program). Residual $\mathbb{Z}_2$ sign handling is tracked at the convention layer and is not promoted to a physical sign datum.

F672 exports that all chartwise $B_{\text{out}}(\text{pair}_m)$ agree (within tolerance) and records the resulting global output projector:

$$B_{\text{out}} = \begin{pmatrix} 1 & -9.149258638450154 \times 10^{-16} \\ -9.149258638450154 \times 10^{-16} & 8.370893363325476 \times 10^{-31} \end{pmatrix} \approx \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

so numerically the global output projector is the pure o_+ ray (machine-level residuals only).

Remark 13.1 (Explicit well-definedness certificate (F672)). *The strict projective closure export F672 includes an explicit well-definedness certificate of chart-independence at projector level. Concretely, for a fixed reference chart (taken as `pair1` in the export), define the max-entry difference*

$$\Delta_m := \max_{a,b \in \{+,-\}} |(B_{\text{out}}(\text{pair}_m) - B_{\text{out}}(\text{pair}_1))_{ab}|.$$

F672 certifies:

$$\max_{m=1,\dots,5} \Delta_m \leq 10^{-12}, \quad \max_m \Delta_m \approx 7.27 \times 10^{-16},$$

and marks the certificate as passed. This is sufficient for projective/ray-level closure scope and does not upgrade to any operator-level groupoid identity (N512 boundary).

This is a *projective* closure statement only: it is insensitive to sign lifts $u_m \mapsto -u_m$ and does not claim any directed/sign-sensitive physical orientation datum.

Theorem 13.1 (Projective strict-core selector closure (N680)). *On the current repo state, there exists one explicit global projective selector closure outcome on C_{v1} (F672) derived from exported strict-core objects, whose closure observable on $Q_{\text{out}} = \text{span}\{o_+, o_-\}$ is a well-defined chartwise-glued rank-1 output projector. Therefore, strict-core selector closure is discharged in projective (ray-level) scope on C_{v1} .*

Hard limits: this does not discharge kernel-alone/global QW-2191, does not promote a directed/sign-sensitive physical orientation datum, and does not claim ToE closure.

14 Directed lifts as gauge/convention layer

Beyond the projective closure object, the repo also exports directed/vector-level representatives and oriented sign-lift data as an explicit *convention layer* (gauge) on top of the projective/ray-level closure. The strict no-false-pass boundary is:

- directed lifts are tracked as gauge/convention data and are not a free strict-core physical sign datum;
- the projective output ray is the invariant content.

Concretely, N692 packages that the directed closure output (with its tracked output sign-lift) induces the same rank-1 output projector as the projective closure output projector of F672. N693 packages the gauge covariance of the directed output sign-lift under chartwise sign relifts.

Theorem 14.1 (Projective/directed closure output-ray invariance (N692)). *Let B_{out} denote the exported projective closure output projector on $Q_{\text{out}} = \text{span}\{o_+, o_-\}$ (F672). For each exported directed closure representative (in its declared scope), let $q_{\text{out}}^{(\text{dir})} \in \mathbb{R}^2$ be the directed output vector (with explicit sign-lift convention). Then the induced rank-1 projector*

$$\Pi(q_{\text{out}}^{(\text{dir})}) := \frac{q_{\text{out}}^{(\text{dir})} q_{\text{out}}^{(\text{dir})\text{T}}}{q_{\text{out}}^{(\text{dir})\text{T}} q_{\text{out}}^{(\text{dir})}}$$

agrees with B_{out} (within the exported numeric tolerance audits). Therefore, the strict global closure outcome on Q_{out} is projectively invariant across the exported directed lifts.

Hard limits: this does not promote a directed/sign-sensitive physical orientation datum in strict core and does not discharge kernel-alone/global QW-2191.

Theorem 14.2 (Chartwise $\mathbb{Z}_2$ gauge covariance of output sign-lift (N693)). *Let $s_{\text{out}}^{\text{prem}}(\text{pair}_m) \in \{\pm 1\}$ denote the exported output sign-lift for the premise-based directed closure on chart pair_m , and let $t(\text{pair}_m) \in \{\pm 1\}$ be the exported chartwise sign relift mapping the premise-based directed representative to the exported sign-fixed directed representative. Then the sign-fixed directed closure output sign-lift satisfies, for $m = 1, \dots, 5$,*

$$s_{\text{out}}^{\text{fix}}(\text{pair}_m) = t(\text{pair}_m) s_{\text{out}}^{\text{prem}}(\text{pair}_m).$$

Therefore, the directed closure output sign-lift is a tracked $\mathbb{Z}_2$ gauge datum and does not upgrade to a strict physical sign.

Hard limits: *no directed physical orientation datum is claimed; no kernel-alone/global QW-2191 discharge is claimed.*

On the current repo state, the directed layer is further stabilized by explicit convention-scoped sign data:

- oriented overlap-edge sign-lifts (T174: F688/P688/N688), with gauge-equivalence packaged (N689),
- deterministic chart sign fixing (0-cochain) from strict-core payload weights (T175: F690/P690/N690),
- a corresponding sign-fixed directed closure export (F692) that still induces the same projective output ray (by N692).

All of these remain *convention-layer* artifacts and do not upgrade any directed sign into strict-core physics.

Explicit $\mathbb{Z}_2$ cochain language for sign lifts (T174/T175)

Let the chart set be $\mathcal{C} = \{\text{pair}_1, \dots, \text{pair}_5\}$ and let $\mathcal{E}$ be the set of 10 overlap edges (the complete graph on $\mathcal{C}$). For each edge $(i, j) \in \mathcal{E}$, the axis-only transition representative exports an operator $O_{ij}^{(\text{axis})}$ (rotation $\alpha_{ij} \bmod \pi$).

Definition 14.1 (Edge sign lift (convention layer)). *An oriented edge sign lift is a $\mathbb{Z}_2$ -valued 1-cochain $s : \mathcal{E} \rightarrow \{\pm 1\}$ defining oriented transitions:*

$$O_{ij}^{(\text{oriented})} := s_{ij} O_{ij}^{(\text{axis})}.$$

Definition 14.2 (Chart sign relift (gauge)). *A chartwise sign relift is a $\mathbb{Z}_2$ -valued 0-cochain $t : \mathcal{C} \rightarrow \{\pm 1\}$ acting on directed representatives by:*

$$u_i \mapsto t_i u_i.$$

It induces the gauge action on edge signs:

$$s_{ij} \mapsto s'_{ij} := t_j s_{ij} t_i.$$

In this language, T174 exports explicit oriented edge sign-lifts s_{ij} that remove sign flips for declared directed representatives (full-edge coherence), and T175 exports one explicit deterministic chart sign-fix t_i derived from strict-core payload weights (Section 12.2). Both remain below any physical sign datum claim.

Exported instance (Release 7). On the current repo state, the deterministic chart sign-fixing 0-cochain exported by F690 is:

$$(t_{\text{pair}1}, t_{\text{pair}2}, t_{\text{pair}3}, t_{\text{pair}4}, t_{\text{pair}5}) = (-1, -1, -1, +1, -1).$$

Anchored to that sign-fixed directed representative, the oriented edge sign-lift exported by F691 flips exactly two overlap edges:

$$s_{\text{pair}2 \rightarrow \text{pair}4} = -1, \quad s_{\text{pair}1 \rightarrow \text{pair}5} = -1,$$

with all remaining overlap edges having $s_{ij} = +1$ (convention scope; no physical sign claim).

15 Open boundary: kernel-alone/global QW-2191 discharge

Theorem 15.1 (Post-T172 strict frontier boundary (N679)). *After the repo exports global selector closure observables on C_{v1} in both:*

- *projective scope (F672, packaged by N672/N673/N674), and*
- *directed premise-tracked scope (F677, packaged by N677/N678), with the raw output obstruction boundary recorded by N675,*

the repo packages the remaining strict frontier as a theorem-level boundary statement (N679):

1. *projective strict-core selector closure is discharged (N680),*
2. *but kernel-alone/global QW-2191 discharge remains explicitly unclaimed,*
3. *and no directed/sign-sensitive physical orientation datum is promoted into strict core beyond premise-tracked sign-lift conventions.*

Hard limits: **N679** does not claim kernel-alone/global QW-2191 discharge, any directed physical sign datum in strict core, or ToE closure.

QW-2191 remains a real strict-core uniqueness obstruction: kernel-alone translation-invariant data leaves an $O(2)$ basis freedom inside each degenerate Fourier pair plane.

The exported global closure in this document is therefore deliberately in *projective* scope (ray/projector semantics), and the repo keeps kernel-alone/global QW-2191 discharge explicitly open. Any future upgrade to strict-core selector closure or a directed physical sign datum must be exported and proven as a separate theorem-level object.

16 Operational physics proxies from projective closure

17 Mass-spectrum proxy (P694)

Let $H_\psi \in \mathbb{R}^{12 \times 12}$ be the exported diagonal/local Psi-sector Hessian value instantiation (F459). We use its symmetrization:

$$H_\psi^{(s)} := \frac{1}{2}(H_\psi + H_\psi^\top).$$

For each Fourier pair plane pair_m ($m = 1, \dots, 5$), the exported global projective selector closure (F672) provides a ray $u_m \in \mathbb{R}^{12}$ (defined only up to sign). Define the quadratic proxy:

$$m_m^2 := \frac{u_m^\top H_\psi^{(s)} u_m}{u_m^\top u_m}.$$

This is invariant under the sign gauge $u_m \mapsto -u_m$.

Probe **P694** computes these “mass-spectrum proxy” values on the current repo state (no Standard Model identification claimed):

pair	m^2 proxy (Rayleigh on $H_\psi^{(s)}$)
pair3	0.6687166508753918
pair5	1.3892705031823167
pair4	1.4474315384116325
pair2	2.0699508733172283
pair1	2.9337556210543343

18 Selector-aligned channel spectrum proxy (P696)

Probe P696 refines P694 into a 12-channel spectrum proxy by constructing a deterministic selector-aligned basis and inspecting $H_\psi^{(s)}$ in that basis.

On $\mathbb{Z}_{12}$, define the standard real orthonormal Fourier vectors:

$$e_0 := \frac{1}{\sqrt{12}}(1, \dots, 1), \quad e_6 := \frac{1}{\sqrt{12}}(1, -1, 1, -1, \dots),$$

and for each $m = 1, \dots, 5$ the real pair-plane basis (c_m, s_m) with scale $\sqrt{2/12}$.

For each pair _{m} , F672 exports coordinates (u_c, u_s) such that (in the declared Fourier convention)

$$u_m = u_c c_m + u_s s_m, \quad u_m^\perp = (-u_s) c_m + (u_c) s_m,$$

so that $u_m^\perp$ is a deterministic orthogonal complement ray in the same plane (used only at ray/projector level).

Define the selector-aligned 12-channel basis:

$$B := (e_0, u_1, u_1^\perp, u_2, u_2^\perp, \dots, u_5, u_5^\perp, e_6),$$

and the induced matrix:

$$M := B^\top H_\psi^{(s)} B.$$

The diagonal entries of M are recorded as a simple operational “channel proxy spectrum”. Mixing is audited explicitly by inspecting the off-diagonal norm.

On the current repo state, P696 exports the sorted diagonal channel proxies (no Standard Model identification claimed):

channel	m^2 proxy
pair3:u	0.6687166508753918
pair5:u	1.3892705031823165
pair4:u	1.4474315384116314
pair2:u	2.069950873317227
e6	2.41906833577034
pair1:u	2.9337556210543316
pair5:u_perp	3.5360177064927694
pair4:u_perp	3.675931516755229
pair2:u_perp	4.2984508516608315
e0	4.761250376916641
pair3:u_perp	4.930970379004997
pair1:u_perp	5.08050282436479

The mixing audit is part of the exported result: on the current repo state, P696 reports an off-diagonal-to-diagonal Frobenius ratio of 1.191284779529568 and a maximum off-block coupling Frobenius norm of 3.4009571444250914, so *no diagonalization claim* is made.

19 Basis-invariant Hessian eigen-spectrum proxy (F704)

The most basis-stable quadratic proxy spectrum is the invariant eigenvalue set of the symmetric Hessian $H_\psi^{(s)}$ itself. If we change coordinates by an orthogonal basis transformation $R \in O(12)$, then

$$H_\psi^{(s)} \mapsto R H_\psi^{(s)} R^\top,$$

so the eigenvalues $\lambda_i(H_\psi^{(s)})$ are unchanged. This avoids any basis-dependent “channel” choices and cleanly separates the invariant spectrum from mixing/assignment questions (which remain open at the physics-interpretation layer).

F704 exports the invariant eigenvalue spectrum on the current repo state (dimensionless m^2 proxies in strict normalization; no physical unit claim):

mode index	$\lambda_i(H_\psi^{(s)})$ (proxy m^2)
0	0.08813142775481989
1	0.11195105316362809
2	0.23796767585569786
3	0.7641952808835573
4	0.7733323123191046
5	1.2068935563568985
6	1.7060617210611557
7	1.8861137936815946
8	2.290146094544166
9	2.989559197836824
10	12.356024722005182
11	12.800940342343853

20 What is and is not claimed physically

- **Claimed (operational):** the repo exports a strict projective selector closure and an OS bundle sufficient to compute the proxy spectra P694/P696 and downstream observer-limit artifacts in a projective (ray-level) gauge-safe scope.
- **Not claimed:** that any proxy number is already identified with a Standard Model mass or coupling.
- **Not claimed:** that the selector-aligned channel basis diagonalizes H_ψ ; mixing is audited and explicitly non-negligible (P696).
- **Not claimed:** kernel-alone/global QW-2191 discharge, directed sign-sensitive physical orientation in strict core, or ToE closure.
- **Host matching discipline:** any attempted identification of proxy values with external physics data is necessarily non-strict and must carry an explicit external dataset + mapping policy; see P702 (channel proxy) and P704 (eigen-spectrum proxy). The current repo freezes this non-strict layer by separating:
 - the external dataset (targets only): `external_data/sm_mass_targets_v1.json`,
 - the external versioned policy (assignment + allowed DOF + optional thresholds): `external_data/sm_ho`

where the current policy explicitly allows *only* a single global scale and sets `pass_criteria=null` (diagnostic metrics only; no pass/FAIL claim).

- **Non-strict diagnostic (current):** under the simplest one-scale centered-log assignment policy against an explicit PDG2024 mass target dataset, P702 computes $\text{RMSE}(\log m^2) \approx 8.10$ and $\max|\Delta m/m| \approx 7.90 \times 10^2$, while P704 computes $\text{RMSE}(\log m^2) \approx 7.23$ and $\max|\Delta m/m| \approx 4.35 \times 10^2$, so *no* “Standard Model match” claim is made at this stage.

21 Emergent observer pipeline (projective scope)

22 Observer-limit readout from projective closure output (P697)

From the exported global projective closure output projector B_{out} in the (o_+, o_-) basis (F672), P697 exports a minimal observer-limit readout consisting of gauge-safe invariants of the 2×2 projector. Write

$$B_{\text{out}} = \begin{pmatrix} p_+ & c \\ c & p_- \end{pmatrix}.$$

Define:

$$y_{\text{bias}} := \frac{1}{2}(p_+ - p_-), \quad y_{\text{total}} := \frac{1}{2}(p_+ + p_-), \quad z_{\text{commit}} := p_+, \quad z_{\text{residual}} := p_-.$$

On the current repo state, P697 reports $p_+ = 1$ and $p_- \approx 8.37 \times 10^{-31}$, so the output projector is numerically the pure o_+ ray, and the readout invariants are $(y_{\text{bias}}, y_{\text{total}}, z_{\text{commit}}, z_{\text{residual}}) \approx (1/2, 1/2, 1, 0)$.

23 Projective emergent-observer chain computability from closure output (P699/N699)

In the declared projective/gauge-safe downstream scope, we represent a ray by a rank-1 (possibly unnormalized) symmetric positive semidefinite matrix P (a projector candidate). For a linear map M , the induced pushforward is:

$$P \mapsto M P M^T.$$

Let Q_{out} be the exported global projective selector-closure output projector in the (o_+, o_-) basis ('F672').

Define the downstream operator chain by the already exported matrices ('F92..F99'):

$$C = \begin{pmatrix} \frac{1}{2} & -\frac{1}{2} \\ \frac{1}{2} & \frac{1}{2} \end{pmatrix}, \quad L = \begin{pmatrix} 1 & 1 \\ -1 & 1 \end{pmatrix}, \quad G = I_2, \quad H = I_2, \quad J = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix},$$

$$K = \begin{pmatrix} 1 & 0 \end{pmatrix}, \quad M = (1), \quad N = (1).$$

Starting from Q_{out} , define the induced chain:

$$Y = C Q_{\text{out}} C^T, \quad Z = L Y L^T, \quad W = G Z G^T, \quad X = H W H^T, \quad U = J X J^T, \quad F = K U K^T, \quad P = M F M^T, \quad C_{\text{cl}} = N P N^T.$$

'P699' computes this chain on the current repo state and audits that the declared residual components vanish (within tolerance) while commitment remains positive. 'N699' packages the computability statement with explicit hard limits (no kernel-alone/global 'QW-2191' discharge; no ToE closure; no actual emergent observer closure).

Theorem 23.1 (Projective emergent-observer chain computability (N699)). *On the current repo state, the repo exports a strict projective downstream chain computability witness: starting from the global projective selector closure output projector (F672), the induced projector pushforward semantics*

$$P \mapsto M P M^T$$

along the exported operator chain (F92..F99) produces a fully computable chain of projectors up to the declared closure-candidate level (P699). This is a projective/ray-level (gauge-safe) computability result only.

Hard limits: *no kernel-alone/global QW-2191 discharge, no ToE closure, and no actual emergent observer closure are claimed.*

24 Operational OS support packet (v1) (F698/P698/N698)

The v1 OS support packet (F698, audited by P698, packaged by N698) collects the minimum strict projective bundle sufficient to compute:

- global projective selector closure on C_{v1} (N680/F672),
- directed-vs-projective output-ray invariance and sign-lift gauge covariance (N692, N693),
- first quadratic proxy spectra (P694, P696),
- observer-limit readout (P697).

The mass-proxy meaning is frozen separately as a scope-limited theorem (N703). A basis-invariant Hessian eigen-spectrum mass observable is exported separately as F704 and included in the v3 OS support packet (F705).

25 Operational OS support packet (v2) (F700/P700/N700)

The strict projective operational ToE “OS” support packet is exported in v2 form:

$$\Lambda_{\text{OS}}^{(\text{proj})} = (\text{N680}, \text{N692}, \text{N693}, \text{P694}, \text{P696}, \text{P697}, \text{N699}),$$

as the object `Lambda_strict_projective_operational_toe_os_support_v2` (F700), audited by P700 and packaged as a theorem by N700. All hard limits remain explicit: no kernel-alone/global QW-2191 discharge, no directed sign-sensitive physical orientation claim in strict core, and no ToE closure.

26 Operational OS support packet (v3) (F705/P705/N705)

The strict projective operational ToE “OS” support packet is exported in v3 form by explicitly including the basis-invariant Hessian eigen-spectrum mass observable F704:

$$\Lambda_{\text{OS}}^{(\text{proj})}(\text{v3}) = (\text{N680}, \text{N692}, \text{N693}, \text{P694}, \text{P696}, \text{F704}, \text{P697}, \text{N699}),$$

as the object `Lambda_strict_projective_operational_toe_os_support_v3` (F705), audited by P705 and packaged as a theorem by N705. All hard limits remain unchanged.

Theorem 26.1 (Strict projective operational OS support v3 (N705)). *On the current repo state, the repo exports one strict projective operational OS support packet v3:*

$$\Lambda_{\text{OS}}^{(\text{proj})}(\text{v3}) = (\text{N680}, \text{N692}, \text{N693}, \text{P694}, \text{P696}, \text{F704}, \text{P697}, \text{N699}),$$

exported as F705, audited by P705, and packaged at theorem level by N705. This is an operational packaging result only: it adds an explicit tracked basis-invariant mass observable (F704) without strengthening any hard-limit claims.

Hard limits: no kernel-alone/global QW-2191 discharge, no directed physical sign datum, no Standard Model host matching, no ToE closure, and no actual emergent observer closure are claimed.

Proposition 26.2 (Release-7 strict projective OS closure dashboard (P706)). *On the current repo state, the repo exports a single consolidated dashboard probe (P706) that checks the strict release-7 operational closure bundle by verifying: N705 (OS support v3), N701 (Operational ToE closure in strict projective OS scope), N703 (mass-proxy meaning discipline), and F704 (basis-invariant mass observable export). The current P706 summary status is PASS_RELEASE_7_STRICT_PROJECTIVE_OPERATIONAL_OS_CLOSURE_DASH*

27 Open target: actual emergent observer closure (future)

The current strict projective pipeline proves *computability* of a downstream observer-facing chain, not the existence/uniqueness of an actual emergent observer. Any future “actual observer closure” claim must be exported as a separate theorem-level artifact with explicit premises and hard limits.

28 Hard limits and open problems

29 Kernel-alone/global QW-2191 discharge

QW-2191 remains open at kernel-alone/global strict-core level. All global closure statements used in the OS bundle are in *projective* scope (rays/projectors) and do not assert strict-core uniqueness of directed selector representatives.

30 Directed physical sign datum

Directed/vector-level sign and orientation data are tracked as a gauge/convention layer. No strict-core theorem in Release 7 upgrades directed sign into a physical datum “for free”.

31 Host matching and Standard Model identification

Any comparison of proxy spectra against external experimental data is non-strict and must carry:

- an explicit external dataset, and
- an explicit identification/mapping policy.

The repo exports such a harness (P702) and keeps it outside strict scope; current one-scale metrics do not support any “Standard Model match” claim.

32 What Release 7 declares as closed vs open

Closed (scope-limited, strict projective OS):

- projective global selector closure on C_{v1} with chart-independent output projector (N680/F672),
- operational OS support packets (v1–v3) and OS computability pipeline from closure output through observer-limit readout and downstream chain (N698, N700, N705, N701),
- strict basis-invariant Hessian eigen-spectrum mass observable (F704),
- residual sign $u_m \mapsto -u_m$ is gauge-irrelevant for the Release-7 OS observables actually used downstream (audited by P709, packaged by N706),
- a disciplined internal meaning of the quadratic “mass proxy” layer (N703).
- release-7 OS closure dashboard readiness (P706) (probe-level).

Open (explicit hard limits):

- kernel-alone/global QW-2191 discharge,
- directed sign-sensitive physical orientation datum,
- Standard Model host matching as a strict claim,
- actual emergent observer closure.

33 Next steps (roadmap, no false pass)

This section records the next concrete moves under strict scientific discipline, separating strict targets from explicitly non-strict policy work.

Priority 1 (strict): T173 — residual sign / kernel-alone QW-2191 frontier

The strict frontier remains: (i) kernel-alone/global discharge of QW-2191, and (ii) any directed/sign-sensitive physical orientation datum in strict core beyond projective/ray semantics. Release 7 exports convention-layer directed representatives (T174/T175), but these do not upgrade sign into strict physics.

The next honest strict move under T173 is therefore ordered as:

- (a) first, export a genuinely strict internal selector source/ingredient that determines the residual sign in a chart-independent way (no hidden convention slots), and
- (b) only if that provider attack stalls, explicitly freeze the directed sign as a gauge/convention layer and keep physics claims restricted to projective/basis-invariant observables already audited downstream.

For option (b), Release 7 now exports an explicit machine audit: P709 enumerates all 2^5 residual sign patterns on $\{\text{pair1}, \dots, \text{pair5}\}$ and verifies that the concrete OS numeric outputs used downstream (quadratic proxy layer P694/P696 and the basis-invariant mass observable F704) are unchanged (within tolerance). The corresponding theorem-level package is N706.

The older methodology is also now audited directly rather than guessed at. The new audit P711 (packaged by N707) shows the strongest honest reading: the `sigma_int`/topological lane survives as a genuine strict local ingredient class, reaching bridge discharge in declared R1 scope, but it still does not provide the missing *global* chartwise sign-coherence provider on the full C_v1 atlas. So it is a clue and a reusable ingredient, not yet a full T173 discharge.

That missing object is now frozen as an explicit target class under the next honest continuation label T176: `GlobalDirectedSignCoherenceProvider_global_C_v1_strict_core_v1`. The new nonexport audit P712 (packaged by N708) confirms that no currently exported strict/global object already satisfies this target.

The next honest narrowing step is also now explicit. The new audit P713 (packaged by N709) shows that the surviving `w_break` + oriented-transport route does *not* globalize to all roots on the exported five-chart lane: only `pair1` and `pair5` provide nonzero root anchors, and those supported roots recover the same convention-layer section. So the surviving candidate is narrower than a full global provider and still does not discharge T176.

The new audit P714 (packaged by N710) then explains why this surviving corridor stops exactly there on current exports: the present `w_break` payload is an *odd* linear weight on Z_12, while the currently exported representatives on `pair2`, `pair3`, `pair4` are even/cosine-axis states. Hence the anchor

$$\sum_x w_{\text{break}}(x) u_m(x)$$

vanishes on those charts by parity and survives only on the odd-axis charts `pair1`, `pair5`. So the next honest strict provider attack must either add an even-or-mixed parity anchor component or leave the single-linear-weight class altogether.

The next audit P715 (packaged by N711) then tests exactly that minimal parity completion using only already-exported strict-core payloads: `w_break` together with the exported even weight `w_ref_unnormalized`. The result is sharper but still scope-limited: all five roots become anchorable, and the recovered rooted sections agree projectively across all roots, yet the exact directed branch still splits into two global sign orbits. Relative to the reference orbit, `pair4` yields the globally negated section. Therefore adding an even anchor component is not sufficient by itself; the remaining strict-provider gap is the final global Z2 orbit split.

The next audit P716 (packaged by N712) localizes that exact split one step further. On current exports, the residual exact orbit split is fully explained by one explicit chart-polarity feature: **pair4** is the unique negative cosine-axis chart, and under the even fallback anchor **w_ref_unnormalized** its root scalar is negative, whereas the corresponding cosine-anchor scalars for **pair2** and **pair3** are positive. So the current failure is no longer diffuse. The remaining strict-provider gap is an exact global branch-fixing rule across chart polarity, not root support and not projective orbit recovery.

The next bridge audit P717 (packaged by N713) then asks the next honest downstream question: whether this localized exact split still matters for the concrete Release-7 OS observables. The answer is now sharper. Because P709/N706 already proves sign-gauge irrelevance over the full residual sign family for the downstream observables actually used in Release 7, the present exact **pair4**-induced branch split is already operationally inert for P694, P696, and F704. So the remaining gap is a strict-core provider problem only, not a Release-7 OS observables problem.

The next provider-side audit P718 (packaged by N714) then tests the narrowest honest class that remains before one has to leave the current exported linear anchor family altogether: any single mixed linear weight in $\text{span}\{w_{\text{break}}, w_{\text{ref,unnormalized}}\}$. This result is again negative but sharper. Every noncritical sector of that span still supports all roots and recovers at most a projective orbit, but no sector upgrades that orbit into one exact global directed section. Some sectors reproduce the current **pair4**-only negated branch; other sectors worsen the split to **pair2/pair3**. So the next strict provider attack must leave the present single-linear span class.

The next nonlinear audit P719 (packaged by N715) then tests the nearest untuned extension of that same current carrier: odd polynomials in the two exported readouts ($\langle w_{\text{break}}, u \rangle, \langle w_{\text{ref}}, u \rangle$) of total degree ≤ 3 , with coefficient alphabet $\{-1, 0, 1\}$. This too is negative, but more informative. Across all 728 nonzero candidates in that scanned class there are 0 exact all-root directed sections, 576 projective-only candidates, and no new negated-root pattern beyond **pair4** or **pair2/pair3**. So the next strict provider attack must either raise complexity in an explicit non-hidden way or leave the current two-readout carrier altogether. From the current Release-7 physics perspective, the preferred continuation is the latter: a new *physically interpretable* observable/provider class rather than further coefficient tuning.

The next observer-facing audit P720 (packaged by N716) then tests the nearest static class on the actually exported output carrier $Q_{\text{out}} = \text{span}\{o_+, o_-\}$: signed projections of the rooted unsigned output vectors onto all nonzero directions with coefficients in $\{-1, 0, 1\}$. This is again negative, but it sharpens the physics-facing picture. There are 0 exact all-root directed sections, 6 projective-only candidates, and every supported direction reproduces only the coarser negated-root pattern **pair4/pair5**; the pure o_- channel is unsupported because the current unsigned rooted outputs already lie almost entirely on o_+ . So even the nearest static observer-facing readout class is too coarse to fix T176, and the next physics-facing provider attack should move toward a more dynamical or flux-like observable rather than further static output-axis tuning.

The next source-topology audit P721 (packaged by N717) then tests the most natural already-exported bridge toward physics: the lane of actual nonzero flow, barrier-protected sign, selector witness, basis-independent promotion, and quotient-safe QW-2191 resolution. This result is mixed but still negative at exact-provider level. Those physics-facing strict ingredients are real and already exported, but the lane remains explicitly downstream-only, τ_{src} is still not identified with the current strict selector carrier, and the best exported object remains a basis-free quotient class only rather than one exact chartwise directed branch on the full atlas. So the next honest physics-facing provider attack should aim at a chart-sensitive transported flux/current-like section rather than another static quotient or axis class.

The next bridge audit P722 (packaged by N718) then sharpens that gap further. The repo already has chart-sensitive transport infrastructure on the **pair1..pair5** atlas, and it already has source-topology physical ingredients. But the current source-topology promotion step explicitly forgets chart basis labels, the current quotient-safe step explicitly quotients chart labels away, and the declared-scope theorem packet keeps the lane downstream-only. So the current source-topology lane is now localized

more precisely as *physics-facing but chart-blind*. The next missing bridge target is therefore explicit: a chart-sensitive transported flux/current-like section on the full atlas.

The next sharpening audit P723 (packaged by N719) then localizes the bridge one level lower again. The source-topology lane already supplies positive sign, positive flow, and selector polarity, and the atlas already supplies full chart transport. But the source-side selector realization still lives in the preLM basis u_T, u_L , the source packets remain pair-chart agnostic, and the atlas exports no distinguished root chart. So the next missing bridge target is now even narrower: a source-topology-to-atlas chart-seed selection bridge telling where the physical source enters the atlas before transport.

The next audit P724 (packaged by N720) then gives the first constructive physical reduction below that bridge. Current positive source polarity already excludes the unique negative atlas-entry branch **pair4** inside the strongest current all-root transported-family candidate. But this still leaves the positive corridor **pair1, pair2, pair3, pair5** unresolved. So the bridge gap is reduced, not closed.

The next audit P725 (packaged by N721) then sharpens that surviving positive corridor further. Current exports already reduce the problem to two remaining positive sublanes: odd-anchor **pair1, pair5** and even-fallback **pair2, pair3**. But the current source-side packets still export no source-to-lane rule selecting between them. So the next missing bridge target is narrower again: a positive-corridor odd/even lane-selection bridge.

The next audit P726 (packaged by N722) then re-reads that same remaining split geometrically on the exported five-chart atlas. The atlas still carries the canonical backbone **pair1-pair2-pair3-pair4-pair5**, so **pair1, pair5** are the outer-edge charts while **pair2, pair3** are the surviving positive interior charts. But the current source-side packets still export no source-to-atlas-position rule selecting between those two chart classes. So the next missing bridge target is narrower again: a positive-corridor outer/interior chart-selection bridge.

The next audit P727 (packaged by N723) then localizes that same surviving positive corridor directly relative to the excluded negative branch itself. Once **pair4** is excluded, the positive charts adjacent to that excluded branch are exactly **pair3, pair5**, while the positive charts shielded from it are **pair1, pair2**. But the current source-side packets still export no source-side rule selecting between those two chart classes. So the next missing bridge target is narrower again: a positive-corridor excluded-negative-boundary adjacency chart-selection bridge.

The next audit P728 (packaged by N724) then injects the first actually exported source-side carrier below that geometric split. The strict residual-datum support carrier F301 already lives on **tau_src_candidate_v1** and carries **pair_index_set={pair1, pair2}**, exactly matching the positive boundary-shielded sublane localized by P727. So the surviving positive corridor is now reduced in a genuinely source-side way to **pair1, pair2**. But F301 remains selector-neutral, **tau_src** is still not identified with the current selector carrier, and the carrier remains exactly symmetric between **pair1** and **pair2**. So the next missing bridge target is narrower again: a residual-datum source-side **pair1/pair2** chart-selection bridge.

The next audit P729 (packaged by N725) then localizes that remaining **pair1/pair2** ambiguity one level further, without inventing any new source object. On current exports, F301 already identifies the two surviving branches as opposite orbit-index realizations on the same residual-datum carrier: **pair1** carries the branch δ_k , while **pair2** carries the branch δ_{-k} . The exported **pair1 $\leftrightarrow$ pair2** lane transport and projector glue still only relate those two branches faithfully; they do not select one of them. So the next missing bridge target is narrower again: a residual-datum **pair1/pair2** orbit-direction selection bridge.

The next audit P730 (packaged by N726) then tests the strongest already-exported strict-side selector ingredient against that same surviving split: the direction-free Shannon ord-reference lane. This result is again negative but sharper: that lane really does export genuine slot-free **pair1/pair2** $0(2)\rightarrow Z2$ cuts, yet **ord_Z12/r_ord** is inversion-invariant and the surviving F301 branches are exactly the inversion pair δ_k / δ_{-k} with identical carrier weights. So the induced source-side **ord** expectation and cross-entropy scores are equal on both branches, and the next missing bridge target narrows again: a direction-free Shannon residual-datum **pair1/pair2** orbit-direction selection bridge.

The next audit P731 (packaged by N727) then tests the strongest already-exported inversion-sensitive witness-side ingredient against that same surviving split: the current reflection-breaking payload **w_break** from F647. This result is constructive but still boundary-only: on the surviving F301 carrier, **w_break** already gives opposite nonzero scores to δ_k and δ_{-k} , so the split is no longer merely invisible on current exports. But this remains only a witness-side separation: F647 is still below admissible **S_sel_int**, F301 remains selector-neutral, and P684 keeps **w_break** only at rooted sign-source / convention strength. So the next missing bridge target narrows again: a typed promotion bridge from that witness-side separation to one selected strict source-side branch.

The next audit P732 (packaged by N728) then tests the strongest already-exported downstream object that could be mistaken for that missing promotion: the current pair1-rooted directed atlas convention state F684. This result is again negative but sharper: the object exists, but it is rooted on the chosen chart **pair1**, remains explicitly convention-scoped, and carries the same transported sign on **pair1** and **pair2** even though P731 already separated those source-side branches by opposite witness-score signs. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the present pair1-rooted convention gauge.

The next audit P733 (packaged by N729) then tests the strongest already-exported full convention-layer transport below that same frontier: the current rooted oriented edge-sign lift and sign-fixed convention-layer lift on the **pair1**→**pair2** edge. This result is again negative but sharper: the current convention layer is real and coherent, yet under every currently exported lift the **pair1**→**pair2** transport sign stays +1 with no sign flip even though P731 already separated the underlying source-side branches by opposite witness-score signs. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the current positive convention-layer pair12 transport either.

The next audit P734 (packaged by N730) then tests the strongest already-exported source-side theorem lane below that same frontier: the declared-scope source-topology selector theorem built from F148/F149/F150. This result is again negative but sharper: that theorem lane is real and quotient-safe, yet it still forgets chart labels, keeps only one basis-free distinguished quotient class, and does not claim raw-theta uniqueness. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the current basis-free quotient-only source-side theorem lane either.

The next audit P735 (packaged by N731) then tests the strongest already-exported local source-side scalar bind lane below that same frontier: the positive barrier-sign witness, the positive nonzero-flow witness, the source-derived contraction map $(\mathbf{a}, \mathbf{b}) = (\cos(\phi), \cos(\phi))$, the current provider-object carrier layer, and the surviving F301 support carrier. This result is again negative but sharper: those local scalar bind data are real and already bind into the pair1/pair2 carrier lane, yet they still factor only through one shared positive $\cos(\phi)$ family, keep $\mathbf{a}=\mathbf{b}$, and inherit identical pair1/pair2 weight profiles. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the current shared local source-side scalar bind lane either.

The next audit P736 (packaged by N732) then tests the strongest already-exported local non-scalar provider-operator lane below that same frontier: the pair-indexed opposite-shift carrier operators themselves on the surviving F301 support carrier. This result is again negative but sharper: that lane already realizes both surviving pair1/pair2 branches from the common seed **delta_0**, yet it still does so with equal source-derived amplitudes $(\mathbf{a}, \mathbf{b}) = (\cos(\phi), \cos(\phi))$, exact inversion-paired carrier data, and explicit selector-neutral status. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the current symmetric local provider-operator shift-direction lane either.

The next audit P737 (packaged by N733) then tests the strongest already-exported local pair1/pair2 atlas/glue lane below that same frontier: the explicit chart-transport object **0_12**, the glued projector-operator section, and the lane-scoped pair1/pair2 atlas stub with overlap declaration. This result is again negative but sharper: that lane is real, yet it remains entirely projector-level and sign-gauge-safe, so it still does not descend the opposite P731 pair1/pair2 witness split as one typed branch distinction. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not

collapse into the current local projector-atlas glue lane either.

The next audit P738 (packaged by N734) then tests the strongest already-exported global projective selector transition/state lane below that same frontier: the global selector transition object F469, the global projective selector state object F470, and the global gluing-consistency audit P474. This result is again negative but sharper: that lane is real and globally glued on C_v1, yet it remains strictly projective/ray-level and sign-gauge-safe, so it still does not descend the opposite P731 pair1/pair2 witness split as one typed branch distinction. So the next missing bridge target narrows again: a typed descent/promotion bridge that does not collapse into the current global projective selector state lane either.

The next audit P739 (packaged by N735) then tests the strongest already-exported global directed selector-state lane below that same frontier: the premise-based directed selector state F474/N524 together with the current directed/projective state audits H37/H38. This result is again negative but sharper: that lane is real and sign-sensitive, yet it remains explicitly premise-based via T164 and descends back to the projective quotient layer, so it still does not upgrade the opposite P731 pair1/pair2 witness split into one strict-core branch distinction. So the next missing bridge target narrows again: a typed strict-core bridge that does not collapse into the current premise-based global directed selector state lane either.

The next audit P740 (packaged by N736) then tests the strongest already-exported global sign-fixed directed closure lane below that same frontier: the sign-fixed directed representative F690/N690, the oriented transition edge sign-lift F691/N691, and the sign-fixed directed closure object F692 together with output-ray invariance N692 and gauge covariance N693. This result is again negative but sharper: that lane is real and globally glued on C_v1, yet it glues only after one explicit output sign-lift, remains entirely in `strict_convention`/gauge scope, collapses back to the same projective output ray, and keeps gauge-covariant output sign-lift data, so it still does not upgrade the opposite P731 pair1/pair2 witness split into one strict-core branch distinction. So the next missing bridge target narrows again: a typed strict-core bridge that does not collapse into the current global sign-fixed directed closure lane either.

The next audit P741 (packaged by N737) then tests the strongest already-exported actual source-side selector witness lane before basis-free quotienting: the actual source-topology selector witness packet F147 on `tau_src_candidate_v1` against the surviving F301 pair1/pair2 carrier. This result is again negative but sharper: that lane is real and already exports a positive signed selector split, yet it remains chart-bound, preobserver/downstream-only, and lives only in the preLM basis `u_T, u_L`, so it still does not promote the opposite P731 pair1/pair2 witness split into one selected source-side branch. So the next missing bridge target narrows again: a typed source-side bridge that does not collapse into the current actual preLM selector witness lane either.

Priority 2 (explicitly non-strict policy): physical units (proxy $\rightarrow$ GeV)

Any unit assignment and any comparison to external experimental data must remain non-strict unless a strict unit map is derived. The repo already exports the required dataset-vs-policy separation:

- dataset: `external_data/sm_mass_targets_v1.json` (units = GeV),
- policy: `external_data/sm_host_matching_policy_v1.json` (one global scale; `pass_criteria=null`).

Additionally, Release 7 exports a dedicated non-strict *unit calibration* policy and a corresponding calibration-map probe for the basis-invariant proxy spectrum F704:

- policy: `external_data/proxy_to_gev_calibration_policy_v1.json` (`pass_criteria=null`),
- calibration map probe: P710 (`fundamental_action_reconstruction/p710_current_nonstrict_proxy_to_`

The v1 calibration uses a single global scale parameter defined by matching geometric means of m^2 (assignment-free), and exports only a proxy $\rightarrow$ GeV interface (no identification/match claim).

Priority 3 (theory extension): “solitons = particles”

To make the slogan scientific, the program must export an explicit nonlinear EOM/Lagrangian in which soliton solutions are defined, a stability criterion, and a map from soliton moduli to the observer-limit proxy layer.

Priority 4 (falsifiable outputs): operational tests under frozen artifacts

The nearest falsifiable layer consistent with current strict scope is the basis-invariant proxy spectra (P694, P696, F704) and the non-strict external host-matching harnesses (P702, P704) producing diagnostics only (no pass claim).

A Export map: key artifacts

A.1 Strict-core selector closure

- Global atlas + transition objects on C_{v1} : F469 (N515); exports `SelectorAtlas_global_C_v1_strict_v1` and `SelectorTransition_global_C_v1_strict_v1`.
- Global projective selector state on C_{v1} : F470 (N516).
- Global projective selector closure output on C_{v1} : F672 (N680).
- Directed/convention-layer lifts and sign-fixing: F690..F692 (with gauge audit P693 and theorem-level packaging N692, N693), with explicit boundaries (no QW-2191 discharge).

A.2 Operational OS support

- Proxy spectra: P694, P696; basis-invariant eigen-spectrum mass observable: F704; observer-limit readout: P697.
- Downstream computability chain: P699 packaged by N699.
- OS support packets: F698/P698/N698 (v1), F700/P700/N700 (v2), F705/P705/N705 (v3).
- Operational ToE closure statement: N701.
- Quadratic mass-proxy meaning definition: N703.
- Residual-sign gauge irrelevance audit (Release-7 OS scope): P709, packaged by N706.
- Previous-methodology survival/global-gap audit for T173: P711, packaged by N707.
- Exact nonexport audit for the next global-provider target T176: P712, packaged by N708.
- Supported-root corridor audit for the strongest current candidate class below T176: P713, packaged by N709.
- Parity-corridor boundary audit for the current `w_break` anchor class: P714, packaged by N710.
- Parity-completed dual-anchor all-root audit below T176: P715, packaged by N711.
- Localized source audit for the remaining exact split below T176: P716, packaged by N712.
- Release-7 OS gauge-irrelevance bridge for the localized exact split below T176: P717, packaged by N713.
- Single mixed linear-weight span insufficiency audit below T176: P718, packaged by N714.

- Low-complexity nonlinear two-readout provider-class audit below T176: P719, packaged by N715.
- Observer-facing signed output-channel projection audit below T176: P720, packaged by N716.
- Source-topology basis-free QW-2191-safe provider nonupgrade audit below T176: P721, packaged by N717.
- Chart-sensitive transported flux/current-like section nonexport audit below T177: P722, packaged by N718.
- Source-topology to atlas chart-seed selection bridge nonexport audit below T178: P723, packaged by N719.
- Positive source-polarity atlas-entry corridor reduction audit below T178: P724, packaged by N720.
- Positive-corridor odd/even lane-selection bridge nonexport audit below T179: P725, packaged by N721.
- Positive-corridor outer/interior chart-selection bridge nonexport audit below T180: P726, packaged by N722.
- Positive-corridor excluded-negative-boundary adjacency chart-selection bridge nonexport audit below T181: P727, packaged by N723.
- Residual-datum source-side boundary-shielded sublane reduction audit below T182: P728, packaged by N724.
- Residual-datum pair1/pair2 orbit-direction selection bridge nonexport audit below T183: P729, packaged by N725.
- Direction-free Shannon residual-datum pair1/pair2 orbit-direction selection bridge nonexport audit below T184: P730, packaged by N726.
- **w_break** witness-payload residual-datum pair1/pair2 orbit-direction promotion-bridge nonexport audit below T185: P731, packaged by N727.
- Pair1-rooted convention-state pair1/pair2 witness-split descent-bridge nonexport audit below T186: P732, packaged by N728.
- Convention-layer pair1/pair2 witness-split transport-descent-bridge nonexport audit below T187: P733, packaged by N729.
- Declared-scope source-topology selector theorem pair1/pair2 orbit-direction descent-bridge nonexport audit below T188: P734, packaged by N730.
- Local source-side scalar-bind pair1/pair2 orbit-direction descent-bridge nonexport audit below T189: P735, packaged by N731.
- Local provider-operator shift-direction pair1/pair2 orbit-direction descent-bridge nonexport audit below T190: P736, packaged by N732.
- Local pair1/pair2 projector-atlas glue orbit-direction descent-bridge nonexport audit below T191: P737, packaged by N733.
- Global projective selector-state pair1/pair2 orbit-direction descent-bridge nonexport audit below T192: P738, packaged by N734.
- Global premise-based directed selector-state pair1/pair2 strict-core-upgrade bridge nonexport audit below T193: P739, packaged by N735.

- Global sign-fixed directed-closure pair1/pair2 strict-core-upgrade bridge nonexport audit below T194: P740, packaged by N736.
- Actual source-topology selector-witness pair1/pair2 promotion-bridge nonexport audit below T195: P741, packaged by N737.
- Non-strict host matching harnesses (external-data dependent): P702, P704.
- Release-7 strict projective OS closure dashboard: P706.

A.3 Emergent observer artifacts

- Observer-limit readout invariants from the projective closure output projector: P697.
- Full induced operator chain computability under $P \mapsto MPM^\top$: P699/N699.

B Explicit numeric artifacts (Release 7)

B.1 Frozen kernel distance profile on $\mathbb{Z}_{12}$

The strict kernel-mode lane uses the frozen symmetric mixing matrix K_{total} on $\mathbb{Z}_{12}$ (specialization route R14, instantiated in F459). On the current repo state, its cyclic-distance profile (for $d = 1, \dots, 6$) is:

d	$k(d)$
1	0.46998567264502017
2	0.19204355169010282
3	0.09142861427792497
4	0.04702916874565040
5	0.02413122336363006
6	0.011070817321442113

The full 12×12 matrix is circulant/symmetric under cyclic distance:

$$(K_{\text{total}})_{ij} = k(d_{\text{cyc}}(i, j)), \quad k(0) = 0, \quad k(d) = k(12 - d) \quad (d = 1, \dots, 6).$$

B.2 Explicit diagonal/local Psi-sector Hessian instantiation (F459)

The diagonal/local Psi-sector Hessian instantiation F459 exports a symmetric matrix $H_\psi \in \mathbb{R}^{12 \times 12}$. On the current repo state it decomposes *exactly* as:

$$(H_\psi)_{ij} = \begin{cases} m_0^2 + d_{\text{local}}(i), & i = j, \\ k(d_{\text{cyc}}(i, j)), & i \neq j, \end{cases}$$

with

$$m_0^2 = 1.013551972358388.$$

The exported diagonal residual profile $d_{\text{local}}(i)$ and the resulting diagonal are:

i	$d_{\text{local}}(i)$	$(H_\psi)_{ii} = m_0^2 + d_{\text{local}}(i)$
0	0.34269049977246646	1.3562424721308544
1	11.518361441687219	12.531913414045606
2	0.5284499241054783	1.5420018964638662
3	-0.1567291917971062	0.8568227805612817
4	-0.5341686256176337	0.4793833467407542
5	1.2939217196807558	2.3074736920391437
6	-0.5936675263840503	0.4198844459743376
7	1.2939217196807553	2.3074736920391430
8	-0.5341686256176337	0.4793833467407542
9	-0.1567291917971062	0.8568227805612817
10	0.5284499241054781	1.5420018964638660
11	11.518361441687219	12.531913414045606

All full numeric matrices (K_{total}, H_ψ) and the invariant eigen-spectrum are exported in `fundamental_action_re`

C Frozen non-strict host matching layer (dataset vs policy)

C.1 External Standard-Model mass dataset (v1)

The repo freezes the external dataset (targets-only) as: `fundamental_action_reconstruction/external_data/sm` with dataset id `SM_mass_targets_v1_pdg2024_summary_tables`. For reader convenience, the current target list (units: GeV) is:

id	m (GeV)	σ_m (GeV)
e	0.00051099895	1.5×10^{-13}
u	0.00216	7.0×10^{-5}
d	0.0047	7.0×10^{-5}
μ	0.1056583755	2.3×10^{-9}
s	0.0935	8.0×10^{-4}
τ	1.77693	9.0×10^{-5}
c	1.273	4.6×10^{-3}
b	4.183	7.0×10^{-3}
W	80.3692	1.33×10^{-2}
Z	91.188	2.0×10^{-3}
H	125.2	1.1×10^{-1}
t	172.57	2.9×10^{-1}

C.2 External host matching policy (v1, frozen no-pass)

The repo freezes the external matching policy as: `fundamental_action_reconstruction/external_data/sm_host` with policy id `SM_host_matching_policy_v1_centered_log_assignment_with_global_scale`. The policy declares:

- an assignment algorithm: linear sum assignment on squared centered $\log m^2$ differences,
- **allowed DOF:** only a single global scale factor (defined by centered $\log m^2$ matching after the assignment),
- **disallowed DOF:** sector scales, kinetic-metric refits, nonlinear mass maps,
- **pass criteria:** `pass_criteria=null`, with semantics: *no pass/FAIL claim is permitted* under this policy version (diagnostic metrics only).

Hard limits: this entire layer remains non-strict and external-data dependent. It is not part of the strict OS closure bundle.

D Release-8 addendum: QW-2191 closure narrative and strict-core-to-SM scaffold

D.1 Scope and status boundary

This addendum records a formal narrative for the Release-8 strict O3 closure line associated with QW-2191. It preserves strict status discipline: closure is asserted in the declared O3 pipeline semantics and is not silently promoted to strongest global ToE closure.

D.2 Operational strict kernel and selector score

The strict operational kernel used in the later-pipeline gate computations is:

$$K_{\text{strict}}(d) = \frac{\cos(\omega d + \phi)}{1 + \beta d^\eta}.$$

The candidate-score map used in the closure lane is:

$$s(\varphi, a) = (\varphi - 0.40) + 0.9(a - 0.58) - 0.6(\varphi - 0.40)^2,$$

with sign rule

$$\text{sign}(s) = \begin{cases} +1, & s \geq 0, \\ -1, & s < 0. \end{cases}$$

D.3 Compressed derivation history to Release 8

1. Kernel-role discipline was made explicit (legacy vs strict), preventing silent role transfer.
2. O3 chain packetization was executed: candidate class definition, normalization map, pairwise matrix, sweep/replay/adversarial diagnostics.
3. Residual ambiguity handling triggered candidate evolution (v2 falsified, v3 degenerate, v4 retained).
4. Formal export obligations were completed including global L2 export completion and checker refresh.
5. Independent replication packet preserved closure output under the same declared rigor.

Hence, the exported O3 endpoint is strict-closed in pipeline semantics.

D.4 Strict-core candidate Lagrangian and SM-shaped extension scaffold

Core nadsoliton candidate density:

$$\mathcal{L}_{\text{core}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \sum_{i=0}^{11} \left[\frac{1}{2} \partial_\mu \psi_i \partial^\mu \psi_i - V(\psi_i) \right] + \frac{1}{2} \sum_{i \neq j} K_{ij} \psi_i \psi_j,$$

with

$$K_{ij} = K_{\text{strict}}(d_{ij}).$$

A full Standard-Model-shaped effective scaffold (for explicit interface bookkeeping) may be written as:

$$\mathcal{L}_{\text{eff}} = -\frac{1}{4}G_{\mu\nu}^A G^{A\mu\nu} - \frac{1}{4}W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu} + \sum_f \bar{\Psi}_f i\gamma^\mu D_\mu \Psi_f + (D_\mu H)^\dagger (D^\mu H) - V(H) + \mathcal{L}_{\text{Yukawa}} + \Delta\mathcal{L}_{\text{nadsoliton}}.$$

where

$$\mathcal{L}_{\text{Yukawa}} = -y_u \bar{Q} \tilde{H} u_R - y_d \bar{Q} H d_R - y_e \bar{L} H e_R + \text{h.c.}$$

and $\Delta\mathcal{L}_{\text{nadsoliton}}$ stores strict-core induced effective corrections/couplings.

This is a documented candidate integration scaffold; it is not, by itself, a claim that host-level SM/GR identification is already fully discharged.

E Full Standard Model + Nadsoliton Effective Lagrangian (Release-8 formal scaffold)

E.1 Purpose and status discipline

This chapter provides a full explicit Standard-Model-shaped Lagrangian embedding with a nadsoliton effective sector. It is a formal *candidate export scaffold*: complete at the symbolic field-theory level, but not automatically an already-discharged strict host-identification theorem.

E.2 Full Standard Model density

The Standard Model density is recorded in the conventional decomposition:

$$\mathcal{L}_{\text{SM}} = \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{fermion}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{Yukawa}} + \mathcal{L}_{\text{gf}} + \mathcal{L}_{\text{ghost}}.$$

Gauge kinetic sector:

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4}G_{\mu\nu}^A G^{A\mu\nu} - \frac{1}{4}W_{\mu\nu}^I W^{I\mu\nu} - \frac{1}{4}B_{\mu\nu} B^{\mu\nu}.$$

Fermion kinetic/covariant sector (three generations):

$$\mathcal{L}_{\text{fermion}} = \sum_{g=1}^3 \left(\bar{Q}_g i\gamma^\mu D_\mu Q_g + \bar{u}_{R,g} i\gamma^\mu D_\mu u_{R,g} + \bar{d}_{R,g} i\gamma^\mu D_\mu d_{R,g} + \bar{L}_g i\gamma^\mu D_\mu L_g + \bar{e}_{R,g} i\gamma^\mu D_\mu e_{R,g} \right).$$

Higgs sector:

$$\mathcal{L}_{\text{Higgs}} = (D_\mu H)^\dagger (D^\mu H) - V(H), \quad V(H) = -\mu_H^2 H^\dagger H + \lambda_H (H^\dagger H)^2.$$

Yukawa sector:

$$\mathcal{L}_{\text{Yukawa}} = -\bar{Q} Y_d H d_R - \bar{Q} Y_u \tilde{H} u_R - \bar{L} Y_e H e_R + \text{h.c.}$$

with $\tilde{H} = i\sigma_2 H^*$. Gauge-fixing and ghost terms are retained symbolically in $\mathcal{L}_{\text{gf}}, \mathcal{L}_{\text{ghost}}$ according to the selected quantization scheme.

E.3 Nadsoliton strict-core sector and coupling interface

Strict-core candidate:

$$\mathcal{L}_{\text{core}} = \frac{1}{2} \partial_\mu \phi \partial^\mu \phi + \sum_{i=0}^{11} \left(\frac{1}{2} \partial_\mu \psi_i \partial^\mu \psi_i - V_i(\psi_i) \right) + \frac{1}{2} \sum_{i \neq j} K_{ij} \psi_i \psi_j,$$

$$K_{ij} = K_{\text{strict}}(d_{ij}) = \frac{\cos(\omega d_{ij} + \varphi)}{1 + \beta d_{ij}^\eta}.$$

A minimal effective interaction bridge to SM-shaped observables is represented as:

$$\Delta\mathcal{L}_{\text{bridge}} = -\frac{c_G}{4}f_G(\phi)G_{\mu\nu}^AG^{A\mu\nu} - \frac{c_W}{4}f_W(\phi)W_{\mu\nu}^IW^{I\mu\nu} - \frac{c_B}{4}f_B(\phi)B_{\mu\nu}B^{\mu\nu} + c_H f_H(\phi)H^\dagger H + \sum_f c_f f_f(\phi)\bar{\Psi}_f\Psi_f.$$

Hence the release-level effective total density is:

$$\mathcal{L}_{\text{tot}}^{(R8)} = \mathcal{L}_{\text{SM}} + \mathcal{L}_{\text{core}} + \Delta\mathcal{L}_{\text{bridge}}.$$

F Theory-closure blocker map (Release 8)

F.1 Purpose of this chapter

This chapter separates what is already closed in the O3 gate from what still blocks the strongest possible global closure claim (global scope, host-identification scope, strict-core completeness scope).

F.2 Current hard blocker classes

1. **O3 semantic boundary:** QW-2191 closure in O3 semantics is not identical to strongest-form global ToE closure.
2. **Host-level SM/GR identification:** one fully exported strict theorem mapping the full observable sector to $\mathcal{L}_{\text{tot}}^{(R8)}$ without extra fitting freedoms is still required.
3. **Selector robustness outside declared pipeline:** additional independent multi-environment audits and cross-regression checks are still required.
4. **Bridge-calibration identifiability:** the functions f_G, f_W, f_B, f_H, f_f and couplings c . still require explicit strict identifiability/injectivity exports.
5. **Global closure theorem packaging:** a single theorem packet stitching all local closures into the strongest global export is still pending.

F.3 Technical execution order

R8-B1 Freeze a reference reproduction test set for O3 plus external replication matrix.

R8-B2 Export theorem-level identifiability for bridge parameters in $\Delta\mathcal{L}_{\text{bridge}}$.

R8-B3 Add a strict host-matching theorem without redundant degrees of freedom (or an explicit non-bridge/non-identification theorem).

R8-B4 Merge all validated parts into one global-closure packet with explicit limits and theorem scope.

R8-B5 Execute an independent external audit (replication + falsification challenge) before any global status upgrade.

F.4 Practical verdict

Release 8 may be published as a professor-grade formal state: O3/QW-2191 strict gate closed in pipeline semantics, full Lagrangian scaffold recorded, and explicit blocker map for strongest-theory closure retained.

G Current closure state theorem map (Release 8, current-only)

G.1 Current-state closure declaration

On the current repository state, the declared Release-8 strict package exports closure in its declared scope, with completion chain:

$$P1343 \rightarrow P1344 \rightarrow P1345 \rightarrow P1346 \rightarrow P1347 \rightarrow P1348.$$

Current closure status object:

$$\text{global_closure_theorem_status_r8_declared_scope} = \text{EXPORTED_CLOSED}.$$

G.2 Audit status boundary (external step still pending)

The only remaining upgrade step is external blind audit execution ('P1349'). Thus:

- closure theorem package in declared scope: exported and closed,
- external blind-audit protocol: exported,
- external blind-audit execution: pending.

G.3 Full Lagrangian status in current Release 8

The current Release-8 documentation includes the full Standard-Model-shaped decomposition plus strict-core nadsoliton sector and bridge terms in the effective total density $\mathcal{L}_{\text{tot}}^{(R8)}$.

In the current-state reading discipline, this Lagrangian package is treated as the active formal physics interface of the closed declared-scope theorem lane.

G.4 No-obsolete-claim discipline

Older narrative fragments that represented pre-closure uncertainty are superseded by the current closure chain exports listed above and should not be interpreted as current status.

H Release-8 professional scientific executive synthesis

H.1 Scope, closure claim, and semantics discipline

This Release-8 document records the strongest currently exported strict lane under explicit no-false-pass discipline. The closure claim is:

$$\text{kernel_alone_fundamental_nonuniqueness_status_strict_internal_source_}(QW-2191) = \text{CLOSED_FULL_STR}$$

This closure is reported in the strict internal-source lane built by the exported chain $P1343 \rightarrow P1344 \rightarrow P1345 \rightarrow P1346 \rightarrow P1347 \rightarrow P1348$, with theorem export, stress validation, independent replication/challenge, and long-horizon stability audit.

H.2 Kernel-first derivation architecture

All current derivation-facing closure claims in this Release-8 synthesis are expressed in the strict-kernel lane:

$$K_{\text{strict}}(d) = \frac{\cos(\omega d + \phi)}{1 + \beta d^\eta}.$$

The selector source is exported internally as $S_{\text{strict_internal_v1}}$, producing a single-valued selector map on declared admissible classes. Thus, for the targeted ambiguity class (real Fourier basis + pair-plane isotropy), closure is kernel-driven in this lane.

H.3 Scientific reading guide for reviewers

For strict external review, the minimal chain to verify is:

1. theorem-level internal selector source export (P1343),
2. stress validation and rollback law (P1344),
3. independent replication and adversarial challenge (P1345),
4. long-horizon drift/regression stability (P1346).

Only after checking this chain should a reviewer accept the Release-8 closure wording.

H.4 Hard limits retained

This release still separates:

- strict kernel-alone closure lane claims,
- host-level identification claims,
- external-data policy layers.

No silent upgrade across those boundaries is admitted.

H.5 Professional publication statement

Within the declared strict internal-source lane and exported audit chain, Release 8 is presented as a closed scientific package for the target fundamental nonuniqueness class, with explicit rollback governance under future contradictory evidence.

Reproducibility

- **As-of date:** 2026-05-12
- **Build smoke probe (PDF + strict dashboards):** P707 (fundamental_action_reconstruction/p707_current_release_7_build_and_closure_smoke_probe.py)
- **Strict OS closure dashboard:** P706 (fundamental_action_reconstruction/p706_current_release_7_strict_projective_operational_toe_os_closure_dashboard_probe.py)
- **Strict T173 frontier dashboard:** P708 (fundamental_action_reconstruction/p708_current_strict_t173_frontier_dashboard_probe.py)
- **Previous-methodology survival audit:** P711 (fundamental_action_reconstruction/p711_current_strict_t173_previous_methodology_survival_and_global_gap_audit_probe.py)
- **Exact global-provider nonexport audit:** P712 (fundamental_action_reconstruction/p712_current_strict_t176_existing_global_directed_sign_coherence_provider_nonexport_audit_probe.py)
- **Supported-root corridor audit for the strongest current candidate class:** P713 (fundamental_action_reconstruction/p713_current_strict_t176_multiroot_rooted_sign_lift_root_independence_audit_probe.py)

- **Parity diagnosis of the current w_break corridor: P714**
(fundamental_action_reconstruction/p714_current_strict_t176_w_break_parity_root_support_profile_audit_probe.py)
- **Parity-completed dual-anchor all-root audit: P715**
(fundamental_action_reconstruction/p715_current_strict_t176_parity_completed_dual_anchor_multiroot_audit_probe.py)
- **Localized source audit for the remaining exact split: P716**
(fundamental_action_reconstruction/p716_current_strict_t176_pair4_negative_cosine_polarity_global_z2_orbit_split_audit_probe.py)
- **Release-7 OS bridge audit for the localized exact split: P717**
(fundamental_action_reconstruction/p717_current_strict_t176_pair4_exact_branch_split_release_7_os_gauge_irrelevance_bridge_audit_probe.py)
- **Single mixed linear-weight span insufficiency audit: P718**
(fundamental_action_reconstruction/p718_current_strict_t176_single_mixed_linear_weight_span_provider_insufficiency_audit_probe.py)
- **Low-complexity nonlinear two-readout provider-class audit: P719**
(fundamental_action_reconstruction/p719_current_strict_t176_low_complexity_odd_polynomial_two_readout_provider_class_audit_probe.py)
- **Observer-facing signed output-channel projection audit: P720**
(fundamental_action_reconstruction/p720_current_strict_t176_observer_facing_signed_output_channel_projection_provider_class_audit_probe.py)
- **Source-topology basis-free QW-2191-safe provider nonupgrade audit: P721**
(fundamental_action_reconstruction/p721_current_strict_t176_source_topology_basis_free_qw2191_safe_provider_nonupgrade_audit_probe.py)
- **Chart-sensitive transported flux/current-like section nonexport audit: P722**
(fundamental_action_reconstruction/p722_current_strict_t177_chart_sensitive_transportied_flux_current_like_section_nonexport_audit_probe.py)
- **Source-topology to atlas chart-seed selection bridge nonexport audit: P723**
(fundamental_action_reconstruction/p723_current_strict_t178_source_topology_to_atlas_chart_seed_selection_bridge_nonexport_audit_probe.py)
- **Positive source-polarity atlas-entry corridor reduction audit: P724**
(fundamental_action_reconstruction/p724_current_strict_t178_positive_source_polarity_atlas_entry_corridor_reduction_audit_probe.py)
- **Positive-corridor odd/even lane-selection bridge nonexport audit: P725**
(fundamental_action_reconstruction/p725_current_strict_t179_positive_corridor_odd_even_lane_selection_bridge_nonexport_audit_probe.py)
- **Positive-corridor outer/interior chart-selection bridge nonexport audit: P726**
(fundamental_action_reconstruction/p726_current_strict_t180_positive_corridor_outer_interior_chart_selection_bridge_nonexport_audit_probe.py)
- **Positive-corridor excluded-negative-boundary adjacency chart-selection bridge nonexport audit: P727** (fundamental_action_reconstruction/p727_current_strict_t181_positive_corridor_excluded_negative_boundary_adjacency_chart_selection_bridge_nonexport_audit_probe.py)

- **Residual-datum source-side boundary-shielded sublane reduction audit: P728**
(`fundamental_action_reconstruction/p728_current_strict_t182_residual_datum_source_side_boundary_shielded_sublane_reduction_audit_probe.py`)
- **Residual-datum pair1/pair2 orbit-direction selection bridge nonexport audit: P729**
(`fundamental_action_reconstruction/p729_current_strict_t183_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe.py`)
- **Direction-free Shannon residual-datum pair1/pair2 orbit-direction selection bridge nonexport audit: P730** (`fundamental_action_reconstruction/p730_current_strict_t184_direction_free_shannon_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe.py`)
- **w_break witness-payload residual-datum pair1/pair2 orbit-direction promotion-bridge nonexport audit: P731** (`fundamental_action_reconstruction/p731_current_strict_t185_w_break_witness_payload_residual_datum_pair12_orbit_direction_promotion_bridge_nonexport_audit_probe.py`)
- **Pair1-rooted convention-state pair1/pair2 witness-split descent-bridge nonexport audit: P732**
(`fundamental_action_reconstruction/p732_current_strict_t186_pair1_rooted_convention_state_pair12_witness_split_descent_bridge_nonexport_audit_probe.py`)
- **Convention-layer pair1/pair2 witness-split transport-descent-bridge nonexport audit: P733**
(`fundamental_action_reconstruction/p733_current_strict_t187_convention_layer_pair12_witness_split_transport_descent_bridge_nonexport_audit_probe.py`)
- **Declared-scope source-topology selector theorem pair1/pair2 orbit-direction descent-bridge nonexport audit: P734** (`fundamental_action_reconstruction/p734_current_strict_t188_declared_scope_source_topology_selector_theorem_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py`)
- **Local source-side scalar-bind pair1/pair2 orbit-direction descent-bridge nonexport audit: P735**
(`fundamental_action_reconstruction/p735_current_strict_t189_local_source_side_scalar_bind_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py`)
- **Local provider-operator shift-direction pair1/pair2 orbit-direction descent-bridge nonexport audit: P736** (`fundamental_action_reconstruction/p736_current_strict_t190_local_provider_operator_shift_direction_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py`)
- **Local pair1/pair2 projector-atlas glue orbit-direction descent-bridge nonexport audit: P737**
(`fundamental_action_reconstruction/p737_current_strict_t191_local_pair12_projector_atlas_glue_orbit_direction_descent_bridge_nonexport_audit_probe.py`)
- **Global projective selector-state pair1/pair2 orbit-direction descent-bridge nonexport audit: P738**
(`fundamental_action_reconstruction/p738_current_strict_t192_global_projective_selector_state_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py`)
- **Global premise-based directed selector-state pair1/pair2 strict-core-upgrade bridge nonexport audit: P739** (`fundamental_action_reconstruction/p739_current_strict`

t193_global_premise_based_directed_selector_state_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe.py)

- **Global sign-fixed directed-closure pair1/pair2 strict-core-upgrade bridge nonexport audit:** P740 (fundamental_action_reconstruction/p740_current_strict_t194_global_sign_fixed_directed_closure_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe.py)
- **Actual source-topology selector-witness pair1/pair2 promotion-bridge nonexport audit:** P741 (fundamental_action_reconstruction/p741_current_strict_t195_actual_source_topology_selector_witness_pair12_witness_split_promotion_bridge_nonexport_audit_probe.py)

To reproduce the PDF and verify the current Release-7 strict dashboard statuses:

```

1 git rev-parse HEAD
2 python3 fundamental_action_reconstruction/\
3 p707_current_release_7_build_and_closure_smoke_probe.py
4 python3 fundamental_action_reconstruction/\
5 p711_current_strict_t173_previous_methodology_survival_and_global_gap_audit_probe.py
6 python3 fundamental_action_reconstruction/\
7 p712_current_strict_t176_existing_global_directed_sign_coherence_provider_nonexport_audit_probe.py
8 python3 fundamental_action_reconstruction/\
9 p713_current_strict_t176_multiroot_rooted_sign_lift_root_independence_audit_probe.py
10 python3 fundamental_action_reconstruction/\
11 p714_current_strict_t176_w_break_parity_root_support_profile_audit_probe.py
12 python3 fundamental_action_reconstruction/\
13 p715_current_strict_t176_parity_completed_dual_anchor_multiroot_audit_probe.py
14 python3 fundamental_action_reconstruction/\
15 p716_current_strict_t176_pair4_negative_cosine_polarity_global_z2_orbit_split_audit_probe.py
16 python3 fundamental_action_reconstruction/\
17 p717_current_strict_t176_pair4_exact_branch_split_release_7_os_gauge_irrelevance_bridge_audit_probe.py
18 python3 fundamental_action_reconstruction/\
19 p718_current_strict_t176_single_mixed_linear_weight_span_provider_insufficiency_audit_probe.py
20 python3 fundamental_action_reconstruction/\
21 p719_current_strict_t176_low_complexity_odd_polynomial_two_readout_provider_class_audit_probe.py
22 python3 fundamental_action_reconstruction/\
23 p720_current_strict_t176_observer_facing_signed_output_channel_projection_provider_class_audit_probe.py
24 python3 fundamental_action_reconstruction/\
25 p721_current_strict_t176_source_topology_basis_free_qw2191_safe_provider_nonupgrade_audit_probe.py
26 python3 fundamental_action_reconstruction/\
27 p722_current_strict_t177_chart_sensitive_transported_flux_current_like_section_nonexport_audit_probe.py
28 python3 fundamental_action_reconstruction/\
29 p723_current_strict_t178_source_topology_to_atlas_chart_seed_selection_bridge_nonexport_audit_probe.py
30 python3 fundamental_action_reconstruction/\
31 p724_current_strict_t178_positive_source_polarity_atlas_entry_corridor_reduction_audit_probe.py
32 python3 fundamental_action_reconstruction/\
33 p725_current_strict_t179_positive_corridor_odd_even_lane_selection_bridge_nonexport_audit_probe.py
34 python3 fundamental_action_reconstruction/\
35 p726_current_strict_t180_positive_corridor_outer_interior_chart_selection_bridge_nonexport_audit_probe.py
36 python3 fundamental_action_reconstruction/\
37 p727_current_strict_t181_positive_corridor_excluded_negative_boundary_adjacency_chart_selection_bridge_nonexport_audit_probe.py
38 python3 fundamental_action_reconstruction/\
39 p728_current_strict_t182_residual_datum_source_side_boundary_shielded_sublane_reduction_audit_probe.py
40 python3 fundamental_action_reconstruction/\
41 p729_current_strict_t183_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe.py
42 python3 fundamental_action_reconstruction/\
43 p730_current_strict_t184_direction_free_shannon_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe.py
44 python3 fundamental_action_reconstruction/\
45 p731_current_strict_t185_w_break_witness_payload_residual_datum_pair12_orbit_direction_promotion_bridge_nonexport_audit_probe.py
46 python3 fundamental_action_reconstruction/\
47 p732_current_strict_t186_pair1_rooted_convention_state_pair12_witness_split_descent_bridge_nonexport_audit_probe.py
48 python3 fundamental_action_reconstruction/\
49 p733_current_strict_t187_convention_layer_pair12_witness_split_transport_descent_bridge_nonexport_audit_probe.py
50 python3 fundamental_action_reconstruction/\
51 p734_current_strict_t188_declared_scope_source_topology_selector_theorem_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py
52 python3 fundamental_action_reconstruction/\
53 p735_current_strict_t189_local_source_side_scalar_bind_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py
54 python3 fundamental_action_reconstruction/\
55 p736_current_strict_t190_local_provider_operator_shift_direction_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py
56 python3 fundamental_action_reconstruction/\
57 p737_current_strict_t191_local_pair12_projector_atlas_glue_orbit_direction_descent_bridge_nonexport_audit_probe.py
58 python3 fundamental_action_reconstruction/\
59 p738_current_strict_t192_global_projective_selector_state_pair12_orbit_direction_descent_bridge_nonexport_audit_probe.py
60 python3 fundamental_action_reconstruction/\
61 p739_current_strict_t193_global_premise_based_directed_selector_state_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe.py
62 python3 fundamental_action_reconstruction/\
63 p740_current_strict_t194_global_sign_fixed_directed_closure_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe.py
64 python3 fundamental_action_reconstruction/\
65 p741_current_strict_t195_actual_source_topology_selector_witness_pair12_witness_split_promotion_bridge_nonexport_audit_probe.py
66 cat fundamental_action_reconstruction/generated/\
67 p707_current_release_7_build_and_closure_smoke_probe_summary.json
68 cat fundamental_action_reconstruction/generated/\
69 p711_current_strict_t173_previous_methodology_survival_and_global_gap_audit_probe_summary.json
70 cat fundamental_action_reconstruction/generated/\
71 p712_current_strict_t176_existing_global_directed_sign_coherence_provider_nonexport_audit_probe_summary.json
72 cat fundamental_action_reconstruction/generated/\
73 p713_current_strict_t176_multiroot_rooted_sign_lift_root_independence_audit_probe_summary.json
74 cat fundamental_action_reconstruction/generated/\

```



```

75 | p714_current_strict_t176_v_break_parity_root_support_profile_audit_probe_summary.json
76 | cat fundamental_action_reconstruction/generated/\
77 | p715_current_strict_t176_parity_completed_dual_anchor_multiroot_audit_probe_summary.json
78 | cat fundamental_action_reconstruction/generated/\
79 | p716_current_strict_t176_pair4_negative_cosine_polarity_global_z2_orbit_split_audit_probe_summary.json
80 | cat fundamental_action_reconstruction/generated/\
81 | p717_current_strict_t176_pair4_exact_branch_split_release_7_os_gauge_irrelevance_bridge_audit_probe_summary.json
82 | cat fundamental_action_reconstruction/generated/\
83 | p718_current_strict_t176_single_mixed_linear_weight_span_provider_insufficiency_audit_probe_summary.json
84 | cat fundamental_action_reconstruction/generated/\
85 | p719_current_strict_t176_low_complexity_odd_polynomial_two_readout_provider_class_audit_probe_summary.json
86 | cat fundamental_action_reconstruction/generated/\
87 | p720_current_strict_t176_observer_facing_signed_output_channel_projection_provider_class_audit_probe_summary.json
88 | cat fundamental_action_reconstruction/generated/\
89 | p721_current_strict_t176_source_topology_basis_free_qv2191_safe_provider_nonupgrade_audit_probe_summary.json
90 | cat fundamental_action_reconstruction/generated/\
91 | p722_current_strict_t177_chart_sensitive_transported_flux_current_like_section_nonexport_audit_probe_summary.json
92 | cat fundamental_action_reconstruction/generated/\
93 | p723_current_strict_t178_source_topology_to_atlas_chart_seed_selection_bridge_nonexport_audit_probe_summary.json
94 | cat fundamental_action_reconstruction/generated/\
95 | p724_current_strict_t178_positive_source_polarity_atlas_entry_corridor_reduction_audit_probe_summary.json
96 | cat fundamental_action_reconstruction/generated/\
97 | p725_current_strict_t179_positive_corridor_odd_even_lane_selection_bridge_nonexport_audit_probe_summary.json
98 | cat fundamental_action_reconstruction/generated/\
99 | p726_current_strict_t180_positive_corridor_outer_interior_chart_selection_bridge_nonexport_audit_probe_summary.json
100 | cat fundamental_action_reconstruction/generated/\
101 | p727_current_strict_t181_positive_corridor_excluded_negative_boundary_adjacency_chart_selection_bridge_nonexport_audit_probe_summary.json
102 | cat fundamental_action_reconstruction/generated/\
103 | p728_current_strict_t182_residual_datum_source_side_boundary_shielded_sublane_reduction_audit_probe_summary.json
104 | cat fundamental_action_reconstruction/generated/\
105 | p729_current_strict_t183_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe_summary.json
106 | cat fundamental_action_reconstruction/generated/\
107 | p730_current_strict_t184_direction_free_shannon_residual_datum_pair12_orbit_direction_selection_bridge_nonexport_audit_probe_summary.json
108 | cat fundamental_action_reconstruction/generated/\
109 | p731_current_strict_t185_v_break_witness_payload_residual_datum_pair12_orbit_direction_promotion_bridge_nonexport_audit_probe_summary.json
110 | cat fundamental_action_reconstruction/generated/\
111 | p732_current_strict_t186_pair1_rooted_convention_state_pair12_witness_split_descent_bridge_nonexport_audit_probe_summary.json
112 | cat fundamental_action_reconstruction/generated/\
113 | p733_current_strict_t187_convention_layer_pair12_witness_split_transport_descent_bridge_nonexport_audit_probe_summary.json
114 | cat fundamental_action_reconstruction/generated/\
115 | p734_current_strict_t188_declared_scope_source_topology_selector_theorem_pair12_orbit_direction_descent_bridge_nonexport_audit_probe_summary.json
116 | cat fundamental_action_reconstruction/generated/\
117 | p735_current_strict_t189_local_source_side_scalar_bind_pair12_orbit_direction_descent_bridge_nonexport_audit_probe_summary.json
118 | cat fundamental_action_reconstruction/generated/\
119 | p736_current_strict_t190_local_provider_operator_shift_direction_pair12_orbit_direction_descent_bridge_nonexport_audit_probe_summary.json
120 | cat fundamental_action_reconstruction/generated/\
121 | p737_current_strict_t191_local_pair12_projector_atlas_glue_orbit_direction_descent_bridge_nonexport_audit_probe_summary.json
122 | cat fundamental_action_reconstruction/generated/\
123 | p738_current_strict_t192_global_projective_selector_state_pair12_orbit_direction_descent_bridge_nonexport_audit_probe_summary.json
124 | cat fundamental_action_reconstruction/generated/\
125 | p739_current_strict_t193_global_premise_based_directed_selector_state_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe_summary.
    | json
126 | cat fundamental_action_reconstruction/generated/\
127 | p740_current_strict_t194_global_sign_fixed_directed_closure_pair12_witness_split_strict_core_upgrade_bridge_nonexport_audit_probe_summary.json
128 | cat fundamental_action_reconstruction/generated/\
129 | p741_current_strict_t195_actual_source_topology_selector_witness_pair12_witness_split_promotion_bridge_nonexport_audit_probe_summary.json
130 | cat fundamental_action_reconstruction/generated/\
131 | p706_current_release_7_strict_projective_operational_toe_os_closure_dashboard_probe_summary.json
132 | cat fundamental_action_reconstruction/generated/\
133 | p708_current_strict_t173_frontier_dashboard_probe_summary.json

```