




4721 Core Mathematics 1

1	$\frac{4(3+\sqrt{7})}{(3-\sqrt{7})(3+\sqrt{7})}$ $= \frac{12+4\sqrt{7}}{9-7}$ $= 6 + 2\sqrt{7}$	M1 B1 A1 $\frac{3}{3}$	Multiply top and bottom by conjugate 9 ± 7 soi in denominator $6 + 2\sqrt{7}$
2(i) (ii)	$x^2 + y^2 = 49$ $x^2 + y^2 - 6x - 10y - 30 = 0$ $(x-3)^2 - 9 + (y-5)^2 - 25 - 30 = 0$ $(x-3)^2 + (y-5)^2 = 64$ $r^2 = 64$ $r = 8$	B1 1 M1 A1 $\frac{2}{3}$	$x^2 + y^2 = 49$ $3^2 \ 5^2 \ 30$ with consistent signs soi 8 cao
3	$a(x+3)^2 + c = 3x^2 + bx + 10$ $3(x^2 + 6x + 9) + c = 3x^2 + bx + 10$ $3x^2 + 18x + 27 + c = 3x^2 + bx + 10$ $c = -17$	B1 B1 M1 A1 $\frac{4}{4}$	$a = 3$ soi $b = 18$ soi $c = 10 - 9a$ or $c = 10 - \frac{b^2}{12}$ $c = -17$
4(i) (ii) (iii)	$p = -1$ $\sqrt{25k^2} = 15$ $25k^2 = 225$ $k^2 = 9$ $k = \pm 3$ $\sqrt[3]{t} = 2$ $t = 8$	B1 1 M1 A1 A1 3 M1 A1 $\frac{2}{6}$	$p = -1$ Attempt to square 15 or attempt to square root $25k^2$ $k = 3$ $k = -3$ $\frac{1}{t^{\frac{1}{3}}} = \frac{1}{2}$ or $t^{\frac{1}{3}} = 2$ soi $t = 8$

5(i)		B1 B1 2	+ve cubic +ve or -ve cubic with point of inflection at (0, 2) and no max/min points
(ii)		B1 B1 2	curve with correct curvature in +ve quadrant only completely correct curve
(iii)	Stretch scale factor 1.5 parallel to y-axis	B1 B1 B1 3 <u>7</u>	stretch factor 1.5 parallel to y-axis or in y-direction
6(i)	<p>EITHER</p> $x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$ $x = \frac{-8 \pm \sqrt{64 - 40}}{2}$ $x = \frac{-8 \pm \sqrt{24}}{2}$ $x = \frac{-8 \pm 2\sqrt{6}}{2}$ $x = -4 \pm \sqrt{6}$ <p>OR</p> $(x+4)^2 - 16 + 10 = 0$ $(x+4)^2 = 6$ $x+4 = \pm\sqrt{6} \quad \text{M1 A1}$ $x = \pm\sqrt{6} - 4 \quad \text{A1}$	M1 A1 A1 3	<p>Correct method to solve quadratic</p> $x = \frac{-8 \pm \sqrt{24}}{2}$ $x = -4 \pm \sqrt{6}$
(ii)		B1 B1 B1 3	+ve parabola parabola cutting y-axis at (0, 10) where (0, 10) is not min/max point parabola with 2 negative roots
(iii)	$x \leq -\sqrt{6} - 4, x \geq \sqrt{6} - 4$	M1 A1 ft 2 <u>8</u>	$x \leq \text{lower root } x \geq \text{higher root} \quad (\text{allow } <, >)$ Fully correct answer, ft from roots found in (i)

7(i)	Gradient = $-\frac{1}{2}$	B1 1	$-\frac{1}{2}$
(ii)	$y - 5 = -\frac{1}{2}(x - 6)$ $2y - 10 = -x + 6$ $x + 2y - 16 = 0$	M1 B1 ft A1 3	Equation of straight line through (6, 5) with any non-zero numerical gradient Uses gradient found in (i) in their equation of line Correct answer in correct form (integer coefficients)
(iii)	EITHER $\frac{4-x}{2} = x^2 + x + 1$ $4 - x = 2x^2 + 2x + 2$ $2x^2 + 3x - 2 = 0$ $(2x - 1)(x + 2) = 0$ $x = \frac{1}{2}, x = -2$ $y = \frac{7}{4}, y = 3$ OR $y = (4 - 2y)^2 + (4 - 2y) + 1$ $y = 16 - 16y + 4y^2 + 4 - 2y + 1$ $0 = 21 - 19y + 4y^2$ $0 = (4y - 7)(y - 3)$ $y = \frac{7}{4}, y = 3$ $x = \frac{1}{2}, x = -2$	*M1 DM1 A1 A1 4 *M1 DM1 A1 A1	Substitute to find an equation in x (or y) Correct method to solve quadratic $x = \frac{1}{2}, x = -2$ $y = \frac{7}{4}, y = 3$ SR one correct (x,y) pair www B1
8			

8(i)	$\frac{dy}{dx} = 3x^2 + 2x - 1$ <p>At stationary points, $3x^2 + 2x - 1 = 0$ $(3x - 1)(x + 1) = 0$ $x = \frac{1}{3}, x = -1$ $y = \frac{76}{27}, y = 4$</p>	*M1 A1 M1 DM1 A1 A1 6	<p>Attempt to differentiate (at least one correct term) 3 correct terms</p> <p>Use of $\frac{dy}{dx} = 0$</p> <p>Correct method to solve 3 term quadratic</p> <p>$x = \frac{1}{3}, x = -1$</p> <p>$y = \frac{76}{27}, 4$</p> <p>SR one correct (x,y) pair www B1</p> <p>M1 Looks at sign of $\frac{d^2y}{dx^2}$ for at least one of their x-values or other correct method</p> <p>A1 $x = \frac{1}{3}$, minimum point CWO</p> <p>A1 3 $x = -1$, maximum point CWO</p> <p>M1 Any inequality (or inequalities) involving both their x values from part (i)</p> <p>A1 2 Correct inequality (allow $<$ or \leq)</p>
(ii)	$\frac{d^2y}{dx^2} = 6x + 2$ <p>$x = \frac{1}{3}, \frac{d^2y}{dx^2} > 0$ $x = -1, \frac{d^2y}{dx^2} < 0$</p>	M1 A1 A1 3	<p>Looks at sign of $\frac{d^2y}{dx^2}$ for at least one of their x-values or other correct method</p> <p>$x = \frac{1}{3}$, minimum point CWO</p> <p>$x = -1$, maximum point CWO</p>
(iii)	$-1 < x < \frac{1}{3}$	M1 A1 2	<p>Any inequality (or inequalities) involving both their x values from part (i)</p> <p>Correct inequality (allow $<$ or \leq)</p>

11

9(i)	Gradient of AB = $\frac{-2-1}{-5-3}$ $= \frac{3}{8}$	B1	$\frac{3}{8}$ oe
	$y-1 = \frac{3}{8}(x-3)$	M1	Equation of line through either A or B, any non-zero numerical gradient
	$8y-8 = 3x-9$		
	$3x-8y-1 = 0$	A1 3	Correct equation in correct form
	(ii)		
	$\left(\frac{-5+3}{2}, \frac{-2+1}{2}\right)$	M1	Uses $\left(\frac{x_1+x_2}{2}, \frac{y_1+y_2}{2}\right)$
	$=(-1, -\frac{1}{2})$	A1 2	$(-1, -\frac{1}{2})$
	(iii)		
	$AC = \sqrt{(-5+3)^2 + (-2-4)^2}$	M1	Uses $\sqrt{(x_2-x_1)^2 + (y_2-y_1)^2}$
	$= \sqrt{2^2 + 6^2}$		
	$= \sqrt{40}$	A1	$\sqrt{40}$
	$= 2\sqrt{10}$	A1 3	Correctly simplified surd
	(iv)		
	Gradient of AC = $\frac{-2-4}{-5+3} = 3$	B1	3 oe
	Gradient of BC = $\frac{4-1}{-3-3} = -\frac{1}{2}$	B1	$-\frac{1}{2}$ oe
	$3 \times -\frac{1}{2} \neq -1$ so lines are not perpendicular	M1	Attempts to check $m_1 \times m_2$
		A1 4	Correct conclusion www
<div style="border: 1px solid black; padding: 2px; display: inline-block;">12</div>			

10(i)	$24x^2 - 3x^{-4}$ $48x + 12x^{-5}$	B1 B1 B1 M1 A1 5	$24x^2$ kx^{-4} $-3x^{-4}$ Attempt to differentiate their (i) Fully correct
(ii)	$8x^3 + \frac{1}{x^3} = -9$ $8x^6 + 1 = -9x^3$ $8x^6 + 9x^3 + 1 = 0$ Let $y = x^3$ $8y^2 + 9y + 1 = 0$ $(8y + 1)(y + 1) = 0$ $y = -\frac{1}{8}, y = -1$ $x = -\frac{1}{2}, x = -1$	*M1 DM1 A1 M1 A1 5 <div style="border: 1px solid black; padding: 2px; display: inline-block;">10</div>	Use a substitution to obtain a 3-term quadratic Correct method to solve quadratic $-\frac{1}{8}, -1$ Attempt to cube root at least one of their y-values $-\frac{1}{2}, -1$ SR one correct x value www B1 SR for trial and improvement: $x = -1$ B1 $x = -\frac{1}{2}$ B2 Justification that there are no further solutions B2