

4724 Core Mathematics 4

<p>1 (a) $2x^2 - 7x - 4 = (2x+1)(x-4)$ or $3x^2 + x - 2 = (3x-2)(x+1)$ $\frac{2x+1}{3x-2}$ as final answer; this answer only</p>	<p>B1 B1 Do not ISW</p>
<p>(b) For correct leading term x in quotient For evidence of correct division process Quotient = $x - 2$ Remainder = $x - 3$</p>	<p>B1 <u>Identity method</u> M1 $M1: x^3 + 2x^2 - 6x - 5 = Q(x^2 + 4x + 1) + R$ A1 $M1: Q = ax + b$ or $x + b, R = cx + d$ & ≥ 2 ops [N.B. If $Q = x + b$, this \Rightarrow 1 of the 2 ops] A1 $A2: a = 1, b = -2, c = 1, d = -3$ SR: <u>B1</u> for two</p>
<p>2 Parts with correct split of $u = \ln x, \frac{dv}{dx} = x^4$ $\frac{x^5}{5} \ln x - \int \frac{x^5}{5} \cdot \frac{1}{x} (dx)$ $\frac{x^5}{5} \ln x - \frac{x^5}{25}$ Correct method with the limits $\frac{4e^5}{25} + \frac{1}{25}$ ISW (Not '+c')</p>	<p>*M1 obtaining result $f(x) + / - \int g(x) dx$ A1 A1 dep*M1 Decimals acceptable here A1 Accept equiv frags; like terms amalgamated</p>
<p>3 (i) $\frac{d}{dx}(x^2 y) = x^2 \frac{dy}{dx} + 2xy$ or $\frac{d}{dx}(xy^2) = 2xy \frac{dy}{dx} + y^2$ Attempt to solve their differentiated equation for $\frac{dy}{dx}$ $\frac{dy}{dx} = \frac{y^2 - 2xy}{x^2 - 2xy}$ only</p>	<p>*B1 dep*M1 A1 WWW AG Must have intermediate line &... ...could imply “=0” on 1st line</p>
<p>(ii)(a) Attempt to solve only $y^2 - 2xy = 0$ & derive $y = 2x$ Clear indication why $y = 0$ is not acceptable</p>	<p>B1 AG Any effort at solving $x^2 - 2xy = 0 \rightarrow B0$ B1 Substituting $y = 2x \rightarrow B0, B0$</p>
<p>(b) Attempt to solve $y = 2x$ simult with $x^2 y - xy^2 = 2$ Produce $-2x^3 = 2$ or $y^3 = -8$ $(-1, -2)$ or $x = -1, y = -2$ only</p>	<p>M1 A1 AEF A1</p>

4	(i)	For (either point) + t (difference between vectors) $\mathbf{r} = (3\mathbf{i} + 2\mathbf{j} + 3\mathbf{k})$ or $\mathbf{i} + 3\mathbf{j} + 4\mathbf{k}) + t(-2\mathbf{i} + \mathbf{j} + \mathbf{k})$ or $2\mathbf{i} - \mathbf{j} - \mathbf{k})$	M1 A1	' t ' can be ' s ', ' λ ' etc. ' \mathbf{r} ' must be ' \mathbf{r} ' but need not be bold Check other formats, e.g. $ta + (1-t)b$
<div>2</div>				
<hr/>				
	(ii)	State/imply that their \mathbf{r} and their $-2\mathbf{i} + \mathbf{j} + \mathbf{k}$ are perpendicular Consider scalar product = 0 Obtain $t = -\frac{1}{6}$ or $\frac{1}{6}$ or $-\frac{5}{6}$ or $\frac{5}{6}$ Subst their t into their equation of AB Obtain $\frac{1}{6}(16\mathbf{i} + 13\mathbf{j} + 19\mathbf{k})$ AEF	*M1 N.B.This *M1 is dep on M1 being earned in (i) dep*M1 A1 M1 A1 Accept decimals if clear	
<div>5</div>				
5	(i)	$(1-x)^{\frac{1}{2}} = 1 - \frac{1}{2}x - \frac{1}{8}x^2$ ignoring x^3 etc $(1+x)^{-\frac{1}{2}} = 1 - \frac{1}{2}x + \frac{3}{8}x^2$ ignoring x^3 etc Product = $1 - x + \frac{1}{2}x^2$ ignoring x^3 etc	B2 B2 B1	SR Allow B1 for $1 - \frac{1}{2}x + kx^2$, $k \neq -\frac{1}{8}$ or 0 SR Allow B1 for $1 - \frac{1}{2}x + kx^2$, $k \neq \frac{3}{8}$ or 0 AG ; with (at least) 1 intermediate step (cf x^2)
<div>5</div>				
<hr/>				
	(ii)	$\sqrt{\frac{5}{9}}$ or $\frac{\sqrt{5}}{3}$ seen $\frac{37}{49}$ or $1 - \frac{2}{7} + \frac{1}{2}\left(\frac{2}{7}\right)^2$ seen $\frac{\sqrt{5}}{3} \approx \frac{37}{49} \Rightarrow \sqrt{5} \approx \frac{111}{49}$	B1 B1 B1 AG	
<div>3</div>				
6	(i)	Produce at least 2 of the 3 relevant equations in t and s Solve for t and s $(t, s) = (4, -3)$ AEF Subst $(4, -3)$ into suitable equation(s) & show consistency	M1 $1 + 2t = 12 + s$, $3t = -4s$, $-5 + 4t = 5 - 2s$ M1 *A1 dep*A1	Either into "3 rd " eqn or into all 3 coordinates. N.B. Intersection coords not asked for
<div>4</div>				
<hr/>				
	(ii)	Method for finding magnitude of any vector Method for finding scalar product of any 2 vectors Using $\cos \theta = \frac{\mathbf{a} \cdot \mathbf{b}}{ \mathbf{a} \mathbf{b} }$ AEF for the correct 2 vectors 137 (136.8359) or 43.2(43.164...)	*M1 Expect $\sqrt{29}$ and $\sqrt{21}$ *M1 Expect -18 dep*M1 Should be $-\frac{18}{\sqrt{29}\sqrt{21}}$ A1 2.39 (2.388236..) or 0.753(0.75335...) rads	
<div>4</div>				

7 (i)	Correct (calc) method for dealing with $\frac{1}{\sin x}$ or $(\sin x)^{-1}$	M1	WWW AG with ≥ 1 line intermed working <div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto; text-align: center; line-height: 20px;">3</div>
	Obtain $-\frac{\cos x}{\sin^2 x}$ or $-(\sin x)^{-2} \cos x$	A1	
	Show manipulation to $-\operatorname{cosec} x \cot x$ (or vice-versa)	A1	
(ii)	Separate variables, $\int (-)\frac{1}{\sin x \tan x} dx = \int \cot t dt$	M1	or $\int \frac{1}{\sin x \tan x} dx = \int (-)\cot t dt$ <u>Style: For the M1 to be awarded, dx and dt must appear on correct sides or there must be \int sign on both sides</u> $\int -\operatorname{cosec} x \cot x dx = \operatorname{cosec} x (+c)$ A1 or $\int \operatorname{cosec} x \cot x dx = -\operatorname{cosec} x$ $\int \cot t dt = \ln \sin t$ or $\ln \sin t $ B1 or $\int -\cot t dt = -\ln \sin t$ or $-\ln \sin t $ Subst $(t, x) = \left(\frac{1}{2}\pi, \frac{1}{6}\pi\right)$ into their equation containing 'c' M1 and attempt to find 'c' $\operatorname{cosec} x = \ln \sin t + 2$ or $\ln \sin t + 2$ A1 WWW ISW ; $\operatorname{cosec} \frac{\pi}{6}$ to be changed to 2 <div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto; text-align: center; line-height: 20px;">5</div>
8 (i)	$A(t+1) + B = 2t$ $A = 2$ $B = -2$	M1 A1 A1	
			<u>Beware</u> : correct values for A and/or B can be obtained from a wrong identity <u>Alt method</u> : subst suitable values into given... ...expressions <div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto; text-align: center; line-height: 20px;">3</div>
(ii)	Attempt to connect dx and dt $dx = t dt$ s.o.i. AEF $x + \sqrt{2x-1} \rightarrow \frac{t^2+1}{2} + t = \frac{(t+1)^2}{2}$ s.o.i. $\int \frac{2t}{(t+1)^2} dt$	M1 A1 B1 A1	But not just $dx = dt$. As AG, look carefully. Any wrong working invalidates AG WWW The 'dt' must be present <div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto; text-align: center; line-height: 20px;">4</div>
(iii)	$\int \frac{1}{t+1} dt = \ln(t+1)$ $\int \frac{1}{(t+1)^2} dt = -\frac{1}{t+1}$ Attempt to change limits (expect 1 & 3) and use $f(t)$ $\ln 4 - \frac{1}{2}$	B1 B1 M1 A1	Or parts $u = 2t, dv = (t+1)^{-2}$ or subst $u = t+1$ or re-substitute and use 1 and 5 on $g(x)$ AEF (like terms amalgamated); if A0 A0 in (i), then final A0 <div style="border: 1px solid black; width: 20px; height: 20px; margin: 0 auto; text-align: center; line-height: 20px;">4</div>

-
- 9 (i) $A : \theta = \frac{1}{2}\pi$ (accept 90°) **B1**
 $B : \theta = 2\pi$ (accept 360°) **B2** SR If B0 awarded for point B, allow B1 SR for any angle s.t. $\sin \theta = 0$

3

- (ii) $\frac{dy}{dx} = \frac{\frac{dy}{d\theta}}{\frac{dx}{d\theta}}$ **M1** or $\frac{dy}{d\theta} \cdot \frac{d\theta}{dx}$ Must be used, not just quoted
 $\frac{dx}{d\theta} = 2 + 2 \cos 2\theta$ **B1**
 $2 + 2 \cos 2\theta = 4 \cos^2 \theta$ with ≥ 1 line intermed work ***B1**
 $\frac{dy}{dx} = \frac{4 \cos \theta}{2 + 2 \cos 2\theta}$ s.o.i. **A1** This & previous line are interchangeable
 $= \sec \theta$ dep***A1** WWW **AG**

5

- (iii) Equating $\sec \theta$ to 2 and producing at least one value of θ **M1** degrees or radians

$$(x =) -\frac{2}{3}\pi - \frac{\sqrt{3}}{2}$$

A1 'Exact' form required

$$(y =) -2\sqrt{3}$$

A1 'Exact' form required

3
