

1	$(4x^2 + 20x + 25) - (x^2 - 6x + 9)$ $= 3x^2 + 26x + 16$ <u>Alternative method using difference of two squares:</u> $(2x + 5 + (x - 3))(2x + 5 - (x - 3))$ $= (3x + 2)(x + 8)$ $= 3x^2 + 26x + 16$	M1 A1 A1 3 3	Square one bracket to give an expression of the form $ax^2 + bx + c$ ($a \neq 0$, $b \neq 0$, $c \neq 0$) One squared bracket fully correct All 3 terms of final answer correct M1 2 brackets with same terms but different signs A1 One bracket correctly simplified A1 All 3 terms of final answer correct
2 (a)(i)		B1	Excellent curve for $\frac{1}{x}$ in either quadrant
(ii)		B1 2	Excellent curve for $\frac{1}{x}$ in other quadrant
		SR B1	Reasonably correct curves in 1 st and 3 rd quadrants
(b)	Stretch Scale factor 8 in y direction or scale factor $\frac{1}{2}$ in x direction	B1 B1 2 5	Correct graph, minimum point at origin, symmetrical
3 (i)	$3\sqrt{20}$ or $3\sqrt{2} \sqrt{5} \times \sqrt{2}$ or $\sqrt{180}$ or $\sqrt{90} \times \sqrt{2}$ $= 6\sqrt{5}$	M1 A1 2	Correctly simplified answer
(ii)	$10\sqrt{5} + 5\sqrt{5}$ $= 15\sqrt{5}$	M1 B1 A1 3 5	Attempt to change both surds to $\sqrt{5}$ One part correct and fully simplified cao

4 (i)	$(-4)^2 - 4 \times k \times k$ $= 16 - 4k^2$	M1 A1 2	Uses $b^2 - 4ac$ (involving k) $16 - 4k^2$
(ii)	$16 - 4k^2 = 0$ $k^2 = 4$ $k = 2$ or $k = -2$	M1 B1 B1 3 5	Attempts $b^2 - 4ac = 0$ (involving k) or attempts to complete square (involving k)
5 (i)	Length = $20 - 2x$ Area = $x(20 - 2x)$ $= 20x - 2x^2$	M1 A1 2	Expression for length of enclosure in terms of x Correctly shows that area = $20x - 2x^2$ AG
(ii)	$\frac{dA}{dx} = 20 - 4x$ For max, $20 - 4x = 0$ $x = 5$ only Area = 50	M1 M1 A1 A1 4 6	Differentiates area expression Uses $\frac{dy}{dx} = 0$
6	Let $y = (x + 2)^2$ $y^2 + 5y - 6 = 0$ $(y + 6)(y - 1) = 0$ $y = -6$ or $y = 1$ $(x + 2)^2 = 1$ $x = -1$ or $x = -3$	B1 M1 A1 M1 A1 A1 6 6	Substitute for $(x + 2)^2$ to get $y^2 + 5y - 6 (= 0)$ Correct method to find roots Both values for y correct Attempt to work out x One correct value Second correct value and no extra real values
7 (a)	$f(x) = x + 3x^{-1}$ $f'(x) = 1 - 3x^{-2}$	M1 A1 A1 A1 4	Attempt to differentiate First term correct x^{-2} soi www Fully correct answer
(b)	$\frac{dy}{dx} = \frac{5}{2}x^{\frac{3}{2}}$ When $x = 4$, $\frac{dy}{dx} = \frac{5}{2}\sqrt{4^{\frac{3}{2}}}$ $= 20$	M1 B1 B1 M1 A1 5 9	Use of differentiation to find gradient $\frac{5}{2}x^{\frac{3}{2}}$ $kx^{\frac{3}{2}}$ $\sqrt{4^{\frac{3}{2}}}$ soi SR If 0 scored for first 3 marks, award B1 if $\sqrt{4^{\frac{3}{2}}}$ correctly evaluated.

8 (i)	$(x + 4)^2 - 16 + 15$ $= (x + 4)^2 - 1$	B1 M1 A1 3	$a = 4$ 15 – their a^2 cao in required form
(ii)	(-4, -1)	B1 ft B1 ft 2 M1 A1	Correct x coordinate Correct y coordinate Correct method to find roots -5, -3
(iii)	$x^2 + 8x + 15 > 0$ $(x + 5)(x + 3) > 0$ $x < -5, x > -3$	M1 A1 4 9	Correct method to solve quadratic inequality eg +ve quadratic graph $x < -5, x > -3$ (not wrapped, strict inequalities, no 'and')
9 (i)	$(x - 3)^2 - 9 + y^2 - k = 0$ $(x - 3)^2 + y^2 = 9 + k$ Centre (3, 0) $9 + k = 4^2$ $k = 7$	B1 B1 M1 A1 4	$(x - 3)^2$ soi Correct centre Correct value for k (may be embedded)
			<u>Alternative method using expanded form:</u> Centre (-g, -f) M1 Centre (3, 0) A1 $4 = \sqrt{f^2 + g^2 - (-k)}$ M1 $k = 7$ A1
(ii)	$(3 - 3)^2 + y^2 = 16$ $y^2 = 16$ $y = 4$ Length of AB = $\sqrt{(-1 - 3)^2 + (0 - 4)^2}$ $= \sqrt{32}$ $= 4\sqrt{2}$	M1 A1 M1 A1 ft A1 5	Attempt to substitute $x = 3$ into original equation or their equation $y = 4$ (do not allow ± 4) Correct method to find line length using Pythagoras' theorem $\sqrt{32}$ or $\sqrt{16 + a^2}$ cao
(iii)	Gradient of AB = 1 or $\frac{a}{4}$ $y - 0 = m(x + 1)$ or $y - 4 = m(x - 3)$ $y = x + 1$	B1 ft M1 A1 3 12	Attempts equation of straight line through their A or B with their gradient Correct equation in any form with simplified constants

10 (i)	$(3x + 1)(x - 5) = 0$ $x = \frac{-1}{3}$ or $x = 5$	M1 A1 A1 3	Correct method to find roots Correct brackets or formula Both values correct SR B1 for $x = 5$ spotted www
(ii)		B1 B1 B1 ft 3	Positive quadratic (must be reasonably symmetrical) y intercept correct both x intercepts correct
(iii)	$\frac{dy}{dx} = 6x - 14$ $6x - 14 = 4$ $x = 3$ On curve, when $x = 3$, $y = -20$ $-20 = (4 \times 3) + c$ $c = -32$ <u>Alternative method:</u> $3x^2 - 14x - 5 = 4x + c$ $3x^2 - 18x - 5 - c = 0$ has one solution $b^2 - 4ac = 0$ $(-18)^2 - (4 \times 3 \times (-5 - c)) = 0$ $c = -32$	M1* M1* A1 A1 ft M1dep A1 6 M1 B1 M1 M1 A1 A1	Use of differentiation to find gradient of curve Equating their gradient expression to 4 Finding y co ordinate for their x value N.B. dependent on both previous M marks Equate curve and line (or substitute for x) Statement that only one solution for a tangent (may be implied by next line) Use of discriminant = 0 Attempt to use a, b, c from their equation Correct equation $c = -32$

12